



Cayley Graphs and Semi-Direct Products

Tushar Muralidharan

Australian National University

May 2024



Table of Contents

1 Preliminaries

- Introduction
- Groups as Symmetries
- Construction of Symmetry Groups

2 Cayley Graphs

- Definitions
- Drawing Groups
- The Relationship Between Groups and Cayley Graphs

3 Semi-Direct Products

- Group Products
- Construction

4 Applications

- Dehn's Three Decision Problems
- Conclusion


What is Group Theory?

group theory 

The study of the symmetries that define the properties of a system.

Figure: Oxford definition of group theory [7]





Groups

A group is a set G together with a binary operation “ \cdot ” on G that satisfies the following three properties:

- *Associativity.* For all $a, b, c \in G$ one has $(a \cdot b) \cdot c = a \cdot (b \cdot c)$.
- *Identity.* There exists an element $e \in G$ such that for all $a \in G$ one has $e \cdot a = a$ and $a \cdot e = a$.
- *Inverses.* For each $a \in G$, there exists an element $b \in G$ such that $a \cdot b = e$ and $b \cdot a = e$, where e is the identity element.



The Symmetric Group

Example

The *symmetric group* S_n is defined as the set of permutations of the set $\{1, 2, 3, \dots, n\}$.

Alternatively, it is the set of symmetries of n points.





The Dihedral Group

Example

The *dihedral group* D_n is **defined as** the set of symmetries of the regular n -gon.

For example, the group D_8 consists of the 8 rotations and 8 reflections of the regular octagon.





Reverse-Engineering


Question: For what object is $\mathbb{Z}/n\mathbb{Z}$ the group of symmetries?



Reverse-Engineering

Question: For what object is $\mathbb{Z}/n\mathbb{Z}$ the group of symmetries?

Guess:






Reverse-Engineering

Question: For what object is $\mathbb{Z}/n\mathbb{Z}$ the group of symmetries?

Guess:




If we choose an n -gon, we get D_n .



Reverse-Engineering

Question: For what object is $\mathbb{Z}/n\mathbb{Z}$ the group of symmetries?

Guess:




If we choose an n -gon, we get D_n . But, we can identify $\mathbb{Z}/n\mathbb{Z}$ as the subgroup of rotational symmetries of D_n .



Reverse-Engineering

Question: For what object is $\mathbb{Z}/n\mathbb{Z}$ the group of symmetries?

Guess:



If we choose an n -gon, we get D_n . But, we can identify $\mathbb{Z}/n\mathbb{Z}$ as the subgroup of rotational symmetries of D_n .


Idea: Draw a little arrow in the middle of each edge.





Reverse-Engineering


Question: For what object is $\mathbb{Z}/n\mathbb{Z}$ the group of symmetries?





Reverse-Engineering

Question: For what object is $\mathbb{Z}/n\mathbb{Z}$ the group of symmetries?



Symmetries must now preserve the arrows.



More Reverse-Engineering

Question: For what object is \mathbb{Z} the group of symmetries?

Guess:



But...



More Reverse-Engineering

Question: For what object is \mathbb{Z} the group of symmetries?

Guess:



But... this doesn't work. Although \mathbb{Z} corresponds to translations left or right by n units, we also have reflections about integer or half-integer points are also symmetries of this line.



More Reverse-Engineering

Question: For what object is \mathbb{Z} the group of symmetries?

Guess:




But... this doesn't work. Although \mathbb{Z} corresponds to translations left or right by n units, we also have reflections about integer or half-integer points are also symmetries of this line.

Idea: Draw a little arrow in the middle of each edge.



More Reverse-Engineering

Question: For what object is \mathbb{Z} the group of symmetries?



More Reverse-Engineering

Question: For what object is \mathbb{Z} the group of symmetries?



Symmetries must now preserve the arrows.




A Different Technique

Question: For what object is $\mathbb{Z} \times \mathbb{Z}$ the group of symmetries?



A Different Technique

Question: For what object is $\mathbb{Z} \times \mathbb{Z}$ the group of symmetries?
Using our previous trick:




But...



A Different Technique

Question: For what object is $\mathbb{Z} \times \mathbb{Z}$ the group of symmetries?
Using our previous trick:




But...this doesn't work. Although every $(a, b) \in \mathbb{Z} \times \mathbb{Z}$ corresponds to a translation (left or right) by a units and (up and down) by b units, we also have reflectional symmetries.



A Different Technique

Question: For what object is $\mathbb{Z} \times \mathbb{Z}$ the group of symmetries?
Using our previous trick:




But...this doesn't work. Although every $(a, b) \in \mathbb{Z} \times \mathbb{Z}$ corresponds to a translation (left or right) by a units and (up and down) by b units, we also have reflectional symmetries. **Idea:** Colors!



A Different Technique

Question: For what object is $\mathbb{Z} \times \mathbb{Z}$ the group of symmetries?



Symmetries must now preserve the arrows *and* the colors.



Where From Here?

I hope to convince you that every finitely presented group is **exactly** the symmetries of some geometric object, namely, its **Cayley graph**.

This should make some sense of the definition of group theory as the study of symmetries.

Let's modify our earlier techniques. (Foreshadowing.)

Drawing arrows \rightarrow Adding directions to edges of a graph

Adding colors \rightarrow Adding labels to edges of a graph



Table of Contents

1 Preliminaries

- Introduction
- Groups as Symmetries
- Construction of Symmetry Groups

2 Cayley Graphs


- Definitions
- Drawing Groups
- The Relationship Between Groups and Cayley Graphs

3 Semi-Direct Products

- Group Products
- Construction

4 Applications


- Dehn's Three Decision Problems
- Conclusion



What is a Cayley Graph?

Definition

The *Cayley graph* for a group G with respect to a generating set S is a directed, labeled graph whose *vertices correspond to the elements* of G , and for each $g \in G$ and $s \in S$ contains an edge from g to gs labeled by s .




Note: The definition of the Cayley graph depends on the generating set S chosen for G . More on this later.

What is a Cayley Graph?

Definition

The *Cayley graph* for a group G with respect to a generating set S is a directed, labeled graph whose *vertices correspond to the elements of G* , and for each $g \in G$ and $s \in S$ contains an edge from g to gs labeled by s .




Sometimes, we color-code (or vary the patterns of) the edges by their generator instead of labeling them. Also, if there is an edge with arrows on both sides, we sometimes omit the arrowheads.




A First Example

The Cayley graph for $C_5 = \langle a \rangle = \{e, a, a^2, a^3, a^4\}$ with respect to the generating set $\{a\}$ is




A First Example

The Cayley graph for $C_5 = \langle a \rangle = \{e, a, a^2, a^3, a^4\}$ with respect to the generating set $\{a\}$ is



A First Example

The Cayley graph for $C_5 = \langle a \rangle = \{e, a, a^2, a^3, a^4\}$ with respect to the generating set $\{a\}$ is




After removing the vertex labels, what are the symmetries (preserving labels and directions) of this graph?




More Cayley Graphs

Can you tell which generators correspond to which edge patterns?



S_3 with respect to $\{(12), (123)\}$




D_4 with respect to a rotation and a reflection


What happens if you look at the graph of the dihedral group in 3D?




Dihedral Groups in 3D



D_4 with respect to a rotation and a reflection



D_5 with respect to a rotation and a reflection



D_6 with respect to a rotation and a reflection

Can you see why these groups might be called “dihedral?”

A More Complicated Cayley Graph




Figure: The Cayley graph of the symmetry group of a cube (generated by three reflections).



Cayley's Theorem

Let $\text{Cay}(G, S)$ denote the Cayley graph of a group G with respect to the generating set S .



Cayley's Theorem

Let $\text{Cay}(G, S)$ denote the Cayley graph of a group G with respect to the generating set S .

Theorem

Every group G with finite generating set S is isomorphic to the (direction, label)-preserving symmetries of $\text{Cay}(G, S)$.




Cayley's Theorem

Theorem

Every group G with finite generating set S is isomorphic to the (direction, label)-preserving symmetries of $\text{Cay}(G, S)$.

Proof Sketch



Cayley's Theorem


Theorem


Every group G with finite generating set S is isomorphic to the (direction, label)-preserving symmetries of $\text{Cay}(G, S)$.

Proof Sketch

Every element $g \in G$ determines a symmetry γ_g of $\text{Cay}(G, S)$,

which sends $\begin{matrix} h \\ \bullet \end{matrix} \rightsquigarrow \begin{matrix} gh \\ \bullet \end{matrix}$ and





Cayley's Theorem

Theorem

Every group G with finite generating set S is isomorphic to the (direction, label)-preserving symmetries of $\text{Cay}(G, S)$.

Proof Sketch

Now we show that every symmetry of $\text{Cay}(G, S)$ arises in this way.

Claim: If ρ is a symmetry of $\text{Cay}(G, S)$, then $\rho = \gamma_g$ for the element g such that $\rho(\bullet_e) = \bullet_g$.

Consider the symmetry $\rho \cdot \gamma_g^{-1}$. This takes \bullet_e to \bullet_e and edges incident to e (since it preserves directions and labels). Hence $\rho \cdot \gamma_g^{-1}$ fixes all vertices adjacent to \bullet_e . But then it will fix edges incident to these, and so on. Thus $\rho \cdot \gamma_g^{-1} = \text{id}$, i.e., $\rho = \gamma_g$. □



Uniqueness

Question: Are Cayley graphs unique?



Uniqueness

Question: Are Cayley graphs unique?


Answer: No, $\text{Cay}(G, S)$ depends on the generating set S .




Uniqueness

Question: Are Cayley graphs unique?

Answer: No, $\text{Cay}(G, S)$ depends on the generating set S .



D_4 with respect to a rotation and reflection




D_4 with respect to two adjacent reflections

But...


Uniqueness

Question: Are Cayley graphs unique?

Answer: No, $\text{Cay}(G, S)$ depends on the generating set S .



D_4 with respect to a rotation and reflection



D_4 with respect to two adjacent reflections

But... Cayley graphs *are* unique up to **quasi-isometry**(!)




Table of Contents

1 Preliminaries

- Introduction
- Groups as Symmetries
- Construction of Symmetry Groups

2 Cayley Graphs

- Definitions
- Drawing Groups
- The Relationship Between Groups and Cayley Graphs

3 Semi-Direct Products

- Group Products
- Construction

4 Applications

- Dehn's Three Decision Problems
- Conclusion




Direct Products


Given groups (G, \star) and (H, \triangle) , recall the notion of the **direct product** $G \times H$, whose binary operation is defined component-wise:
 $(g_1, h_1) \cdot (g_2, h_2) = (g_1 \star g_2, h_1 \triangle h_2)$.

What does the Cayley graph of $G \times H$ look like? Let's look at the case where G and H are cyclic.

Direct Products




Cayley graph of $C_2 \times C_6$




Cayley graph of $C_3 \times C_6$

Direct Products




Cayley graph of $C_2 \times C_6$



Cayley graph of $C_3 \times C_6$

Let $C_2 = \langle a \rangle$ and $C_6 = \langle b \rangle$. Then multiplying by the generator (a, e) takes us along the white lines and multiplying by (e, b) takes us around the hexagon.





Generalizing the Direct Product


Let's compare the Cayley graph of $C_2 \times C_6$ with that of D_6 .

Generalizing the Direct Product


Let's compare the Cayley graph of $C_2 \times C_6$ with that of D_6 .


 $C_2 \times C_6$

 D_6

Generalizing the Direct Product




$$C_2 \times C_6$$




$$D_6$$

They are the same, except that we traverse the back hexagon in the opposite direction.

Generalizing the Direct Product



$$C_2 \times C_6$$



$$D_6$$

Recall that in $C_2 \times C_6$, moving around the hexagon corresponds to multiplying by (e, h) for some h .



Generalizing the Direct Product

That is, we have

$$(e, h) \cdot (a, b) = (a, hb),$$

so the first component is fixed and the second is multiplied by h .




Generalizing the Direct Product

That is, we have

$$(e, h) \cdot (a, b) = (a, hb),$$

so the first component is fixed and the second is multiplied by h .
What if we relax the condition that we must stay at the same first component?



Generalizing the Direct Product

That is, we have

$$(e, h) \cdot (a, b) = (a, hb),$$

so the first component is fixed and the second is multiplied by h .
What if we relax the condition that we must stay at the same first component? In the general case for two groups G and H , we write

$$(e, h_1) \cdot (g_2, h_2) = (f_{h_1}(g_2), h_1 h_2),$$

where $f_{h_1}(g_2)$ need not be equal to g_2 .



Generalizing the Direct Product

That is, we have

$$(e, h) \cdot (a, b) = (a, hb),$$

so the first component is fixed and the second is multiplied by h . What if we relax the condition that we must stay at the same first component? In the general case for two groups G and H , we write

$$(e, h_1) \cdot (g_2, h_2) = (f_{h_1}(g_2), h_1 h_2),$$

where $f_{h_1}(g_2)$ need not be equal to g_2 . As we vary g_2 , we might expect $f_{h_1}(g_2)$ to vary across all elements of G , so we enforce that f_{h_1} is a **bijection** for all $h_1 \in H$.



Generalizing the Direct Product

Let's denote our new group by $G \rtimes H$. Since we want our multiplication to be well-defined, we must have

$$\begin{aligned}
 (f_{h_1}(g_1) \cdot f_{h_1}(g_2), h_1 h_2) &= (f_{h_1}(g_1), e) \cdot (f_{h_1}(g_2), h_1 h_2) \\
 &= (e, h_1) \cdot (g_1, h_1^{-1}) \cdot (e, h_1) \cdot (g_2, h_2) \\
 &= (e, h_1) \cdot (g_1, e) \cdot (g_2, h_2) \\
 &= (e, h_1) \cdot (g_1 g_2, h_2) \\
 &= (f_{h_1}(g_1 g_2), h_1 h_2),
 \end{aligned}$$

so $f_{h_1}(g_1) \cdot f_{h_1}(g_2) = f_{h_1}(g_1 g_2)$, i.e. f_{h_1} must be a **homomorphism**.



Generalizing the Direct Product

Let's denote our new group by $G \rtimes H$. Since we want our multiplication to be well-defined, we must have

$$\begin{aligned}
 (f_{h_1}(g_1) \cdot f_{h_1}(g_2), h_1 h_2) &= (f_{h_1}(g_1), e) \cdot (f_{h_1}(g_2), h_1 h_2) \\
 &= (e, h_1) \cdot (g_1, h_1^{-1}) \cdot (e, h_1) \cdot (g_2, h_2) \\
 &= (e, h_1) \cdot (g_1, e) \cdot (g_2, h_2) \\
 &= (e, h_1) \cdot (g_1 g_2, h_2) \\
 &= (f_{h_1}(g_1 g_2), h_1 h_2),
 \end{aligned}$$

so $f_{h_1}(g_1) \cdot f_{h_1}(g_2) = f_{h_1}(g_1 g_2)$, i.e. f_{h_1} must be a **homomorphism**. Since it is also a bijection, f_{h_1} must be an **automorphism** for all $h_1 \in H$.



Generalizing the Direct Product

Hence we have a map $\varphi: H \rightarrow \text{Aut}(G)$ defined by $h \mapsto f_h$.
Moreover,



Generalizing the Direct Product

Hence we have a map $\varphi: H \rightarrow \text{Aut}(G)$ defined by $h \mapsto f_h$.
Moreover,

$$\begin{aligned}
 (f_{h_1 h_2}(g_1), h_1 h_2) &= (e, h_1 h_2) \cdot (g_1, e) \\
 &= (e, h_1) \cdot (e, h_2) \cdot (g_1, e) \\
 &= (e, h_1) \cdot (f_{h_2}(g_1), h_2) \\
 &= (f_{h_1}(f_{h_2}(g_1)), h_1 h_2) \\
 &= ((f_{h_1} \circ f_{h_2})(g_1), h_1 h_2)
 \end{aligned}$$

$$\text{so } f_{h_1 h_2}(g_1) = (f_{h_1} \circ f_{h_2})(g_1).$$




Generalizing the Direct Product

Hence we have a map $\varphi: H \rightarrow \text{Aut}(G)$ defined by $h \mapsto f_h$.
Moreover,

$$\begin{aligned}
 (f_{h_1 h_2}(g_1), h_1 h_2) &= (e, h_1 h_2) \cdot (g_1, e) \\
 &= (e, h_1) \cdot (e, h_2) \cdot (g_1, e) \\
 &= (e, h_1) \cdot (f_{h_2}(g_1), h_2) \\
 &= (f_{h_1}(f_{h_2}(g_1)), h_1 h_2) \\
 &= ((f_{h_1} \circ f_{h_2})(g_1), h_1 h_2)
 \end{aligned}$$

so $f_{h_1 h_2}(g_1) = (f_{h_1} \circ f_{h_2})(g_1)$. In other words, the map φ is a homomorphism.



Generalizing the Direct Product

Finally, we derive the group operation for $G \rtimes H$.

$$\begin{aligned}
 (g_1, h_1) \cdot (g_2, h_2) &= (g_1, e) \cdot (e, h_1) \cdot (g_2, h_2) \\
 &= (g_1, e) \cdot (f_{h_1}(g_2), h_1 h_2) \\
 &= (g_1 f_{h_1}(g_2), h_1 h_2) \\
 &= (g_1 \varphi(h_1)(g_2), h_1 h_2).
 \end{aligned}$$

Therefore, we arrive at the definition of a semi-direct product of groups.



The Definition


Definition

The *semi-direct product* of two groups G and H with respect to a homomorphism $\varphi: H \rightarrow \text{Aut}(G)$, denoted $G \rtimes_{\varphi} H$, is the set $G \times H$ with the binary operation defined by


$$(g_1, h_1) \cdot (g_2, h_2) = (g_1 \varphi(h_1)(g_2), h_1 h_2).$$

If we choose φ to be the trivial homomorphism, then the semi-direct product reduces to the direct product.

Our Original Example




$$C_2 \times C_6 \cong C_6 \times C_2$$




$$D_6$$




A Twist



Our Original Example




$$C_2 \times C_6 \cong C_6 \times C_2$$




$$D_6$$



Our Original Example




$$C_2 \times C_6 \cong C_6 \times C_2$$




$$D_6$$

If we define $\varphi: C_2 \rightarrow \text{Aut}(C_6) \cong C_2$ to be the nontrivial homomorphism, then $C_6 \rtimes_{\varphi} C_2 \cong D_6$!

Our Original Example



$$C_2 \times C_6 \cong C_6 \times C_2$$



$$D_6$$

If we define $\varphi: C_2 \rightarrow \text{Aut}(C_6) \cong C_2$ to be the nontrivial homomorphism, then $C_6 \rtimes_{\varphi} C_2 \cong D_6$! In this case, φ acts like a **twist** that reverses the directions of the arrows on one of the hexagons.



Table of Contents

1 Preliminaries

- Introduction
- Groups as Symmetries
- Construction of Symmetry Groups

2 Cayley Graphs

- Definitions
- Drawing Groups
- The Relationship Between Groups and Cayley Graphs

3 Semi-Direct Products

- Group Products
- Construction

4 Applications

- Dehn's Three Decision Problems
- Conclusion



Why Should We Care?

Are Cayley graphs just for show?




Why Should We Care?

Are Cayley graphs just for show? **No.**



Decision Problems

In 1912, German mathematician Max Dehn posed the following questions.




Decision Problems

In 1912, German mathematician Max Dehn posed the following questions.

- *Word Problem.* Is it possible to decide which words (i.e. finite products of the generators and their inverses) represent the identity element?
- *Conjugacy Problem.* Is it possible to decide whether two elements are conjugate?
- *Isomorphism Problem.* Is it possible to decide whether two groups are isomorphic?

All three problems are...



Decision Problems

In 1912, German mathematician Max Dehn posed the following questions.

- *Word Problem.* Is it possible to decide which words (i.e. finite products of the generators and their inverses) represent the identity element?
- *Conjugacy Problem.* Is it possible to decide whether two elements are conjugate?
- *Isomorphism Problem.* Is it possible to decide whether two groups are isomorphic?

All three problems are... undecidable, in general. However...



The Decision Problems

Cayley graphs can help. Much of the efforts of geometric group theorists has been to solve the word problem for certain classes of groups.



The Decision Problems

Cayley graphs can help. Much of the efforts of geometric group theorists has been to solve the word problem for certain classes of groups.

If a group G is generated by the set S , then every word w in the generators determines a path in the Cayley graph from \bullet_e to \bullet_w . Conversely, every edge path in the Cayley graph determines a word (where we add inverses if the path travels in the opposite direction of an edge).

Words that represent
the identity element



Closed edge paths in
the Cayley graph



The Decision Problems

Cayley graphs can help. Much of the efforts of geometric group theorists has been to solve the word problem for certain classes of groups.

If a group G is generated by the set S , then every word w in the generators determines a path in the Cayley graph from \bullet_e to \bullet_w . Conversely, every edge path in the Cayley graph determines a word (where we add inverses if the path travels in the opposite direction of an edge).

Words that represent
the identity element




Closed edge paths in
the Cayley graph

The latter can be analyzed via geometry and topology!



Group-Theoretic and Geometric Connections


The latter can be analyzed via geometry and topology!



Group-Theoretic and Geometric Connections

The latter can be analyzed via geometry and topology!

The shortest path metric on the Cayley graph is called the *word metric*. Treating Cayley graphs (1-complexes) as geometric objects can illuminate geometric properties that are useful in solving the word problem. For example, short-cuts in Cayley graphs in hyperbolic space can be shown to correspond to length-reducing applications of group relations [3].



Group-Theoretic and Geometric Connections

The latter can be analyzed via geometry and topology!

The shortest path metric on the Cayley graph is called the *word metric*. Treating Cayley graphs (1-complexes) as geometric objects can illuminate geometric properties that are useful in solving the word problem. For example, short-cuts in Cayley graphs in hyperbolic space can be shown to correspond to length-reducing applications of group relations [3].

My research involves Cayley graphs embedded in $CAT(0)$ spaces (complexes), which are defined by non-positive curvature.



A Parting Reflection

So, my interest in symmetry has not been misplaced.

*H. S. M. Coxeter,
upon learning that his brain displayed
a high degree of bilateral symmetry*

References I



Matt Clay and Dan Margalit.
Office Hours with a Geometric Group Theorist.
 Princeton University Press, 2017.



M. Dehn.
 Über unendliche diskontinuierliche gruppen.
Mathematische Annalen, 71(1):116–144, Mar 1911.



M. Gromov.
Hyperbolic Groups, pages 75–263.
 Springer New York, New York, NY, 1987.

References II



John Meier.

Groups, Graphs and Trees: An Introduction to the Geometry of Infinite Groups.

London Mathematical Society Student Texts. Cambridge University Press, 2008.



Jules Poon.

Cayley graphs.



Jules Poon.

Cayley graphs and pretty things.



References III



Richard Rennie and Jonathan Law.

A Dictionary of Physics.

Oxford University Press, 2019.