

CLASS 10 · ICSE · SELINA CONCISE MATHEMATICS · PYQ SOLUTIONS

Chapter 21: Trigonometrical Identities

Step-by-step solutions to the previous year questions

The question numbers match the PYQ question sheet.

1 The three basic identities

1. Assertion (A): If $\sin^2 A + \sin A = 1$, then $\cos^4 A + \cos^2 A = 1$. Reason (R): $1 - \sin^2 A = \cos^2 A$. (a) A true, R false (b) A false, R true (c) both true, and R is the correct reason for A (d) both true, but R is not the correct reason for A **ICSE 2025, 1 mark**

SOLUTION

From the given relation, $\sin A = 1 - \sin^2 A = \cos^2 A$ (using R).

Then $\cos^4 A + \cos^2 A = \sin^2 A + \sin A = 1$. So A is true, and R is exactly what proves it.

Answer: (c)

2. Statement 1: $\sin^2 \theta + \cos^2 \theta = 1$. Statement 2: $\operatorname{cosec}^2 \theta + \cot^2 \theta = 1$. Which is valid? (a) only 1 (b) only 2 (c) both (d) neither **ICSE 2024, 1 mark**

SOLUTION

Statement 1 is a basic identity. The correct form of the second is $\operatorname{cosec}^2 \theta - \cot^2 \theta = 1$, so statement 2 is not valid.

Answer: (a)

3. $(1 + \sin A)(1 - \sin A)$ equals: (a) $\operatorname{cosec}^2 A$ (b) $\sin^2 A$ (c) $\sec^2 A$ (d) $\cos^2 A$ **ICSE 2023, 1 mark**

SOLUTION

$$(1 + \sin A)(1 - \sin A) = 1 - \sin^2 A = \cos^2 A$$

Answer: (d)

2 Proving identities

4. Prove that $\frac{(\cot A + \tan A - 1)(\sin A + \cos A)}{\sin^3 A + \cos^3 A} = \sec A \operatorname{cosec} A$. **ICSE 2025, 4 marks**

SOLUTION

$$\begin{aligned}\cot A + \tan A - 1 &= \frac{\cos^2 A + \sin^2 A - \sin A \cos A}{\sin A \cos A} = \frac{1 - \sin A \cos A}{\sin A \cos A} \\ \sin^3 A + \cos^3 A &= (\sin A + \cos A)(\sin^2 A - \sin A \cos A + \cos^2 A) = (\sin A + \cos A)(1 - \sin A \cos A) \\ \text{LHS} &= \frac{(1 - \sin A \cos A)(\sin A + \cos A)}{\sin A \cos A (\sin A + \cos A)(1 - \sin A \cos A)} = \frac{1}{\sin A \cos A} \\ &= \sec A \operatorname{cosec} A = \text{RHS. Proved.}\end{aligned}$$

5. Factorise $\sin^3 \theta + \cos^3 \theta$. Hence prove that $\frac{\sin^3 \theta + \cos^3 \theta}{\sin \theta + \cos \theta} + \sin \theta \cos \theta = 1$. **ICSE 2024, 4 marks**

SOLUTION

$$\begin{aligned}\text{Using } a^3 + b^3 &= (a + b)(a^2 - ab + b^2): \sin^3 \theta + \cos^3 \theta = (\sin \theta + \cos \theta)(\sin^2 \theta - \sin \theta \cos \theta + \cos^2 \theta) \\ &= (\sin \theta + \cos \theta)(1 - \sin \theta \cos \theta) \\ \text{LHS} &= \frac{(\sin \theta + \cos \theta)(1 - \sin \theta \cos \theta)}{\sin \theta + \cos \theta} + \sin \theta \cos \theta = 1 - \sin \theta \cos \theta + \sin \theta \cos \theta \\ &= 1 = \text{RHS. Proved.}\end{aligned}$$

6. Prove that $(\sin^2 \theta - 1)(\tan^2 \theta + 1) + 1 = 0$. **ICSE 2023, 4 marks**

SOLUTION

$$\begin{aligned}\sin^2 \theta - 1 &= -\cos^2 \theta \text{ and } \tan^2 \theta + 1 = \sec^2 \theta \\ \text{LHS} &= -\cos^2 \theta \times \sec^2 \theta + 1 = -1 + 1 \\ &= 0 = \text{RHS. Proved.}\end{aligned}$$

7. Prove that $\frac{\sin A}{1 + \cot A} - \frac{\cos A}{1 + \tan A} = \sin A - \cos A$. **ICSE 2020, 3 marks**

SOLUTION

$$\begin{aligned}\frac{\sin A}{1 + \cot A} &= \frac{\sin A}{\frac{\sin A + \cos A}{\sin A}} = \frac{\sin^2 A}{\sin A + \cos A} \text{ and } \frac{\cos A}{1 + \tan A} = \frac{\cos^2 A}{\sin A + \cos A} \\ \text{LHS} &= \frac{\sin^2 A - \cos^2 A}{\sin A + \cos A} = \frac{(\sin A - \cos A)(\sin A + \cos A)}{\sin A + \cos A} \\ &= \sin A - \cos A = \text{RHS. Proved.}\end{aligned}$$

8. Prove that $\left(\frac{1 - \tan \theta}{1 - \cot \theta}\right)^2 = \tan^2 \theta$. **ICSE 2020, 4 marks**

SOLUTION

$$1 - \cot \theta = 1 - \frac{1}{\tan \theta} = \frac{\tan \theta - 1}{\tan \theta}$$

$$\frac{1 - \tan \theta}{1 - \cot \theta} = \frac{(1 - \tan \theta) \tan \theta}{\tan \theta - 1} = -\tan \theta$$

Squaring: LHS = $(-\tan \theta)^2 = \tan^2 \theta = \text{RHS}$. **Proved.**

9. Prove that $(\operatorname{cosec} \theta - \sin \theta)(\sec \theta - \cos \theta)(\tan \theta + \cot \theta) = 1$. **ICSE 2019, 3 marks**

SOLUTION

$$\operatorname{cosec} \theta - \sin \theta = \frac{1 - \sin^2 \theta}{\sin \theta} = \frac{\cos^2 \theta}{\sin \theta} \text{ and } \sec \theta - \cos \theta = \frac{\sin^2 \theta}{\cos \theta}$$

$$\tan \theta + \cot \theta = \frac{\sin^2 \theta + \cos^2 \theta}{\sin \theta \cos \theta} = \frac{1}{\sin \theta \cos \theta}$$

$$\text{LHS} = \frac{\cos^2 \theta}{\sin \theta} \times \frac{\sin^2 \theta}{\cos \theta} \times \frac{1}{\sin \theta \cos \theta} = 1 = \text{RHS. Proved.}$$

10. Prove that $\sqrt{\sec^2 \theta + \operatorname{cosec}^2 \theta} = \tan \theta + \cot \theta$. **ICSE 2018, 3 marks**

SOLUTION

$$\sec^2 \theta + \operatorname{cosec}^2 \theta = \frac{1}{\cos^2 \theta} + \frac{1}{\sin^2 \theta} = \frac{\sin^2 \theta + \cos^2 \theta}{\sin^2 \theta \cos^2 \theta} = \frac{1}{\sin^2 \theta \cos^2 \theta}$$

$$\text{LHS} = \frac{1}{\sin \theta \cos \theta}$$

$$\text{RHS} = \frac{\sin \theta}{\cos \theta} + \frac{\cos \theta}{\sin \theta} = \frac{\sin^2 \theta + \cos^2 \theta}{\sin \theta \cos \theta} = \frac{1}{\sin \theta \cos \theta}$$

LHS = RHS. **Proved.**

11. Prove that $(1 + \cot \theta - \operatorname{cosec} \theta)(1 + \tan \theta + \sec \theta) = 2$. **ICSE 2018, 4 marks**

SOLUTION

$$\begin{aligned} \text{LHS} &= \left(\frac{\sin \theta + \cos \theta - 1}{\sin \theta}\right) \left(\frac{\cos \theta + \sin \theta + 1}{\cos \theta}\right) = \frac{(\sin \theta + \cos \theta)^2 - 1}{\sin \theta \cos \theta} \\ &= \frac{1 + 2 \sin \theta \cos \theta - 1}{\sin \theta \cos \theta} = 2 = \text{RHS. Proved.} \end{aligned}$$

12. Prove that $\frac{\sin \theta - 2 \sin^3 \theta}{2 \cos^3 \theta - \cos \theta} = \tan \theta$. **ICSE 2017, 3 marks**

SOLUTION

$$\text{LHS} = \frac{\sin \theta (1 - 2 \sin^2 \theta)}{\cos \theta (2 \cos^2 \theta - 1)}$$

Using $1 = \sin^2 \theta + \cos^2 \theta$: $1 - 2 \sin^2 \theta = \cos^2 \theta - \sin^2 \theta$ and $2 \cos^2 \theta - 1 = \cos^2 \theta - \sin^2 \theta$.

The brackets cancel: LHS = $\frac{\sin \theta}{\cos \theta} = \tan \theta = \text{RHS}$. **Proved.**

3 Complementary angles

13. Without using tables, evaluate $\sin^2 28^\circ + \sin^2 62^\circ + \tan^2 38^\circ - \cot^2 52^\circ + \frac{1}{4} \sec^2 30^\circ$. ICSE 2017, 3 marks

SOLUTION

$\sin 62^\circ = \cos 28^\circ$, so $\sin^2 28^\circ + \sin^2 62^\circ = \sin^2 28^\circ + \cos^2 28^\circ = 1$.

$\cot 52^\circ = \tan 38^\circ$, so $\tan^2 38^\circ - \cot^2 52^\circ = 0$.

$$\frac{1}{4} \sec^2 30^\circ = \frac{1}{4} \times \frac{4}{3} = \frac{1}{3}$$

Answer: $1 + 0 + \frac{1}{3} = \frac{4}{3}$

These questions are based on questions asked in past board examinations. They have been reworded by MathSetu, the figures are redrawn, and the solutions are MathSetu's own. MathSetu is an independent study resource and is not affiliated with or endorsed by CBSE, CISCE, NCERT or any publisher.