

CLASS 9 · ICSE · SELINA CONCISE MATHEMATICS · SHORT NOTES

Chapter 6: Simultaneous Linear Equations

Topic-wise short notes with solved examples

Solving a pair of linear equations in two variables by substitution, elimination and cross-multiplication, equations that reduce to linear form, and word problems.

1 A pair of linear equations

- A **linear equation in two variables** has the form $ax + by + c = 0$, where a and b are not both zero.
- Two such equations taken together are **simultaneous linear equations**. A **solution** is a pair (x, y) that satisfies both.
- Always check the answer by substituting in **both** equations.

2 Method of substitution

- From one equation, express one variable in terms of the other.
- Substitute this in the other equation and solve the resulting equation in one variable.
- Substitute back to find the second variable.

Example 2.1 Solve $x + y = 7$ and $3x - 2y = 11$ by substitution.

SOLUTION

From the first, $y = 7 - x$.

$3x - 2(7 - x) = 11$ gives $5x = 25$, so $x = 5$.

$x = 5, y = 2$

3 Method of elimination

- Multiply the equations by suitable numbers so that the coefficients of one variable become **equal in size**.
- **Add** the equations if those coefficients have opposite signs; **subtract** if they have the same sign.
- Solve for the remaining variable and substitute back.

Example 3.1 Solve $3x + 4y = 10$ and $2x - 3y = 1$ by elimination.

SOLUTION

Multiply the first by 3 and the second by 4: $9x + 12y = 30$ and $8x - 12y = 4$.

Add: $17x = 34$, so $x = 2$. Then $6 + 4y = 10$.

$$x = 2, y = 1$$

Example 3.2 Solve $41x + 53y = 135$ and $53x + 41y = 147$.

SOLUTION

Add: $94x + 94y = 282$, so $x + y = 3$. Subtract: $12x - 12y = 12$, so $x - y = 1$.

$$x = 2, y = 1$$

4 Method of cross-multiplication

- For $a_1x + b_1y + c_1 = 0$ and $a_2x + b_2y + c_2 = 0$:

$$\frac{x}{b_1c_2 - b_2c_1} = \frac{y}{c_1a_2 - c_2a_1} = \frac{1}{a_1b_2 - a_2b_1}$$

- Write both equations with zero on the right before using the formula.
- The method works when $a_1b_2 - a_2b_1 \neq 0$.

Example 4.1 Solve $2x + 3y - 17 = 0$ and $3x - 2y - 6 = 0$ by cross-multiplication.

SOLUTION

$$\frac{x}{(3)(-6) - (-2)(-17)} = \frac{y}{(-17)(3) - (-6)(2)} = \frac{1}{(2)(-2) - (3)(3)}$$

$$\frac{x}{-52} = \frac{y}{-39} = \frac{1}{-13}$$

$$x = 4, y = 3$$

5 Equations reducible to linear form

- If the variables appear as $\frac{1}{x}$ and $\frac{1}{y}$, put $\frac{1}{x} = a$ and $\frac{1}{y} = b$, solve for a and b , then invert.
- If they appear as $\frac{1}{x+y}$ and $\frac{1}{x-y}$, use these as the new variables in the same way.
- Equations with xy on the right, such as $2x + 3y = 5xy$: divide every term by xy .

Example 5.1 Solve $\frac{2}{x} + \frac{3}{y} = 13$ and $\frac{5}{x} - \frac{4}{y} = -2$.

SOLUTION

With $a = \frac{1}{x}$, $b = \frac{1}{y}$: $2a + 3b = 13$ and $5a - 4b = -2$.

Multiply by 4 and 3 and add: $23a = 46$, so $a = 2$ and $b = 3$.

$$x = \frac{1}{2}, y = \frac{1}{3}$$

Example 5.2 Solve $\frac{5}{x+y} - \frac{2}{x-y} = -1$ and $\frac{15}{x+y} + \frac{7}{x-y} = 10$.

SOLUTION

With $a = \frac{1}{x+y}$, $b = \frac{1}{x-y}$: $5a - 2b = -1$ and $15a + 7b = 10$.

Three times the first: $15a - 6b = -3$. Subtract: $13b = 13$, so $b = 1$ and $a = \frac{1}{5}$.

$$x + y = 5 \text{ and } x - y = 1$$

$$x = 3, y = 2$$

6 Word problems

- Choose the two unknowns, form two equations from the conditions, solve and check against the words.

Type	How to set up
Two-digit number	Tens digit x , units digit y : number $10x + y$, reversed $10y + x$
Fraction	Numerator x , denominator y
Ages	Present ages x and y ; after t years $x + t$, $y + t$
Boat and stream	Speed downstream $x + y$, upstream $x - y$
Fixed charge plus rate	Total = fixed charge + rate $\times$ quantity

Example 6.1 The sum of the digits of a two-digit number is 9. The number obtained by reversing the digits is 27 more than the number. Find the number.

SOLUTION

$$x + y = 9 \text{ and } (10y + x) - (10x + y) = 27, \text{ that is } y - x = 3.$$

$$x = 3, y = 6; \text{ the number is } 36$$

Example 6.2 A fraction becomes $\frac{1}{2}$ when 1 is added to the numerator and becomes $\frac{1}{3}$ when 1 is added to the denominator. Find it.

SOLUTION

$$2(x + 1) = y \text{ and } 3x = y + 1. \text{ So } 3x = 2x + 3.$$

$$x = 3, y = 8; \text{ the fraction is } \frac{3}{8}$$

Example 6.3 A boat goes 30 km downstream in 2 hours and 18 km upstream in 2 hours. Find the speed of the boat in still water and the speed of the stream.

SOLUTION

$$x + y = 15 \text{ and } x - y = 9$$

Boat 12 km/h; stream 3 km/h

Example 6.4 Five years ago a man was 7 times as old as his son. Five years from now he will be 3 times as old. Find their present ages.

SOLUTION

$$x - 5 = 7(y - 5) \text{ and } x + 5 = 3(y + 5), \text{ that is } x - 7y = -30 \text{ and } x - 3y = 10.$$

$$\text{Subtract: } 4y = 40$$

Son 10 years, father 40 years

7 Common mistakes

- Changing the sign of only some terms when subtracting one equation from another.
- Multiplying only one side of an equation.
- In cross-multiplication, leaving the constants on the right-hand side.
- Giving a and b as the answer in reducible equations instead of x and y .
- Writing a two-digit number as xy or $x + y$ instead of $10x + y$.
- Not checking the solution in the second equation.