

CLASS 10 · CBSE · NCERT MATHEMATICS · PYQ SOLUTIONS

Chapter 8: Introduction to Trigonometry

Step-by-step solutions to the previous year questions

The question numbers match the PYQ question sheet.

1 Trigonometric ratios

1. If $\sin \theta = \frac{a}{b}$, write $\cos \theta$ in terms of a and b . (a) $\frac{b}{\sqrt{b^2 - a^2}}$ (b) $\frac{b}{a}$ (c) $\frac{\sqrt{b^2 - a^2}}{b}$ (d) $\frac{a}{\sqrt{b^2 - a^2}}$

CBSE 2026, 1 mark

SOLUTION

Take a right triangle with the side opposite θ equal to a and the hypotenuse equal to b .

The adjacent side is $\sqrt{b^2 - a^2}$, so $\cos \theta = \frac{\sqrt{b^2 - a^2}}{b}$.

Answer: (c)

2. If $\tan \theta = \frac{24}{7}$, find $\sin \theta + \cos \theta$. CBSE 2026, 2 marks

SOLUTION

Take the opposite side as 24 and the adjacent side as 7. Hypotenuse = $\sqrt{24^2 + 7^2} = \sqrt{625} = 25$.

$$\sin \theta = \frac{24}{25} \text{ and } \cos \theta = \frac{7}{25}$$

Answer: $\frac{31}{25}$

3. If $\cot \theta = \frac{7}{8}$, evaluate $\frac{(1 + \sin \theta)(1 - \sin \theta)}{(1 + \cos \theta)(1 - \cos \theta)}$. CBSE 2026, 2 marks

SOLUTION

Numerator = $1 - \sin^2 \theta = \cos^2 \theta$ and denominator = $1 - \cos^2 \theta = \sin^2 \theta$

$$\text{Value} = \frac{\cos^2 \theta}{\sin^2 \theta} = \cot^2 \theta = \left(\frac{7}{8}\right)^2$$

Answer: $\frac{49}{64}$

4. If $2 \tan A = 3$, find the value of $\frac{4 \sin A + 3 \cos A}{4 \sin A - 3 \cos A}$. (a) $\frac{7}{\sqrt{13}}$ (b) $\frac{1}{\sqrt{13}}$ (c) 3 (d) does not exist

CBSE 2023, 1 mark

SOLUTION

Divide the numerator and the denominator by $\cos A$: $\frac{4 \tan A + 3}{4 \tan A - 3}$

$\tan A = \frac{3}{2}$, so $4 \tan A = 6$ and the value is $\frac{6+3}{6-3} = 3$.

Answer: (c)

5. If $\sin \alpha = \frac{1}{\sqrt{2}}$ and $\cot \beta = \sqrt{3}$, find $\operatorname{cosec} \alpha + \operatorname{cosec} \beta$. CBSE 2023, 2 marks

SOLUTION

$$\operatorname{cosec} \alpha = \frac{1}{\sin \alpha} = \sqrt{2}$$

$\cot \beta = \sqrt{3}$ gives $\beta = 30^\circ$, so $\operatorname{cosec} \beta = 2$.

Answer: $2 + \sqrt{2}$

6. If $4 \tan \theta = 3$, evaluate $\frac{4 \sin \theta - \cos \theta + 1}{4 \sin \theta + \cos \theta - 1}$. CBSE 2018, 3 marks

SOLUTION

$\tan \theta = \frac{3}{4}$. Take the opposite side as 3 and the adjacent side as 4, so the hypotenuse is 5.

$$\sin \theta = \frac{3}{5} \text{ and } \cos \theta = \frac{4}{5}$$

$$\text{Numerator} = \frac{12}{5} - \frac{4}{5} + 1 = \frac{13}{5} \text{ and denominator} = \frac{12}{5} + \frac{4}{5} - 1 = \frac{11}{5}$$

Answer: $\frac{13}{11}$

2 Ratios of standard angles

7. If $\cos A = \frac{1}{2}$, find the value of $\sin^2 A + 2 \cos^2 A$. (a) $\frac{3}{2}$ (b) $\frac{5}{4}$ (c) -1 (d) $\frac{1}{2}$ CBSE 2026, 1 mark

SOLUTION

$$\cos^2 A = \frac{1}{4}, \text{ so } \sin^2 A = 1 - \frac{1}{4} = \frac{3}{4}$$

$$\sin^2 A + 2 \cos^2 A = \frac{3}{4} + \frac{2}{4} = \frac{5}{4}$$

Answer: (b)

8. Find the acute angle θ for which $7 + 4 \sin \theta = 9$. CBSE 2025, 1 mark

SOLUTION

$$4 \sin \theta = 2, \text{ so } \sin \theta = \frac{1}{2}$$

Answer: $\theta = 30^\circ$

- 9.** Given that $x \cos 60^\circ + y \cos 0^\circ + \sin 30^\circ - \cot 45^\circ = 5$, find the value of $x + 2y$. **CBSE 2025, 2 marks**

SOLUTION

Put in the values: $x \times \frac{1}{2} + y \times 1 + \frac{1}{2} - 1 = 5$

$\frac{x}{2} + y = \frac{11}{2}$. Multiply both sides by 2.

Answer: $x + 2y = 11$

- 10.** Evaluate $\frac{\tan^2 60^\circ}{\sin^2 60^\circ + \cos^2 30^\circ}$. **CBSE 2025, 2 marks**

SOLUTION

$\tan 60^\circ = \sqrt{3}$, $\sin 60^\circ = \frac{\sqrt{3}}{2}$ and $\cos 30^\circ = \frac{\sqrt{3}}{2}$

Value = $\frac{3}{\frac{3}{4} + \frac{3}{4}} = \frac{3}{\frac{3}{2}}$

Answer: 2

- 11.** Given $\cos(\alpha + \beta) = 0$, find $\cos\left(\frac{\alpha + \beta}{2}\right)$. (a) $\frac{1}{\sqrt{2}}$ (b) $\frac{1}{2}$ (c) 0 (d) $\sqrt{2}$ **CBSE 2024, 1 mark**

SOLUTION

$\cos(\alpha + \beta) = 0 = \cos 90^\circ$, so $\alpha + \beta = 90^\circ$.

$\cos\left(\frac{\alpha + \beta}{2}\right) = \cos 45^\circ = \frac{1}{\sqrt{2}}$

Answer: (a)

- 12.** Find the value of $2\sqrt{2} \cos 45^\circ \sin 30^\circ + 2\sqrt{3} \cos 30^\circ$. **CBSE 2024, 2 marks**

SOLUTION

$= 2\sqrt{2} \times \frac{1}{\sqrt{2}} \times \frac{1}{2} + 2\sqrt{3} \times \frac{\sqrt{3}}{2}$

$= 1 + 3$

Answer: 4

- 13.** For $A = 60^\circ$ and $B = 30^\circ$, verify that $\sin(A + B) = \sin A \cos B + \cos A \sin B$. **CBSE 2024, 2 marks**

SOLUTION

LHS = $\sin(60^\circ + 30^\circ) = \sin 90^\circ = 1$

RHS = $\sin 60^\circ \cos 30^\circ + \cos 60^\circ \sin 30^\circ = \frac{\sqrt{3}}{2} \times \frac{\sqrt{3}}{2} + \frac{1}{2} \times \frac{1}{2} = \frac{3}{4} + \frac{1}{4} = 1$

LHS = RHS. Verified.

- 14.** Evaluate $\frac{3}{4} \tan^2 30^\circ - \sec^2 45^\circ + \sin^2 60^\circ$. (a) -1 (b) $\frac{5}{6}$ (c) $-\frac{3}{2}$ (d) $\frac{1}{6}$ **CBSE 2023, 1 mark**

SOLUTION

$$\tan^2 30^\circ = \frac{1}{3}, \sec^2 45^\circ = 2 \text{ and } \sin^2 60^\circ = \frac{3}{4}$$

$$\text{Value} = \frac{3}{4} \times \frac{1}{3} - 2 + \frac{3}{4} = \frac{1}{4} - 2 + \frac{3}{4} = -1$$

Answer: (a)

3 Trigonometric identities

- 15.** Given $x = h + a \cos \theta$ and $y = k + b \sin \theta$, prove that $\left(\frac{x-h}{a}\right)^2 + \left(\frac{y-k}{b}\right)^2 = 1$. **CBSE 2026, 3 marks**

SOLUTION

From the first relation, $x - h = a \cos \theta$, so $\frac{x-h}{a} = \cos \theta$.

From the second, $y - k = b \sin \theta$, so $\frac{y-k}{b} = \sin \theta$.

Squaring and adding: LHS = $\cos^2 \theta + \sin^2 \theta = 1 = \text{RHS}$. **Proved.**

- 16.** Prove that $\frac{\tan A}{1 + \sec A} - \frac{\tan A}{1 - \sec A} = 2 \operatorname{cosec} A$. **CBSE 2026, 3 marks**

SOLUTION

$$\text{LHS} = \tan A \left[\frac{(1 - \sec A) - (1 + \sec A)}{(1 + \sec A)(1 - \sec A)} \right] = \frac{-2 \tan A \sec A}{1 - \sec^2 A}$$

Since $\sec^2 A - 1 = \tan^2 A$, the denominator is $-\tan^2 A$.

$$\begin{aligned} \text{LHS} &= \frac{-2 \tan A \sec A}{-\tan^2 A} = \frac{2 \sec A}{\tan A} = \frac{2}{\cos A} \times \frac{\cos A}{\sin A} = \frac{2}{\sin A} \\ &= 2 \operatorname{cosec} A = \text{RHS. } \mathbf{Proved.} \end{aligned}$$

- 17.** Simplify $\tan^2 \theta - \left(\frac{1}{\cos \theta} \times \sec \theta\right)$. **CBSE 2025, 1 mark**

SOLUTION

$$\frac{1}{\cos \theta} = \sec \theta, \text{ so the bracket is } \sec^2 \theta.$$

$$\tan^2 \theta - \sec^2 \theta = -(\sec^2 \theta - \tan^2 \theta) = -1$$

Answer: -1

- 18.** Prove that $\frac{\tan \theta}{1 - \cot \theta} + \frac{\cot \theta}{1 - \tan \theta} = 1 + \sec \theta \operatorname{cosec} \theta$. **CBSE 2025, 3 marks**

SOLUTION

Write everything in terms of $\sin \theta$ and $\cos \theta$.

$$\text{First term: } \frac{\sin \theta}{\cos \theta} \div \frac{\sin \theta - \cos \theta}{\sin \theta} = \frac{\sin^2 \theta}{\cos \theta (\sin \theta - \cos \theta)}$$

$$\text{Second term: } \frac{\cos \theta}{\sin \theta} \div \frac{\cos \theta - \sin \theta}{\cos \theta} = -\frac{\cos^2 \theta}{\sin \theta (\sin \theta - \cos \theta)}$$

$$\text{LHS} = \frac{\sin^3 \theta - \cos^3 \theta}{\sin \theta \cos \theta (\sin \theta - \cos \theta)}$$

$$\begin{aligned} \text{Using } a^3 - b^3 &= (a - b)(a^2 + ab + b^2): \text{LHS} = \frac{\sin^2 \theta + \sin \theta \cos \theta + \cos^2 \theta}{\sin \theta \cos \theta} = \frac{1 + \sin \theta \cos \theta}{\sin \theta \cos \theta} \\ &= \frac{1}{\sin \theta \cos \theta} + 1 = 1 + \sec \theta \operatorname{cosec} \theta = \text{RHS. Proved.} \end{aligned}$$

- 19.** Prove that $\frac{\sin A + \cos A}{\sin A - \cos A} + \frac{\sin A - \cos A}{\sin A + \cos A} = \frac{2}{2 \sin^2 A - 1}$. **CBSE 2025, 3 marks**

SOLUTION

Take the common denominator $(\sin A - \cos A)(\sin A + \cos A) = \sin^2 A - \cos^2 A$.

$$\text{Numerator} = (\sin A + \cos A)^2 + (\sin A - \cos A)^2 = 2 \sin^2 A + 2 \cos^2 A = 2$$

$$\begin{aligned} \text{LHS} &= \frac{2}{\sin^2 A - \cos^2 A} = \frac{2}{\sin^2 A - (1 - \sin^2 A)} \\ &= \frac{2}{2 \sin^2 A - 1} = \text{RHS. Proved.} \end{aligned}$$

- 20.** Given $\sec \theta - \tan \theta = m$, find $\sec \theta + \tan \theta$. (a) $1 - \frac{1}{m}$ (b) $m^2 - 1$ (c) $\frac{1}{m}$ (d) $-m$ **CBSE 2024, 1 mark**

SOLUTION

$$\sec^2 \theta - \tan^2 \theta = 1, \text{ so } (\sec \theta + \tan \theta)(\sec \theta - \tan \theta) = 1.$$

$$\sec \theta + \tan \theta = \frac{1}{\sec \theta - \tan \theta} = \frac{1}{m}$$

Answer: (c)

- 21.** Prove the identity $\frac{\tan \theta}{1 - \cot \theta} + \frac{\cot \theta}{1 - \tan \theta} = 1 + \sec \theta \operatorname{cosec} \theta$. **CBSE 2024, 3 marks**

SOLUTION

$$\text{Write } s = \sin \theta \text{ and } c = \cos \theta. \text{ Then } \frac{\tan \theta}{1 - \cot \theta} = \frac{s/c}{(s-c)/s} = \frac{s^2}{c(s-c)} \text{ and } \frac{\cot \theta}{1 - \tan \theta} = \frac{c/s}{(c-s)/c} = -\frac{c^2}{s(s-c)}.$$

$$\begin{aligned} \text{LHS} &= \frac{s^3 - c^3}{sc(s-c)} = \frac{(s-c)(s^2 + sc + c^2)}{sc(s-c)} = \frac{s^2 + c^2 + sc}{sc} \\ &= \frac{1 + sc}{sc} = \frac{1}{sc} + 1 \\ &= 1 + \sec \theta \operatorname{cosec} \theta = \text{RHS} \end{aligned}$$

- 22.** If $\sin \theta + \cos \theta = \sqrt{3}$, find the value of $\sin \theta \cos \theta$. **CBSE 2023, 2 marks**

SOLUTION

Square both sides: $\sin^2 \theta + \cos^2 \theta + 2 \sin \theta \cos \theta = 3$

$$1 + 2 \sin \theta \cos \theta = 3$$

Answer: $\sin \theta \cos \theta = 1$

- 23.** Prove that $\frac{\tan \theta + \sec \theta - 1}{\tan \theta - \sec \theta + 1} = \frac{1 + \sin \theta}{\cos \theta}$. **CBSE 2023, 3 marks**

SOLUTION

In the numerator, replace 1 by $\sec^2 \theta - \tan^2 \theta$.

$$\text{Numerator} = (\sec \theta + \tan \theta) - (\sec \theta + \tan \theta)(\sec \theta - \tan \theta) = (\sec \theta + \tan \theta)(1 - \sec \theta + \tan \theta)$$

The second bracket is the same as the denominator, so LHS = $\sec \theta + \tan \theta$.

$$= \frac{1}{\cos \theta} + \frac{\sin \theta}{\cos \theta} = \frac{1 + \sin \theta}{\cos \theta} = \text{RHS. Proved.}$$

- 24.** Prove that $(\sin \theta + \operatorname{cosec} \theta)^2 + (\cos \theta + \sec \theta)^2 = 7 + \tan^2 \theta + \cot^2 \theta$. **CBSE 2019, 3 marks**

SOLUTION

$$\text{LHS} = \sin^2 \theta + \operatorname{cosec}^2 \theta + 2 \sin \theta \operatorname{cosec} \theta + \cos^2 \theta + \sec^2 \theta + 2 \cos \theta \sec \theta$$

$$\sin \theta \operatorname{cosec} \theta = 1 \text{ and } \cos \theta \sec \theta = 1, \text{ so LHS} = (\sin^2 \theta + \cos^2 \theta) + 4 + \operatorname{cosec}^2 \theta + \sec^2 \theta$$

$$= 1 + 4 + (1 + \cot^2 \theta) + (1 + \tan^2 \theta)$$

$$= 7 + \tan^2 \theta + \cot^2 \theta = \text{RHS. Proved.}$$

- 25.** Prove that $(1 + \cot A - \operatorname{cosec} A)(1 + \tan A + \sec A) = 2$. **CBSE 2019, 3 marks**

SOLUTION

$$\begin{aligned} \text{LHS} &= \left(\frac{\sin A + \cos A - 1}{\sin A} \right) \left(\frac{\cos A + \sin A + 1}{\cos A} \right) \\ &= \frac{(\sin A + \cos A)^2 - 1}{\sin A \cos A} = \frac{\sin^2 A + \cos^2 A + 2 \sin A \cos A - 1}{\sin A \cos A} \\ &= \frac{2 \sin A \cos A}{\sin A \cos A} = 2 = \text{RHS. Proved.} \end{aligned}$$

- 26.** Prove that $\frac{\sin A - \cos A + 1}{\sin A + \cos A - 1} = \frac{1}{\sec A - \tan A}$. **CBSE 2019, 4 marks**

SOLUTION

$$\text{Divide the numerator and the denominator by } \cos A: \text{LHS} = \frac{\tan A - 1 + \sec A}{\tan A + 1 - \sec A}$$

$$\text{In the denominator, write } 1 = \sec^2 A - \tan^2 A = (\sec A - \tan A)(\sec A + \tan A).$$

$$\text{Denominator} = (\tan A - \sec A) + (\sec A - \tan A)(\sec A + \tan A) = (\sec A - \tan A)(\sec A + \tan A - 1)$$

$$\text{LHS} = \frac{\tan A + \sec A - 1}{(\sec A - \tan A)(\sec A + \tan A - 1)} = \frac{1}{\sec A - \tan A} = \text{RHS. Proved.}$$

27. Prove that $\frac{\sin A - 2 \sin^3 A}{2 \cos^3 A - \cos A} = \tan A$. **CBSE 2018, 4 marks**

SOLUTION

$$\text{LHS} = \frac{\sin A(1 - 2 \sin^2 A)}{\cos A(2 \cos^2 A - 1)}$$

Using $1 = \sin^2 A + \cos^2 A$: $1 - 2 \sin^2 A = \cos^2 A - \sin^2 A$ and $2 \cos^2 A - 1 = \cos^2 A - \sin^2 A$.

The two brackets are equal and cancel: $\text{LHS} = \frac{\sin A}{\cos A} = \tan A = \text{RHS}$. **Proved.**

These questions are based on questions asked in past board examinations. They have been reworded by MathSetu, the figures are redrawn, and the solutions are MathSetu's own. MathSetu is an independent study resource and is not affiliated with or endorsed by CBSE, CISCE, NCERT or any publisher.