

CLASS 9 · CBSE · GANITA MANJARI PART 1 · SHORT NOTES

Chapter 4: Exploring Algebraic Identities

Topic-wise short notes with solved examples

An identity is an equation that is true for every value of its variables. This chapter builds the standard identities for squares and cubes and uses them to calculate quickly, factorise expressions and simplify algebraic fractions.


1 Identity and equation

- An **equation** is true only for some values of the variable. $x^2 - 1 = 24$ is true only for $x = 5$ and $x = -5$.
- An **identity** is true for **all** values of the variables. $(x + y)^2 = x^2 + 2xy + y^2$ holds whatever numbers x and y are: positive, negative or fractions.
- An identity can be checked with numbers, but it is **proved** with the distributive property.
- In general $(a + b)^2 \neq a^2 + b^2$. The two differ by $2ab$, so they are equal only when $a = 0$ or $b = 0$.

2 Squares of a sum and a difference

$$(a + b)^2 = a^2 + 2ab + b^2 \quad (a - b)^2 = a^2 - 2ab + b^2$$

- Proof: $(a + b)^2 = (a + b)(a + b) = a^2 + ab + ba + b^2 = a^2 + 2ab + b^2$. Replacing b by $-b$ gives the second identity.
- Picture: a square of side $a + b$ is made of a square of area a^2 , a square of area b^2 and two rectangles of area ab each.



A square of side $a + b$

Example 2.1 Expand $(3x + 5y)^2$ and $(4a - 3b)^2$.

SOLUTION

$$(3x)^2 + 2(3x)(5y) + (5y)^2 \text{ and } (4a)^2 - 2(4a)(3b) + (3b)^2$$

$$9x^2 + 30xy + 25y^2 \text{ and } 16a^2 - 24ab + 9b^2$$

Example 2.2 Find 52^2 and 38^2 without multiplying directly.

SOLUTION

$$52^2 = (50 + 2)^2 = 2500 + 200 + 4 = 2704$$

$$38^2 = (40 - 2)^2 = 1600 - 160 + 4 = 1444$$

2704 and 1444

Example 2.3 Take any three consecutive square numbers. Show that (smallest + largest) $- 2 \times$ (middle) is always 2.

SOLUTION

Let the numbers be $n - 1$, n and $n + 1$.

$$(n - 1)^2 + (n + 1)^2 = (n^2 - 2n + 1) + (n^2 + 2n + 1) = 2n^2 + 2$$

$$2n^2 + 2 - 2n^2 = 2, \text{ whatever } n \text{ is.}$$

3 Square of a sum of three terms

$$(a + b + c)^2 = a^2 + b^2 + c^2 + 2ab + 2bc + 2ca$$

- Proof: put $d = b + c$. Then $(a + d)^2 = a^2 + 2a(b + c) + (b + c)^2$, which expands to the result.
- For a minus sign, treat the term as negative: the squares stay positive and only the products change sign.

Example 3.1 Expand $(2p + q - 3r)^2$.

SOLUTION

Take $a = 2p$, $b = q$, $c = -3r$.

$$4p^2 + q^2 + 9r^2 + 2(2p)(q) + 2(q)(-3r) + 2(-3r)(2p)$$

$$4p^2 + q^2 + 9r^2 + 4pq - 6qr - 12pr$$

Example 3.2 Find 112^2 using this identity.

SOLUTION

$$(100 + 10 + 2)^2 = 10000 + 100 + 4 + 2000 + 40 + 400$$

12544

4 Difference of two squares

$$a^2 - b^2 = (a + b)(a - b)$$

- Written as $a^2 = (a + b)(a - b) + b^2$, it gives a quick way to square a number. Śhrīdharāchārya gave this method around 750 CE.

Example 4.1 Find 47×53 and 45^2 .

SOLUTION

$$47 \times 53 = (50 - 3)(50 + 3) = 2500 - 9 = 2491$$

$$45^2 = (45 + 5)(45 - 5) + 5^2 = 50 \times 40 + 25 = 2025$$

2491 and 2025

5 Factorising with the square identities

- **Factorising** is writing an expression as a product. Read the identities from right to left.
- First take out any common factor. Then look for two perfect squares and check the middle term.

Example 5.1 Factorise $49m^2 + 42m + 9$ and $16y^2 - 24y + 9$.

SOLUTION

$$49m^2 = (7m)^2, 9 = 3^2 \text{ and } 42m = 2(7m)(3)$$

$$16y^2 = (4y)^2, 9 = 3^2 \text{ and } 24y = 2(4y)(3)$$

$$(7m + 3)^2 \text{ and } (4y - 3)^2$$

Example 5.2 Factorise $12a^2 - 60ab + 75b^2$ and $36s^2 - 49t^2$.

SOLUTION

$$12a^2 - 60ab + 75b^2 = 3(4a^2 - 20ab + 25b^2) = 3(2a - 5b)^2$$

$$36s^2 - 49t^2 = (6s)^2 - (7t)^2$$

$$3(2a - 5b)^2 \text{ and } (6s + 7t)(6s - 7t)$$

6 Products and factors of the form $x^2 + (a + b)x + ab$

$$(x + a)(x + b) = x^2 + (a + b)x + ab \quad (px + a)(qx + b) = pqx^2 + (pb + qa)x + ab$$

- **Algebra tiles** show this as a rectangle: one x^2 -tile, $(a + b)$ x -tiles and ab unit tiles fit into a rectangle of sides $x + a$ and $x + b$.
- To factorise $x^2 + sx + p$, find two numbers whose **sum is** s and whose **product is** p . Then split the middle term.
- To factorise $ax^2 + bx + c$, find two numbers whose sum is b and whose product is ac , split the middle term and group.

Example 6.1 Factorise $x^2 + 9x + 20$ and $x^2 - x - 12$.

SOLUTION

Sum 9, product 20: the numbers are 4 and 5.

Sum -1 , product -12 : the numbers are -4 and 3 .

$$(x + 4)(x + 5) \text{ and } (x - 4)(x + 3)$$

Example 6.2 Factorise $6x^2 + 11x + 3$.

SOLUTION

Sum 11, product $6 \times 3 = 18$: the numbers are 9 and 2.

$$6x^2 + 9x + 2x + 3 = 3x(2x + 3) + 1(2x + 3)$$

$$(3x + 1)(2x + 3)$$

Example 6.3 A rectangular garden is 5 m shorter in breadth than in length, and its area is 84 m^2 . Find its length and breadth.

SOLUTION

Let the length be x m. Then $x(x - 5) = 84$, so $x^2 - 5x - 84 = 0$.

Sum -5 , product -84 : the numbers are -12 and 7 . So $(x - 12)(x + 7) = 0$.

$x = 12$ or $x = -7$. A length cannot be negative.

Length 12 m; breadth 7 m

7 Cubes of a sum and a difference

$$(a + b)^3 = a^3 + 3a^2b + 3ab^2 + b^3 \quad (a - b)^3 = a^3 - 3a^2b + 3ab^2 - b^3$$

- Proof: $(a + b)^3 = (a + b)(a^2 + 2ab + b^2)$, then use the distributive property.
- Picture: a cube of edge $a + b$ splits into a cube of volume a^3 , a cube of volume b^3 , three cuboids of volume a^2b and three of volume ab^2 .
- In $(a - b)^3$ the signs alternate: $+$, $-$, $+$, $-$.

Example 7.1 Expand $(2x + 3)^3$ and find 98^3 .

SOLUTION

$$(2x)^3 + 3(2x)^2(3) + 3(2x)(3)^2 + 3^3 = 8x^3 + 36x^2 + 54x + 27$$

$$98^3 = (100 - 2)^3 = 1000000 - 3(10000)(2) + 3(100)(4) - 8 = 1000000 - 60000 + 1200 - 8$$

$$8x^3 + 36x^2 + 54x + 27 \text{ and } 941192$$

Example 7.2 A cube has volume $27a^3 + 54a^2b + 36ab^2 + 8b^3$ cubic units. Find its edge.

SOLUTION

$$27a^3 = (3a)^3, 8b^3 = (2b)^3, 54a^2b = 3(3a)^2(2b) \text{ and } 36ab^2 = 3(3a)(2b)^2$$

The volume is $(3a + 2b)^3$, so the edge is $3a + 2b$ units.

8 Sum and difference of two cubes

$$x^3 + y^3 = (x + y)(x^2 - xy + y^2) \quad x^3 - y^3 = (x - y)(x^2 + xy + y^2)$$

- Check by multiplying out: $(x - y)(x^2 + xy + y^2) = x^3 + x^2y + xy^2 - x^2y - xy^2 - y^3 = x^3 - y^3$.
- The sign in the first bracket matches the sign between the cubes. The middle term of the second bracket has the opposite sign.

Example 8.1 Factorise $8x^3 + 27$ and $a^3 - 64b^3$.

SOLUTION

$$8x^3 + 27 = (2x)^3 + 3^3 \text{ and } a^3 - 64b^3 = a^3 - (4b)^3$$

$$(2x + 3)(4x^2 - 6x + 9) \text{ and } (a - 4b)(a^2 + 4ab + 16b^2)$$

9 The identity for three cubes

$$x^3 + y^3 + z^3 - 3xyz = (x + y + z)(x^2 + y^2 + z^2 - xy - yz - zx)$$

- **Special case:** if $x + y + z = 0$, then $x^3 + y^3 + z^3 = 3xyz$.

Example 9.1 Three numbers have sum 6, the sum of their products in pairs is 11 and their product is 6. Find the sum of their cubes.

SOLUTION

$$x^2 + y^2 + z^2 = (x + y + z)^2 - 2(xy + yz + zx) = 36 - 22 = 14$$

$$x^3 + y^3 + z^3 - 3(6) = 6 \times (14 - 11) = 18$$

$$x^3 + y^3 + z^3 = 36$$

Example 9.2 Find $7^3 + (-4)^3 + (-3)^3$ without cubing.

SOLUTION

$$7 + (-4) + (-3) = 0, \text{ so the sum of the cubes is } 3xyz = 3 \times 7 \times (-4) \times (-3).$$

252

10 Simplifying rational expressions

- A **rational expression** is a fraction whose numerator and denominator are polynomials.

- To simplify, **factorise** the numerator and the denominator and cancel the common factors. A factor can be cancelled only if it is not zero.

Example 10.1 Simplify $\frac{x^2 - 9}{x^2 + x - 12}$, assuming the denominator is not zero.

SOLUTION

Numerator: $(x - 3)(x + 3)$. Denominator: sum 1, product -12 gives $(x + 4)(x - 3)$.

Cancel the common factor $(x - 3)$.

$$\frac{x + 3}{x + 4}$$

Example 10.2 Simplify $\frac{4y^2 - 20y + 25}{4y^2 - 25}$, assuming the denominator is not zero.

SOLUTION

Numerator: $(2y - 5)^2$. Denominator: $(2y - 5)(2y + 5)$.

$$\frac{2y - 5}{2y + 5}$$

11 Common mistakes

- Writing $(a + b)^2 = a^2 + b^2$ or $(a + b)^3 = a^3 + b^3$. The middle terms are missing.
- Squaring only the letter: $(3x)^2 = 9x^2$, not $3x^2$.
- Getting the sign wrong in $x^3 + y^3 = (x + y)(x^2 - xy + y^2)$. The middle term of the second bracket is $-xy$, not $-2xy$.
- Cancelling terms instead of factors: in $\frac{x^2 - 9}{x^2 + x - 12}$ the x^2 cannot be cancelled. Factorise first.
- Keeping a negative length. After solving, reject the value that does not fit the problem.