



Assignment Project Exam Help

Introduction to Database Systems – Part 2

<https://tutorcs.com>

Math Concepts

WeChat: cstutorcs



What are the Math Concepts behind Databases?

Assignment Project Exam Help

- Set

- Tuple

<https://tutorcs.com>

- Cartesian Product of Sets

WeChat: cstutorcs

- Relation



Set Notation

Assignment Project Exam Help

Container



<https://tutorcs.com>

WeChat: cstutorcs





Set Notation

Assignment Project Exam Help

- We need set notation to represent formal definitions in this course.

- A **set** is a collection of distinct elements.

<https://tutorcs.com>

- Two basic properties of sets

- The elements in a set have no order.

e.g., $\{1, 2, 3\} = \{2, 3, 1\}$

- Each element can not be in the set more than once.

e.g., $\{\text{Monday}, \text{Monday}, \text{Tuesday}, \text{Wednesday}, \text{Thursday}, \text{Friday}, \text{Friday}\}$ is Not a set. Note that **Multisets** allow to have duplicate elements.

WeChat: cstutorcs



Set Notation

Assignment Project Exam Help

Two ways of specifying a set

- 1 $\{x_1, \dots, x_n\}$ (i.e., list all the elements in a set)

<https://tutorcs.com>

- $\{2, 3, 4, 5\}$
- $\{\text{Sydney, Melbourne, Canberra}\}$
- $\{\}$ or \emptyset , i.e., the *empty* set.

WeChat: cstutorcs

- 2 $\{x \mid \varphi\}$ (i.e., describe the elements that satisfy a property φ)

- $\{x \mid x \text{ is a student currently enrolled in COMP7240}\}$
- $\{x \mid x \text{ is an integer and } x > 0\}$



Set Operations

Assignment Project Exam Help

• Membership: $x \in A$ if x is in the set A ; $x \notin A$ if x is not in the set A .

<https://tutorcs.com>



WeChat: cstutorcs

$1 \in \{1,2,3\}$ $3 \in \{1,2,3\}$

$2 \in \{1,2\}$ $3 \notin \{1,2\}$



Set Operations

- **Equality:** If A and B have the same elements, we write $A = B$; otherwise we write $A \neq B$.

- $\{x \mid x \text{ is an integer, } x > 1 \text{ and } x < 6\} = \{2, 3, 4, 5\}$

- If one set contains some element that is not in the other set, then they are different.



$$\{1, 2\} \neq \{1, 2, 3\}$$



Set Operations

- **Subset:** A is called a **subset** of B if every element of A is in B and we write $A \subseteq B$.
- **Proper subset:** A is called a **proper subset** of B if $A \subseteq B$ and A and B are not equal, and we write $A \subset B$.

<https://tutorcs.com>

WeChat: cstutorcs



$$\begin{aligned}\{1,2\} &\subseteq \{1,2,3\} & \{1,2\} &\subseteq \{1,2\} \\ \{1,2\} &\subset \{1,2,3\} & &\end{aligned}$$



Set Operations

- **Subset:** A is called a **subset** of B if every element of A is in B and we write $A \subseteq B$.
- **Proper subset:** A is called a **proper subset** of B if $A \subseteq B$ and A and B are not equal, and we write $A \subset B$.

<https://tutorcs.com>

\subseteq means \subset or $=$

$$\begin{aligned}\{1,2\} &\subseteq \{1,2,3\} & \{1,2\} &\subseteq \{1,2\} \\ \{1,2\} &\subset \{1,2,3\}\end{aligned}$$





Set Operations

Assignment Project Exam Help

• Union: $A \cup B$ for the set containing everything in A and everything in B .

• $\{3, 4, 5\} \cup \{3, 5, 7, 9\} = \{3, 4, 5, 7, 9\}.$

<https://tutorcs.com>

WeChat:    



Set Operations


Assignment Project Exam Help

• Intersection: $A \cap B$ for the set of elements that are in both A and B

• $\{3, 4, 5\} \cap \{3, 5, 7, 9\} = \{3, 5\}$.

<https://tutorcs.com>

WeChat: [cmtutorcs](#)





Set Operations


Assignment Project Exam Help

• Difference: $A - B$ is the elements from A that are *not* in B .

• $\{3, 4, 5\} - \{3, 5, 7, 9\} = \{4\}$.

<https://tutorcs.com>

WeChat: tutorcs





Set Operations – Exercise

Assignment Project Exam Help

- Let $A = \{1, 2, 3\}$ and $B = \{true, false\}$.

- Which of the following are correct?

1 $\{2\} \in A$ No! $\{2\} \subset A$ and $2 \in A$

2 $true \subset B$ No! $true \in B$ and $\{true\} \subset B$

3 $\{2, 3\} \subseteq A \cup B$ Yes! $A \cup B = \{1, 2, 3, true, false\}$

4 $2 \in A \cap B$ No! $A \cap B = \{\}$

5 $2 \in A - \{1, 3, 5\}$ Yes! $A - \{1, 3, 5\} = \{2\}$

6 $\{1, 4\} \subseteq A - B$ No! $A - B = \{1, 2, 3\}$

7 $\emptyset \cap B = \emptyset$ Yes! $\emptyset = \{\}$, the empty set



Australian
National
University

Tuple Notation

Assignment Project Exam Help

In Order



<https://tutorcs.com>

WeChat: cstutorcs





Tuple Notation

Assignment Project Exam Help

- A tuple is an ordered list of n elements.
 - $(1, 2, 3, 4, 5)$
 - $(\text{Melbourne}, \text{Sydney}, \text{Canberra})$
- Two tuples are **equal** if they have the same elements in the same order.
 - $(1, 2, 3) \neq (2, 3, 1)$ (i.e., the order does matter!)
- The same element can be in a tuple twice.
 - $(\text{Monday}, \text{Monday}, \text{Tuesday}, \text{Wednesday}, \text{Thursday}, \text{Friday}, \text{Friday})$ is a tuple.
- Ordered pairs are special cases of tuples.

<https://tutorcs.com>

WeChat: cstutorcs



Cartesian Product of Sets

Assignment Project Exam Help



<https://tutorcs.com>

WeChat: cstutorcs



Cartesian Product of Sets

Assignment Project Exam Help

<https://tutorcs.com>

WeChat: cstutorcs

$\{2, 3, 4, 5, 6, 7, 8, 9, 10, J, Q, K, A\}$





Cartesian Product of Sets

Assignment Project Exam Help

<https://tutorcs.com>

WeChat: cstutorcs



$\{2, 3, 4, 5, 6, 7, 8, 9, 10, J, Q, K, A\}$

$\{\spadesuit, \diamondsuit, \clubsuit, \heartsuit\}$



Cartesian Product of Sets

Assignment Project Exam Help

- The Cartesian product operation takes an ordered list of sets, and returns a set of tuples.

- **Cartesian product** $D_1 \times \dots \times D_n$ is the set of all possible combinations of values from the sets D_1, \dots, D_n .

- It contains all the tuples with the first element from the first set, the second element from the second set, ...

- For example $A \times B = \{(a, b) \mid a \in A \text{ and } b \in B\}$.

If $A = \{2, 3\}$ and $B = \{\text{Clubs}, \text{Diamonds}, \text{Hearts}, \text{Spades}\}$

Then $A \times B = \{(2, \text{Clubs}), (2, \text{Diamonds}), (2, \text{Hearts}), (2, \text{Spades}), (3, \text{Clubs}), (3, \text{Diamonds}), (3, \text{Hearts}), (3, \text{Spades})\}$.

$(2, \text{Clubs}) \in A \times B$, $(\text{Spades}, 3) \notin A \times B$, $(4, \text{Hearts}) \notin A \times B$

$\{(3, \text{Clubs}), (3, \text{Diamonds}), (3, \text{Hearts}), (3, \text{Spades})\} \subseteq A \times B$

<https://tutorcs.com>

WeChat: estutorcs



Relation Notation

Assignment Project Exam Help

$\{2, 3, 4, 5, 6, 7, 8, 9, 10, J, Q, K, A\}$

\times

$\{\spadesuit, \diamondsuit, \clubsuit, \heartsuit\}$

<https://tutorcs.com>

WeChat: cstutorcs





Relation Notation

Assignment Project Exam Help
 $\{2, 3, 4, 5, 6, 7, 8, 9, 10, J, Q, K, A\}$

$\{\spadesuit, \diamondsuit, \clubsuit, \heartsuit\}$

<https://tutorcs.com>

WeChat: cstutorcs



ROYAL FLUSH



STRAIGHT FLUSH



FOUR OF A KIND



FULL HOUSE



FLUSH



STRAIGHT



THREE OF A KIND



TWO PAIRS



ONE PAIR



HIGH HAND



Relation Notation

Assignment Project Exam Help

- A relation is a subset of a Cartesian product of sets.

- **Example**

- Let $X = \{Canberra, Paris, Tokyo, Kyoto\}$, and $Y = \{Australia, France, Japan\}$.

- Let $R = \{(a, b) | a \in X, b \in Y \text{ and } a \text{ is a city in } b\}$.

- It is easy to see that R is a relation

- $R \subseteq X \times Y$.

- $(Canberra, Australia) \in R, (Paris, France) \in R$
but $(Tokyo, France) \notin R, (France, Japan) \notin R$



Relation Notation

Assignment Project Exam Help

- A relation is a subset of a Cartesian product of sets.

- **Example**

• Let $\mathbb{Z} = \{\dots, -1, 0, 1, 2, \dots\}$ the set of all integers

- Let $R = \{(x, y) \mid x \in \mathbb{Z}, y \in \mathbb{Z} \text{ and } x < y\}$.

- It is easy to see that R is a relation.

WeChat: cstutorcs

- $R \subseteq \mathbb{Z} \times \mathbb{Z}$.

- $(0, 1) \in R, (-4, -2) \in R$
but $(0, 0) \notin R, (100, -2) \notin R$.