

Copyright ©Copyright University of New South Wales 2020. All rights reserved.

Course materials subject to Copyright

UNSW Sydney owns copyright in these materials (unless stated otherwise). The material is subject to copyright under Australian law and overseas under international treaties.

The materials are provided for use by enrolled UNSW students. The materials, or any part, may not be copied, shared or distributed, in print or digitally, outside the course without permission.

Students may only copy a reasonable portion of the material for personal research or study or for criticism or review. Under no circumstances may these materials be copied or reproduced for sale or commercial purposes without prior written permission of UNSW Sydney.

Statement on class recording

To ensure the free and open discussion of ideas, students may not record, by any means, classroom lectures, discussion and/or activities without the advance written permission of the instructor, and any such recording properly approved in advance can be used solely for the student's own private use.

WARNING: Your failure to comply with these conditions may lead to disciplinary action, and may give rise to a civil action or a criminal offence under the law.

THE ABOVE INFORMATION MUST NOT BE REMOVED FROM THIS MATERIAL.

WeChat: cstutorcs

Assignment Project Exam Help

Financial Econometrics

Slides-01: RETURN PROPERTIES Part II

<https://tutores.com>

Rachida Ouyse
School of Economics

¹©Copyright University of New South Wales 2020. All rights reserved. This copyright notice must not be removed from this material.

WeChat: cstutorcs



Shape Characteristics: Population

Let X_t be a random variable with pdf $f(x)$

Assignment Project Exam Help

$$\mu = E[X_t] : \text{center}$$

$$\sigma^2 = \text{var}(X_t) = E[(X_t - \mu)^2] : \text{spread}$$

$$\text{skewness}(X_t) = S(X) = E \left[\frac{(X_t - \mu)^3}{\sigma^3} \right] : \text{symmetry}$$

$$\text{kurtosis}(X_t) = K(X) = E \left[\frac{(X_t - \mu)^4}{\sigma^4} \right] : \text{tail thickness}$$

$$K(X) - 3 : \text{Excess kurtosis}$$

Note: The k^{th} moment and central moment of X_t are:

$$m'_k = E[X_t^k]$$

$$m_k = E[(X_t - \mu)^k]$$

Shape Characteristics of Random Variable

Assignment Project Exam Help

- Why are the mean and variance of returns important?

They are concerned with long-term return and risk, respectively.

- Why is return symmetry of interest in financial study?

Symmetry has important implications in holding short or long financial positions and in risk management.

- Why is kurtosis important?

Related to volatility forecasting, efficiency in estimation and tests, etc.

High kurtosis implies heavy (or long) tails in distribution.

Example: Normal Random Variable

Assignment Project Exam Help

Normal Distribution

$$X \sim N(\mu, \sigma^2)$$

$$f(x) = \frac{1}{\sqrt{2\pi\sigma^2}} \exp\left(-\frac{(x-\mu)^2}{2\sigma^2}\right) \quad -\infty \leq x \leq \infty$$

WeChat: cstutorcs

$$E[X] = \mu$$

$$\text{var}(X) = \sigma^2$$

$$\text{skew}(X) = 0$$

$$\text{kurt}(X) = 3$$

$$m_k = 0 \text{ for } k \text{ odd}$$

Shape Characteristics: Sample moments

Assignment Project Exam Help

Sample moments

Let $\{r_t, \dots, r_T\}$ denote a random sample of size T where r_t is a realization of the random variable \tilde{r} .

$$\hat{\mu} = \frac{1}{T} \sum_{t=1}^T r_t, \quad \hat{\sigma}^2 = \frac{1}{T-1} \sum_{t=1}^T (r_t - \hat{\mu})^2 = \hat{m}_2$$


$$\widehat{\text{skew}} = \frac{\hat{m}_3}{\hat{\sigma}^3}, \quad \widehat{\text{kurt}} = \frac{\hat{m}_4}{\hat{\sigma}^4}$$

$$\hat{m}_k = \frac{1}{T-1} \sum_{t=1}^T (r_t - \hat{\mu})^k,$$

Note: we divide by $T - 1$ to get unbiased estimates. Check software to see how moments are computed.

Shape Characteristics: Visually

Assignment Project Exam Help



Testing for normality

- **QQ plot:** plot standardized empirical quantiles vs. theoretical quantiles from specified distribution. Note: Shapiro-Wilks (SW) test for normality: correlation coefficient between values used in QQ-plot
- **Jarque-Bera (JB) test** for normality

$$JB = \frac{T}{6} \left(\widehat{skew}^2 + \frac{(\widehat{kurt} - 3)^2}{4} \right) \sim_A \chi^2(2)$$

Note: if r_t is $N(u, \sigma^2)$ then:

$$\sqrt{T}\widehat{skew} \sim N(0, 6), \text{ and } \sqrt{T}(\widehat{kurt} - 3) \sim N(0, 24)$$

Shape Characteristics: Normality test

Assignment Project Exam Help

The null hypothesis:

H_0 : Data (the return) X_t are Normally distributed.

- ① **Skewness test:** $Z_{sk} = \frac{\widehat{skew}}{\sqrt{6/T}} \sim N(0, 1)$

Reject H_0 if $|z_{sk}|$ is too large (> 1.96 , at 5%)

- ② **Kurtosis test:** $Z_{kt} = \frac{\widehat{kurt}-3}{\sqrt{24/T}} \sim N(0, 1)$

Reject H_0 if $|z_{kt}|$ is too large (> 1.96 , at 5%).

- ③ **Jarque-Bera test:** $JB = Z_{ks}^2 + Z_{kt}^2 \sim \chi^2_2$


Reject JB is too large (> 5.99 at 5%)

Example: Descriptive Statistics

- Descriptive statistics

e.g. NYSE index prices: (1995:01:03-2012:08:30)

Composite, Industrial,
Trans, Utility, Finance.



Descriptive statistics of log returns.

	Composite	Industrial	Trans	Utility	Finance
Mean	0.035	0.034	0.031	0.007	0.052
Std. Dev.	1.006	1.009	1.320	1.087	1.310
Skewness	-0.316	-0.386	-1.044	-0.275	-0.042
Kurtosis	7.224	7.755	18.103	5.637	5.772

Correlations of log returns

	Composite	Industrial	Trans	Utility	Finance
Composite	1				
Industrial	0.983	1			
Trans	0.731	0.708	1		
Utility	0.769	0.711	0.505	1	
Finance	0.885	0.800	0.668	0.623	1

Portfolio variance and diversification:

$$Z = \frac{1}{2}(Y + X),$$

$$\text{var}(Z) = \frac{1}{4}[\text{Var}(X) + \text{Var}(Y) + 2\text{Cov}(X, Y)]$$

Example: Descriptive Statistics

Assignment Project Exam Help



Normality test

eg. Comp. index log return

time series plot

<https://tutorcs.com>

histogram



Series:	RC
Sample:	1 1931
Observations:	1930
Mean	0.035370
Median	0.012255
Maximum	2.78714
Minimum	-8.71112
Std. Dev.	1.006207
Skewness	-0.315728
Kurtosis	7.224376
Jarque-Bera	1467.129
Probability	0.000000

p value =
 $P(\chi^2_{(2)} > JB)$

Stylized Fact: Large kurtosis



Assignment Project Exam Help

= Some stylised facts about index return series

- concentration around zero with a few large "outliers"
- large standard deviations (volatile)
- negative skewness (longer tail at the negative side)
- large kurtosis (tail probabilities larger than normal)
- large variation followed by large ones (clustering)

leptokurtic

<https://tutorcs.com>



Series: RC	
Sample 1 1931	
Observations 1930	
Mean	0.035300
Median	0.052285
Maximum	5.178704
Minimum	-6.791142
Std. Dev.	1.006207
Skewness	-0.315728
Kurtosis	7.224376
Jarque-Bera	1467.129
Probability	0.000000

Descriptive statistics: Autocorrelation

• Predictability

- We say X_{t+1} is predictable if information at t (eg. $\{X_t, X_{t-1}, \dots\}$), helps to improve our prediction of X_{t+1} .
- In particular, X_{t+1} is predictable if X_{t+1} is correlated with X_{t-j} for some $j > 0$ (ie. $Cov(X_{t+1}, X_{t-j}) \neq 0$).

• Autocorrelation Function (ACF)

- Autocovariance: $\gamma_j = Cov(X_t, X_{t-j}) = Cov(X_t, X_{t+j})$
Sample autocovariance: $\hat{\gamma}_j = \frac{1}{T} \sum_{t=j+1}^T (X_t - \bar{X})(X_{t-j} - \bar{X})$

- Autocorrelation: $\rho_j = \frac{\gamma_j}{\gamma_0}$

Sample Autocorrelation: $\hat{\rho}_j = \frac{\hat{\gamma}_j}{\hat{\gamma}_0}$

• Partial autocorrelation (PAC)

- PAC ρ_j is a measure of the direct relation between X_t and X_{t-j} for $j = 1, 2, \dots$
- ρ_j is the correlation between X_t and X_{t-j} after controlling for the effects of X_t and $X_{t-1} \dots X_{t-j+1}$
- $\hat{\rho}_1 = \hat{\phi}_{11}$ in $X_t = \phi_{10} + \phi_{11}X_{t-1} + e_{1t}$
- $\hat{\rho}_2 = \hat{\phi}_{21}$ in $X_t = \phi_{20} + \phi_{21}X_{t-1} + \phi_{22}X_{t-2} + e_{2t}, \dots$

Test for autocorrelation

Assignment Project Exam Help

The null hypothesis: H_0 : There is no autocorrelation (White noise process)

- 1 **Autocorrelation test:** $\sqrt{T}\hat{\rho}_j \sim N(0,1)$ under the null hypothesis
Reject if $|\hat{\rho}_j|$ is too large ($> 1.96/\sqrt{T}$, at 5% significance level)

WeChat: cstutorcs

Joint Hypothesis Tests

- We can also test the joint hypothesis that all m of the ρ_k correlation coefficients are simultaneously equal to zero using the Q -statistic developed by Box and Pierce:

$$Q = T \sum_{k=1}^m \hat{\rho}_k^2$$

where T = sample size, m = maximum lag length

- The Q -statistic is asymptotically distributed as a χ_m^2 .
- However, the Box Pierce test has poor small sample properties, so a variant has been developed, called the Ljung-Box statistic:

$$Q^* = T(T+2) \sum_{k=1}^m \frac{\hat{\rho}_k^2}{T-k} \sim \chi_m^2$$

- This statistic is very useful as a portmanteau (general) test of linear dependence in time series.

An ACF Example

- Question:

Suppose that a researcher had estimated the first 5 autocorrelation coefficients using a series of length 100 observations, and found them to be (from 1 to 5): 0.207, -0.013, 0.086, 0.005, -0.022.

Test each of the individual coefficient for significance, and use both the Box-Pierce and Ljung-Box tests to establish whether they are jointly significant.

Solution

A coefficient would be significant if it lies outside $(-0.196, +0.196)$ at the 5% level, so only the first autocorrelation coefficient is significant.

$$Q = 5.09 \text{ and } Q^* = 5.26$$

Compared with a tabulated $\chi^2(5)=11.1$ at the 5% level, so the 5 coefficients are jointly insignificant.

Example: ACF/PACF

- Descriptive statistics

eg. NYSE Composite return

AC test at 5% level:

$$1.96/\sqrt{T} = 0.04462$$

H_0 is rejected at

$$j = 1, 2, 5, 12$$

LB test at 5% level:

H_0 is rejected for

all m , as all p-values

are less than 0.05.

Correlogram of RC					
Sample: 1 1931					
Included observations: 1930					
Autocorrelation	Partial Correlation	J	AC	PAC	Q-Stat Prob
1	0.068	0.068	9.0448	0.003	
2	0.055	0.051	13.106	0.001	
3	-0.031	-0.024	14.940	0.002	
4	-0.001	0.001	14.942	0.005	
5	-0.052	-0.055	20.226	0.001	
6	-0.014	-0.008	20.624	0.002	
7	-0.033	-0.037	22.761	0.002	
8	0.011	0.012	22.992	0.003	
9	0.033	0.028	25.155	0.003	
10	0.028	0.021	26.723	0.003	
11	-0.013	-0.044	30.332	0.001	
12	-0.064	-0.061	35.987	0.000	
13	0.015	0.004	36.403	0.001	
14	-0.013	-0.009	36.720	0.001	
15	-0.002	0.008	36.726	0.001	
16	-0.027	-0.031	38.181	0.001	
17	0.014	0.025	38.588	0.002	
18	-0.007	-0.015	38.677	0.003	
19	0.022	0.026	39.639	0.004	
20	0.006	0.004	39.710	0.005	

Example: ACF/PACF of squared Returns

- Descriptive statistics

Assignment Project Exam Help

What about squared returns?

Usually strongly correlated.


— Why squared returns?

$E(r_t^2) \approx \text{Var}(r_t)$

eg. NYSE Composite

return squared

WeChat: cstutorcs



Summary of stylized Facts

Assignment Project Exam Help

KEY stylised facts about financial return series

- 1 the returns have small, often non-significant autocorrelations (no linear return predictability)
- 2 the squared returns have strong positive autocorrelations (predictability in volatility, volatility clustering)
- 3 large kurtosis (heavy tails, tail probabilities larger than normal)

<https://tutorcs.com>
WeChat: cstutorcs

Summary

Assignment Project Exam Help

- Characterizing Financial time series:
 - asset price and returns
 - stylised facts about index return series
- Normality tests: Z_{ks}, Z_{kt}, LB
- Predictability in returns
 - Autocovariance and autocorrelation
 - Tests for autocorrelation: AC test and Q_m
- Next week: Application of linear regression in Finance (asset pricing)