

**Copyright** ©Copyright University of New South Wales 2020. All rights reserved.

**Course materials subject to Copyright**

UNSW Sydney owns copyright in these materials (unless stated otherwise). The material is subject to copyright under Australian law and overseas under international treaties. The material

are provided for use by enrolled UNSW students. The materials, or any part, may not be copied, shared or distributed, in print or digitally, outside the course without permission. Students may only copy a reasonable portion of the material for personal research or study or for criticism or review. Under no circumstances may these materials be copied or reproduced for sale or commercial purposes without prior written permission of UNSW Sydney.

**Statement on class recording**

To ensure the free and open discussion of ideas, students may not record, by any means, classroom lectures, discussion and/or activities without the advance written permission of the instructor, and any such recording properly approved in advance can be used solely for the student's own private use.

WARNING: Your failure to comply with these conditions may lead to disciplinary action, and may give rise to a civil action or a criminal offence under the law.

THE ABOVE INFORMATION MUST NOT BE REMOVED FROM THIS MATERIAL.

# Assignment Project Exam Help

<https://tutorcs.com>

## WeChat: cstutorcs

# Assignment Project Exam Help

Financial Econometrics

Slides 10: Modeling Return Volatility: Testing/ Estimating/ Forecasting  
ARCH and Introduction to GARCH

<https://tutorcs.com>

Dr. Rachida Ouysse  
School of Economics<sup>1</sup>

<sup>1</sup>©Copyright University of New South Wales 2020. All rights reserved. This copyright notice must not be removed from this material.

WeChat: cstutorcs



## Lecture Plan

# Assignment Project Exam Help

- ARCH LM-test
- Forecasting with ARCH
- Generalised ARCH: why and how
  - Formulation of GARCH: parameter restrictions
- Properties of GARCH(1,1)
  - Mean, variance, ARMA(1,1) representation
  - ML estimation of GARCH
- Forecasting with GARCH

## LM test for ARCH effect

# Assignment Project Exam Help

Testing for ARCH effects: The ARCH-LM test

- Obtain the residuals  $\hat{\mu}_t$  from a regression, e.g.

$$y_t = \beta_1 + \beta_2 x_{2t} + \beta_3 x_{3t} + \beta_4 x_{4t} + \mu_t$$

- Obtain  $R^2$  of the auxiliary regression

$$\hat{\mu}_t^2 = \gamma_0 + \gamma_1 \hat{\mu}_{t-1}^2 + \gamma_2 \hat{\mu}_{t-1}^2 + \dots + \gamma_q \hat{\mu}_{t-q}^2 + \nu_t$$

- Calculate test statistic  $T'R^2$  with  $T'$  the number of observations in the auxiliary regression.
- Under the null hypothesis of no ARCH,  $T'R^2 \sim \chi^2(q)$ .

## ARCH-LM TEST

## LM test for ARCH effect: Example

# Assignment Project Exam Help

eg. NYSE composite return:

- 1 Estimate the model for mean (eg. AR(1)) and save the residual series  $\hat{\mu}_t$ .
- 2 OLS auxiliary regression:  $\hat{\mu}_t^2 = \gamma_0 + \gamma_1 \hat{\mu}_{t-1}^2 + \cdots + \gamma_q \hat{\mu}_{t-q}^2 + error_t$   
Save the  $R^2$ . ( $q$  depends on  $T$  and data frequency)
- 3  $T' = T - q$ , with  $q = 5$  reject when  $T' R^2$  exceeds  $\chi_{(5)}^2$

eg. NYSE composite return: LM test with  $q = 5$

ARCH Test:

F-statistic	37.43273	Probability	0.000000
Obs*R-squared	171.0570	Probability	0.000000

ARCH Test:

F-statistic	1.318328	Probability	0.253338
Obs*R-squared	6.589613	Probability	0.252993

Performed on “V” to check the adequacy of variance equation

WeChat: cstutorcs

## Forecasting with ARCH Models

## Assignment Project Exam Help

- Using repeated substitutions, we can make multi-step forecasts for the return and its volatility
- Example. AR(1)-ARCH(2)

<https://tutorcs.com>

$$y_t = c + \phi_1 y_{t-1} + \mu_t, \mu_t | \Omega_{t-1} \sim N(0, \sigma_t^2)$$

$$\sigma_t^2 = \alpha_0 + \alpha_1 \mu_{t-1}^2 + \alpha_2 \mu_{t-2}^2$$

$$y_{t+1|t} = c + \phi_1 y_t,$$

[WeChat: tutorcs](https://tutorcs.com)

$$y_{t+2|t} = c + \phi_1 y_{t+1|t} + \dots$$

$$\sigma_{t+1|t}^2 = \alpha_0 + \alpha_1 \mu_t^2 + \alpha_2 \mu_{t-1}^2,$$

$$\sigma_{t+2|t}^2 = \alpha_0 + \alpha_1 \sigma_{t+1|t}^2 + \alpha_2 \mu_t^2,$$

$$\sigma_{t+3|t}^2 = \alpha_0 + \alpha_1 \sigma_{t+2|t}^2 + \alpha_2 \sigma_{t+1|t}^2, \dots$$

## Forecasting with ARCH models: Example




## Assignment Project Exam Help

– Forecasting with ARCH models

eg. NYSE composite return:

AR(1)-ARCH(5) forecasts

reverts to unconditional  
(mean reverting)



Remember the limitations of ARCH!

# Assignment Project Exam Help

## Advantages of ARCH

- It is able to capture 'clustering' in return series or the autocorrelation in squared returns.
- It facilitates volatility forecasting.
- It explains partially, non-normality in return series.

## Limitations of ARCH

- ▶ In ARCH( $q$ ), the  $q$  may be selected by AIC, SIC or LR test. The correct value of  $q$  might be very large. The model might not be parsimonious. (eg. ARCH(1) would not work for the composite return)
- ▶ The conditional variance  $\sigma_t^2$  cannot be negative: Requires non-negativity constraints on the coefficients. Sufficient (but not necessary) condition is:  $\alpha_i \geq 0$  for all  $i = 0, 1, 2, \dots, q$ . Especially for large values of  $q$  this might be violated



## GARCH Models: Introduction

# Assignment Project Exam Help

Generalised ARCH (GARCH) models allow the conditional variance to depend upon previous own lags.

- Let  $\mu_t$  be the error term or shock in a model.

$$\text{ARCH}(q): \text{Var}(\mu_t | \Omega_{t-1}) = \sigma_t^2,$$

$$\sigma_t^2 = \alpha_0 + \alpha_1 \mu_{t-1}^2 + \alpha_2 \mu_{t-2}^2 + \cdots + \alpha_q \mu_{t-q}^2,$$

is not parsimonious as a large  $q$  is often required.

- If  $\sigma_{t-1}^2$  is a summary of volatility info in  $\Omega_{t-2}$ , then  $\Omega_{t-1} = \{\mu_{t-1}, \mu_{t-2}, \mu_{t-3}, \dots\} = \{\mu_{t-1}, \sigma_{t-1}^2\}$  (volatility wise!)
- This leads to the GARCH(1,1) model:

$$\sigma_t^2 = \alpha_0 + \alpha_1 \mu_{t-1}^2 + \beta_1 \sigma_{t-1}^2$$

## GARCH: Introduction

## Assignment Project Exam Help

- More generally, GARCH( $p, q$ ) model

$$\text{Var}(\mu_t | \Omega_{t-1}) = \sigma_t^2,$$

$$\sigma_t^2 = \alpha_0 + \alpha_1 \mu_{t-1}^2 + \cdots + \alpha_q \mu_{t-q}^2 + \beta_1 \sigma_{t-1}^2 + \cdots + \beta_p \sigma_{t-p}^2,$$

where the parameters should satisfy:

- (1) Positivity constraint:  $\alpha_0 > 0, \alpha_i \geq 0, \beta_j \geq 0$  for all  $i = 1, \dots, q$  and  $j = 1, \dots, p$
- (2) Finite Variance  $\sum_{i=1}^q \alpha_i + \sum_{j=1}^p \beta_j < 1$ 
  - In practice, the models for asset returns rarely go beyond GARCH(1,1).

## Properties of GARCH

# Assignment Project Exam Help

The generalisation implied by GARCH can be seen from backward iterating the GARCH(1,1) model:

$$\sigma_t^2 = \frac{\alpha_0}{1 - \beta_1} + \alpha_1 \sum_{j=1}^{\infty} \beta_1^{j-1} \mu_{t-j}^2.$$

This shows that the GARCH model is an ARCH( $\infty$ ) with geometrically declining coefficients (for  $|\beta_1| < 1$ ).

## Properties of GARCH(1,1)

## Assignment Project Exam Help

Alternatively, if we define the surprise in the squared innovations as  $\omega_t = \mu_t^2 - \sigma_t^2$ , the GARCH(1,1) model can be rewritten as

$$\begin{aligned}\mu_t^2 - \omega_t &= \alpha_0 + \alpha_1 \mu_{t-1}^2 + \beta_1 (\mu_{t-1}^2 - \omega_{t-1}) \\ \mu_t^2 &= \alpha_0 + (\alpha_1 + \beta_1) \mu_{t-1}^2 + \omega_t - \beta_1 \omega_{t-1}\end{aligned}$$

which shows that the **squared errors** follow an ARMA(1,1) model. As the root of the autoregressive part is  $\alpha_1 + \beta_1$ , the squared residuals are stationary provided  $|\alpha_1 + \beta_1| < 1$ .

Under stationarity  $E(\mu_t^2) = E(\mu_{t-1}^2) = E(\mu_{t-2}^2) = \sigma^2$ , the unconditional variance of  $\mu_t$  is given by

$$\begin{aligned}\sigma^2 &= \alpha_0 + \alpha_1 \sigma^2 + \beta_1 \sigma^2 \\ &= \frac{\alpha_0}{1 - (\alpha_1 + \beta_1)}\end{aligned}$$

## Properties of GARCH(1,1)

# Assignment Project Exam Help

Two general cases can be distinguished

- ▶  $\alpha_1 + \beta_1 < 1 \rightarrow$  the unconditional variance is defined, i.e. finite

<https://tutorcs.com>

- ▶  $\alpha_1 + \beta_1 \geq 1 \rightarrow$  the unconditional variance is not defined, i.e. infinite

The latter case is denoted **non-stationarity in variance**

- ▶ Variance does not converge to an unconditional mean
- ▶ The special case where  $\alpha_1 + \beta_1 = 1$  is known as a unit root in variance or **integrated GARCH** (IGARCH)

WeChat: cstutorcs

## Properties of GARCH(1,1)

## Assignment Project Exam Help

- GARCH(1,1):  $\mu_t | \Omega_{t-1} \sim N(0, \sigma_t^2)$ ,

$$\sigma_t^2 = \alpha_0 + \alpha_1 \mu_{t-1}^2 + \beta_1 \sigma_{t-1}^2$$

$$\alpha_0 > 0 \quad \alpha_1 \geq 0 \quad \beta_1 \geq 0 \quad \alpha_1 + \beta_1 < 1$$

<https://tutores.com>

- Its conditional variance is time varying:

$$E(\mu_t | \Omega_{t-1}) = 0, \quad \text{Var}(\mu_t | \Omega_{t-1}) = \sigma_t^2,$$

$$C(95\%) = E(y_{t+1} | \Omega_{t-1}) \pm 2\sigma_t$$

WeChat: cstutores

- $\mu_t$  is a White Noise:  $E(\mu_t) = 0$ ,  $\text{Var}(\mu_t) = \frac{\alpha_0}{1 - (\alpha_1 + \beta_1)}$ ,  $\text{Cov}(\mu_t, \mu_{t-j}) = 0$
- **But** it is NOT an independent WN or iid WN. It is **NOT unconditionally Normally distributed**:  $\text{kurt}(\mu_t) > 3$

## Properties of GARCH(1,1)

# Assignment Project Exam Help

- GARCH(1,1) can be expressed in terms of **standardised shocks**  $\nu_t$ :

$$\mu_t = \sigma_t^2 \nu_t \text{ and } \nu_t \sim \text{iid } N(0,1)$$

- When model is correct,  $\nu_t^2$  should have no autocorrelation.

### Advantages of the GARCH model (compared to ARCH)

- Avoids overfitting, i.e. a higher order ARCH model may have a more parsimonious GARCH representation
- Due to less estimated parameters, violations of the non-negativity constraint are less likely

## GARCH(1,1) Estimation

# Assignment Project Exam Help

## Estimating GARCH models

For instance, estimate the following AR(1)-GARCH(1,1) model

$$\begin{aligned} y_t &= \mu + \phi y_{t-1} + u_t \\ u_t &= \nu_t \sigma_t, \quad \nu_t \sim N(0, 1) \\ \sigma_t^2 &= \alpha_0 + \alpha_1 \mu_{t-1}^2 + \beta_1 \sigma_{t-1}^2 \end{aligned}$$

**OLS is inappropriate**

OLS minimises the RSS,  $\sum \hat{\mu}_t^2 = \sum (y_t - \hat{\mu} - \hat{\phi} y_{t-1})^2$ , which is a function of the parameters in the conditional mean equation only and not in the conditional variance equation

- ▶ In fact, OLS assumes that the residuals are homoscedastic, i.e. all slope coefficients in the conditional variance equation are set to zero



## GARCH(1,1) Estimation

## Assignment Project Exam Help

## Maximum Likelihood

- ▶ Make assumptions about **conditional distribution** of  $\mu_t$ , e.g.

$$\nu_t \sim N(0, 1) \quad \text{such that} \quad \mu_t \sim N(0, \sigma_t^2)$$

This means that conditional on information available at  $t-1$ ,  $\mu_t$  is normally distributed with mean zero and variance  $\sigma_t^2$  with the latter being known at time  $t-1$ . Note that this does not imply that the **unconditional distribution** of  $\mu_t$  is normal, as  $\sigma_t$  becomes a random variable if we do not condition on all information available on  $t-1$ .

- ▶ The conditional distribution of  $y_t$  is then also normal, given by

$$f(y_t | y_{t-1}, \dots, \mu_{t-1}, \dots) = \frac{1}{\sqrt{2\pi\sigma_t^2}} \exp\left(-\frac{1}{2} \frac{\mu_t^2}{\sigma_t^2}\right)$$

with  $\mu_t = y_t - \mu - \phi y_{t-1}$  and  $\sigma_t^2 = \alpha_0 + \alpha_1 \mu_{t-1}^2 + \beta_1 \sigma_{t-1}^2$ .

## GARCH(1,1) Estimation

# Assignment Project Exam Help


The **loglikelihood function** is given by the sum over all  $t$  of the log of the conditional distribution of  $y_t$

$$L = -\frac{T}{2} \log(2\pi) - \frac{T}{2} \sum_t \log(\sigma_t^2) - \frac{1}{2} \sum_{t=1}^T \frac{\mu_t^2}{\sigma_t^2}$$

- ▶ The ML estimator is obtained by **maximizing** the loglikelihood with respect to the unknown parameters ( $\mu, \phi, \alpha_0, \alpha_1, \beta_1$ ).
- ▶ Analytical solution not possible: use **numerical procedures**
  - ▶ These algorithms 'search' over the parameter space, from an **initial guess**, until a maximum for the loglikelihood function is found
  - ▶ Potential problem: the loglikelihood function may have several **local maxima** such that alternative initial guesses may yield different results.
  - ▶ In practice: use linear regression to get initial estimates of the parameters in the conditional mean equation and choose some (alternative) parameter value for the parameters in the conditional variance equation  $\neq 0$ .

## GARCH(1,1) Estimation

## Assignment Project Exam Help



<https://tutorcs.com>

WeChat: cstutorcs

## GARCH(1,1) Estimation

# Assignment Project Exam Help

- Fortunately, first order conditions are, under some weak assumptions, valid even when  $\nu_t$  is not normally distributed.

- ▶ The parameter estimates are still consistent
- ▶ Adjustments have to be made to the standard errors, i.e. use Bollersley-Woodridge variance-covariance matrix, also known as **Quasi Maximum Likelihood Estimation**, which is robust for non-normality.

<https://tutorcs.com>

WeChat: cstutorcs

— — — — —

## Example 1

## Example 1: GARCH(1,1) Estimation

## Assignment Project Exam Help

eg. NYSE composite return

Dependent Variable: RC  
Method: ML - ARCH (Marquardt) - Normal distribution

Sample (adjusted): 3 1931

Included observations: 1929 after adjustments

Meaningful change after 14 iterations

Bershad-Walding robust standard errors &amp; covariance

Variance inflation factor (VIF)

GARCH = C(1) + C(2)\*RESID(-1)^2

C(1) = 0.072383 C(2) = 0.102468

	Coefficient	Std. Error	z-Statistic	Prob.
C	0.072383	0.018494	3.913969	0.0001
AR(1)	0.102468	0.025514	4.016133	0.0001

## Variance Equation

	Coefficient	Std. Error	z-Statistic	Prob.
C	0.013305	0.004529	2.937750	0.0033
RESID(-1)^2	0.119181	0.024748	4.815836	0.0000
GARCH-1	0.877598	0.021117	41.71302	0.0000

R-squared	0.0004	Mean dependent variable	0.00516
Adjusted R-squared	0.00034	S.D. dependent variable	1.00945
S.E. of regression	0.289	Akaike information criterion	2.230091
Sum squared resid	1210.240	Schwarz criterion	2.230091
Log likelihood	-2523.598	F-statistic	1.164594
Durbin-Watson stat	2.055220	Prob(F-statistic)	0.324498

Inverted AR Roots	10
-------------------	----

ARCH Test: q = 5

F-statistic	0.764181	Probability	0.575599
Obs*R-squared	3.825239	Probability	0.574842

Test Equation:  
Dependent Variable: STD\_RESID\*2


## Correlogram of Standardized Residuals Squared

Sample: 3 1931

Included observations: 1929

Q-statistic probabilities adjusted for 1 ARMA term(s)

Autocorrelation	Partial Correlation	AC	PAC	Q-Stat	Prob
1	1	-0.001	-0.001	0.0005	
2	0.36	0.036	0.036	2.4991	0.114
3	-0.14	-0.014	-0.014	2.8675	0.238
4	-0.11	-0.012	-0.012	3.1058	0.376
5	-0.20	-0.019	-0.019	3.8964	0.420
6	0.009	0.010	0.031	4.0531	0.542
7	0.003	0.004	0.068	4.0658	0.668
8	-0.001	-0.002	0.0670	4.0670	0.772
9	-0.011	-0.011	4.3009	0.629	
10	-0.010	-0.010	4.4826	0.877	
11	-0.008	-0.006	4.5929	0.917	
12	-0.016	-0.016	5.1005	0.526	
13	-0.004	-0.004	5.1268	0.954	
14	-0.011	-0.011	5.3589	0.966	
15	-0.019	-0.019	6.0452	0.965	
16	-0.009	-0.009	6.2090	0.976	



Series: Standardized Residuals

Sample 3 1931

Observations 1929

Mean	-0.048341
Median	-0.039867
Maximum	2.850528
Minimum	-6.601836
Std. Dev.	0.996820
Skewness	-0.547486
Kurtosis	4.973199
Jarque-Bera	409.3080
Probability	0.000000

## Example 1

## Example 1: GARCH(1,1) Estimation

## Assignment Project Exam Help

— ML Estimation of GARCH(1,1)

eg. NYSE composite return (continued)

Large  $\beta_1$  estimate: about 0.9

Small  $\alpha_1$  estimate about 0.1

$\alpha_1 + \beta_1$  estimate: very close to 1

<https://tutorcs.com>

GARCH(1,1) is preferred by AIC and SIC.

WeChat: cstutorcs

	AIC	SIC
AR(1)-ARCH(5)	2.664	2.687
AR(1)-GARCH(1,1)	2.622	2.636

## Example 1

## Example 1: GARCH(1,1) Estimation

# Assignment Project Exam Help

ML Estimation of GARCH(1,1)

eg. NYSE composite return (continued)

GARCH(1,1)  $\sigma_t$  plot is smoother than ARCH(5).



Large  $\beta_1$  estimate implies *persistence*:

$\sigma_t$  tends to continue at the current level.

<https://tutorcs.com>

$$\sigma_t^2 = \alpha_0 + \alpha_1 \varepsilon_{t-1}^2 + \beta_1 \sigma_{t-1}^2$$

WeChat: cstutorcs



## Summary facts about GARCH models

# Assignment Project Exam Help

- GARCH(1,1) is usually preferred to ARCH or higher order GARCH, because of its parsimony.
- Usually, GARCH  $\beta_1$  estimate is about 0.9 or more and  $\alpha_1 + \beta_1$  estimate is very close to 1, for daily returns.
- Standardised residuals are usually non-normal, with negative skewness and excessive kurtosis.
- GARCH(1,1) is able to capture clustering in returns but unable to account for  
Asymmetry: negative returns tend to cause more volatility;  
Non-normality; Structural change
- Coefficient restrictions are hard to impose in MLE