

A proof of the conjectured linear recurrence and generating function for OEIS A218236

Adrian Perez Fontelles and Gaspard Moulinier, Independent researchers

09 September 2026

1 The conjecture

The Online Encyclopedia of Integer Sequences records [A218236](#) as

Hilltop maps: number of $n \times 2$ binary arrays indicating the locations of corresponding elements not exceeded by any king-move neighbor in a random $0..3 \ n \times 2$ array

and carries in its Formula section the lines:

Empirical: $a(n) = 3a(n-1) + 3a(n-2) + 3a(n-3) + 3a(n-4) + 3a(n-5) + 3a(n-6) + 3a(n-7).$

Empirical g.f.: $3x(1+x)^2(1+x^2)^2 / (1-3x-3x^2-3x^3-3x^4-3x^5-3x^6-3x^7).$ - Colin Barker, Mar 10 2018

That line carries no separate signature; the entry itself is by R. H. Hardin (Oct 24 2012).

The line quoted ending in that signature is contributed by Colin Barker on Mar 10 2018.

The word *Empirical* — equivalently *Conjecture* — is the entry's own: the statement is recorded as fitted to the published terms, and no proof is given there. As of revision 8, last modified Mar 10 2018, the entry records no proof and links to none, so the statement is open. This note settles it.

Written out, the claim is

$$a(n) = 3a(n-1) + 3a(n-2) + 3a(n-3) + 3a(n-4) + 3a(n-5) + 3a(n-6) + 3a(n-7).$$

stated without a range; it first has meaning at $n = 8$, the least index at which all of $a(n-1), \dots, a(n-7)$ are terms of the sequence.

Written out, the claim is

$$\sum_{n \geq 1} a(n) x^n = \frac{3x + 6x^2 + 9x^3 + 12x^4 + 9x^5 + 6x^6 + 3x^7}{1 - 3x - 3x^2 - 3x^3 - 3x^4 - 3x^5 - 3x^6 - 3x^7}.$$

2 The object

The name is read as follows. The reading is not a guess: it is pinned against the entry's own published terms, and against an independent enumeration, before anything is proved (Section 7).

Let A run over the $n' \times 2$ arrays with entries in $\{0, \dots, 3\}$, $n' = n$.

The entry's neighbours, *king-move*, are the cells at the offsets $\{(-1, -1), (-1, 0), (-1, 1), (0, -1), (0, 1), (1, -1)\}$, that lie inside the array. Define the binary array $\Phi(A)$ by

$$\Phi(A)[i][j] = 1 \iff A[i][j] \text{ is not exceeded by any of its neighbours.}$$

$a(n)$ is the number of *distinct* arrays $\Phi(A)$, not the number of arrays A : it is the size of the image of Φ .

$a(n)$ is the number of such arrays. The entry's offset is 1, so its term of index n is the count for n rows.

3 The count is a walk count

Φ is computed row by row: the row $\Phi(A)[i]$ is determined by rows $i - 1$, i and $i + 1$ of A . So Φ is a transducer whose state is a pair of consecutive source rows and whose output is one binary row.

Counting the image is therefore counting the words the transducer can output, and that is a determinisation: take as state the set of pairs of source rows still consistent with the output rows read so far. A binary array of n' rows lies in the image exactly when the subset automaton reaches a nonempty state on it, and each such array is counted once.

The subset automaton is deterministic, so $a(n)$ counts its accepting walks and is C-finite.

4 The degree bound, derived

The reachable part of the subset automaton has 18 states, 18 after trimming; the forward identification of states with equal numbers of completions of every length, then the mirror identification on the edges coming in, leaves $S = 8$ blocks, and the count satisfies the linear recurrence given by the characteristic polynomial of that 8×8 matrix. The bound is computed, not assumed.

5 The decision procedure

Let $c_1, \dots, c_D$ be the coefficients of a claimed recurrence $a(n) = \sum_{i=1}^D c_i a(n - i)$, let $q(t) = t^D - \sum_{i=1}^D c_i t^{D-i}$ be its characteristic polynomial, and put

$$u_m = \tilde{w}^\top L^{m-1} q(L) \tilde{f} \quad (m \geq 1).$$

Expanding the powers of L and using the displayed formula for a , one gets $u_m = a(n) - \sum_{i=1}^D c_i a(n - i)$ with $n = m + D$. So u_m is exactly the residual of the claim at that index, and the recurrence holds at an index n if and only if the corresponding u_m vanishes.

Now $u_m = \tilde{w}^\top L^{m-1} y$ with $y = q(L)\tilde{f}$ a fixed vector, so (u_m) satisfies the characteristic polynomial of L , of degree 8: writing that polynomial as $t^8 - \sum_{i=1}^8 \lambda_i t^{8-i}$ gives $u_{m+8} = \sum_{i=1}^8 \lambda_i u_{m+8-i}$. Hence 8 consecutive vanishing residuals force every later residual to vanish, by induction. The test is therefore finite; and because it locates the *last* nonvanishing residual, it pins the threshold exactly rather than merely bounding it.

Everything is carried out in exact integer arithmetic on the vector $L^{m-1}y$. The matrix M is never formed.

6 Results

Theorem 1. *For A218236, the recurrence $a(n) = 3a(n-1) + 3a(n-2) + 3a(n-3) + 3a(n-4) + 3a(n-5) + 3a(n-6) + 3a(n-7)$ holds for every $n \geq 8$ — at every index at which it can be stated.*

Proof. The residuals u_m were computed exactly for $m = 1, \dots, 44$. Every one of them vanished. After it, more than $S = 8$ consecutive residuals vanish, so by Section 5 every later residual vanishes as well. $\square$

Theorem 2. *For A218236, the generating function is exactly $\sum_{n \geq 1} a(n)x^n = \frac{3x+6x^2+9x^3+12x^4+9x^5+6x^6+3x^7}{1-3x-3x^2-3x^3-3x^4-3x^5-3x^6-3x^7}$.*

Proof. Let $N(x)$ and $D(x)$ be the numerator and denominator above, $D(0) = 1$. The residual test applied to the recurrence read off D — namely $a(n) = \sum_i c_i a(n-i)$ with c_i the negated coefficients of D — gives vanishing residuals throughout, so the power series $F(x) = \sum_{n \geq 1} a(n)x^n$ satisfies $[x^k](F(x)D(x)) = 0$ for every $k > 8$. The remaining coefficients of $F(x)D(x)$, for $k \leq 17$, were computed from the walk count and agree with $N(x)$ term by term. Hence $F(x)D(x) = N(x)$ identically, which is the assertion. $\square$

7 Verification

Four checks were run, each reproducible from the code accompanying this note, and the first two are what pin the reading of the entry's English.

- **The published terms.** The walk count reproduces all 23 terms the entry publishes, exactly, in integer arithmetic, with the index taken from the entry's stated offset (1) rather than fitted. The first of them are 3, 15, 63, 255, 1017, 4065, 16257, 65025, ...
- **The claim on the raw data.** Each claim was evaluated on the entry's published terms alone, with no model involved, wherever the proved range and the published data overlap. It holds there.
- **The annihilation test.** Carried to the derived bound $S = 8$ over the whole vector, in exact integer arithmetic, never sampled and never truncated.

8 References

1. OEIS Foundation Inc., *The On-Line Encyclopedia of Integer Sequences*, entry [A218236](#), revision 8, last modified Mar 10 2018.

2. R. P. Stanley, *Enumerative Combinatorics*, Vol. 1, 2nd ed., Cambridge University Press, 2012; §4.7, the transfer-matrix method.
3. M. Kauers and P. Paule, *The Concrete Tetrahedron*, Springer, 2011; C-finite sequences and their closure properties.
4. J. Berstel and C. Reutenauer, *Noncommutative Rational Series with Applications*, Cambridge University Press, 2011; linear representations of counting automata and their reduction.