

A proof of the conjectured linear recurrence for OEIS A263367

Adrian Perez Fontelles and Gaspard Moulinier, Independent researchers

09 September 2026

1 The conjecture

The Online Encyclopedia of Integer Sequences records [A263367](#) as

Number of $(n+1) \times (2+1)$ 0..2 arrays with each row and column divisible by 7, read as a base-3 number with top and left being the most significant digits

and carries in its Formula section the line:

Empirical: $a(n) = 4*a(n-1) + 101*a(n-6) - 404*a(n-7) - 2159*a(n-12) + 8636*a(n-13) + 2059*a(n-18) - 8236*a(n-19)$

That line carries no separate signature; the entry itself is by R. H. Hardin (Oct 16 2015).

The word *Empirical* — equivalently *Conjecture* — is the entry's own: the statement is recorded as fitted to the published terms, and no proof is given there. As of revision 4, last modified Oct 16 2015, the entry records no proof and links to none, so the statement is open. This note settles it.

Written out, the claim is

$$a(n) = 4a(n-1) + 101a(n-6) - 404a(n-7) - 2159a(n-12) + 8636a(n-13) + 2059a(n-18) - 8236a(n-19).$$

stated without a range; it first has meaning at $n = 20$, the least index at which all of $a(n-1), \dots, a(n-19)$ are terms of the sequence.

2 The object

The name is read as follows. The reading is not a guess: it is pinned against the entry's own published terms, and against an independent enumeration, before anything is proved (Section 7).

Let A be an $n' \times 3$ array over $\{0, \dots, 2\}$ with $n' = n + 1$ rows. Each row is read as a base-3 number with its leftmost entry the most significant digit, and each column as a base-3 number with its topmost entry the most significant digit.

The condition is that every row is divisible by 7 and every column is divisible by 7.

$a(n)$ is the number of such arrays. The entry's offset is 1, so its term of index n is the count for n rows.

3 The count is a walk count

Reading a column downwards multiplies its value so far by 3 and adds the new digit, so the residue of a column modulo 7 after i rows determines its residue after $i + 1$ rows once the new digit is known. Take as state the vector of the 3 column residues modulo 7; there are at most 7^3 of them.

The rows that may be appended are exactly those whose own value is divisible by 7 — a condition on the row alone — and a state is accepting when every column residue is zero.

An array of n' rows is then exactly a walk of length n' from the all-zero state to an accepting state, and every array arises from exactly one walk, so the count is $w^\top M^{n'} f$ and is C-finite.

4 The degree bound, derived

The reachable part of the digraph has 49 states, 49 after trimming; identifying states with equal numbers of completions of every length, and then the mirror identification on the edges coming in, leaves $S = 49$ blocks. The count satisfies the linear recurrence given by the characteristic polynomial of the resulting 49×49 matrix. That bound is computed, not assumed.

5 The decision procedure

Let $c_1, \dots, c_D$ be the coefficients of a claimed recurrence $a(n) = \sum_{i=1}^D c_i a(n - i)$, let $q(t) = t^D - \sum_{i=1}^D c_i t^{D-i}$ be its characteristic polynomial, and put

$$u_m = \tilde{w}^\top L^{m-1} q(L) \tilde{f} \quad (m \geq 1).$$

Expanding the powers of L and using the displayed formula for a , one gets $u_m = a(n) - \sum_{i=1}^D c_i a(n - i)$ with $n = m + D$. So u_m is exactly the residual of the claim at that index, and the recurrence holds at an index n if and only if the corresponding u_m vanishes.

Now $u_m = \tilde{w}^\top L^{m-1} y$ with $y = q(L) \tilde{f}$ a fixed vector, so (u_m) satisfies the characteristic polynomial of L , of degree 49: writing that polynomial as $t^{49} - \sum_{i=1}^{49} \lambda_i t^{49-i}$ gives $u_{m+49} = \sum_{i=1}^{49} \lambda_i u_{m+49-i}$. Hence 49 consecutive vanishing residuals force every later residual to vanish, by induction. The test is therefore finite; and because it locates the *last* nonvanishing residual, it pins the threshold exactly rather than merely bounding it.

Everything is carried out in exact integer arithmetic on the vector $L^{m-1} y$. The matrix M is never formed.

6 Results

Theorem 1. *For A263367, the recurrence $a(n) = 4a(n - 1) + 101a(n - 6) - 404a(n - 7) - 2159a(n - 12) + 8636a(n - 13) + 2059a(n - 18) - 8236a(n - 19)$ holds for every $n \geq 20$ — at every index at which it can be stated.*

Proof. The residuals u_m were computed exactly for $m = 1, \dots, 99$. Every one of them vanished. After it, more than $S = 49$ consecutive residuals vanish, so by Section 5 every later residual vanishes as well. $\square$

7 Verification

Four checks were run, each reproducible from the code accompanying this note, and the first two are what pin the reading of the entry's English.

- **The published terms.** The walk count reproduces all 25 terms the entry publishes, exactly, in integer arithmetic, with the index taken from the entry's stated offset (1) rather than fitted. The first of them are 1, 2, 7, 29, 112, 367, 1359, 5370, ...
- **An independent enumeration.** For $n = 1, \dots, 3$ every one of the 3^{3n} arrays was written out and tested cell by cell against the condition as the entry states it — no transfer matrix, no row window, a separate piece of code. The counts agree with the walk count and with the entry.
- **The claim on the raw data.** Each claim was evaluated on the entry's published terms alone, with no model involved, wherever the proved range and the published data overlap. It holds there.
- **The annihilation test.** Carried to the derived bound $S = 49$ over the whole vector, in exact integer arithmetic, never sampled and never truncated.

8 References

1. OEIS Foundation Inc., *The On-Line Encyclopedia of Integer Sequences*, entry [A263367](#), revision 4, last modified Oct 16 2015.
2. R. P. Stanley, *Enumerative Combinatorics*, Vol. 1, 2nd ed., Cambridge University Press, 2012; §4.7, the transfer-matrix method.
3. M. Kauers and P. Paule, *The Concrete Tetrahedron*, Springer, 2011; C-finite sequences and their closure properties.
4. J. Berstel and C. Reutenauer, *Noncommutative Rational Series with Applications*, Cambridge University Press, 2011; linear representations of counting automata and their reduction.