

# A proof of the conjectured linear recurrence and generating function for OEIS A269078

Adrian Perez Fontelles and Gaspard Moulinier, Independent researchers

09 September 2026

## 1 The conjecture

The Online Encyclopedia of Integer Sequences records [A269078](#) as

*Number of  $4 \times n$  binary arrays with some element plus some horizontally, diagonally or antidiagonally adjacent neighbor totalling two not more than once*

and carries in its Formula section the lines:

Empirical:  $a(n) = 4a(n-1) + 28a(n-2) - 78a(n-3) - 264a(n-4) + 296a(n-5) + 527a(n-6) - 252a(n-7) - 324a(n-8)$ .

Empirical g.f.:  $x(16 + 25x - 267x^2 - 104x^3 + 691x^4 + 275x^5 - 576x^6 - 324x^7) / (1 - 2x - 16x^2 + 7x^3 + 18x^4)^2$ . - Colin Barker, Jan 19 2019

That line carries no separate signature; the entry itself is by R. H. Hardin (Feb 19 2016).

The line quoted ending in that signature is contributed by Colin Barker on Jan 19 2019.

The word *Empirical* — equivalently *Conjecture* — is the entry's own: the statement is recorded as fitted to the published terms, and no proof is given there. As of revision 8, last modified Jan 19 2019, the entry records no proof and links to none, so the statement is open. This note settles it.

Written out, the claim is

$$a(n) = 4a(n-1) + 28a(n-2) - 78a(n-3) - 264a(n-4) + 296a(n-5) + 527a(n-6) - 252a(n-7) - 324a(n-8).$$

stated without a range; it first has meaning at  $n = 9$ , the least index at which all of  $a(n-1), \dots, a(n-8)$  are terms of the sequence.

Written out, the claim is

$$\sum_{n \geq 1} a(n) x^n = \frac{16x + 25x^2 - 267x^3 - 104x^4 + 691x^5 + 275x^6 - 576x^7 - 324x^8}{1 - 4x - 28x^2 + 78x^3 + 264x^4 - 296x^5 - 527x^6 + 252x^7 + 324x^8}.$$

## 2 The object

The name is read as follows. The reading is not a guess: it is pinned against the entry's own published terms, and against an independent enumeration, before anything is proved (Section 7).

Let  $A$  be an  $n' \times 4$  array over  $\{0, \dots, 1\}$  with  $n' = n$  rows, the entry's array transposed so the growing direction is downwards, the adjacency offsets transposed with it.

The entry's *horizontally, diagonally or antidiagonally* adjacency is used once per pair, so the offsets taken are

$$\mathcal{D} = \{(1, -1), (1, 0), (1, 1)\}.$$

Let

$$N(A) = \#\{((i, j), (i+\delta, j+\varepsilon)) : (\delta, \varepsilon) \in \mathcal{D}, \text{ both cells inside the array, } A[i][j] + A[i+\delta][j+\varepsilon] = 2\}.$$

The condition is  $N(A) \leq 1$ .

$a(n)$  is the number of such arrays. The entry's offset is 1, so its term of index  $n$  is the count for  $n$  columns.

## 3 The count is a walk count

The quantity counted is global, but only its value up to 1 matters, so it can be carried in the state capped at 2. Every offset moves at most one row, so a pair is detected when the lower of its two cells is read. Taking as state the last row together with that capped counter makes the whole condition decidable one row at a time.

An array of  $n'$  rows is then exactly a walk of length  $n'$  from the empty state to a state whose counter satisfies the bound, and every array arises from exactly one walk, so  $a(n) = w^\top M^{n'} f$  and the count is C-finite.

## 4 The degree bound, derived

The reachable part of the digraph has 49 states, 33 after trimming; the forward identification of states with equal numbers of completions of every length, followed by the mirror identification on the edges coming in, leaves  $S = 10$  blocks. The count satisfies the linear recurrence given by the characteristic polynomial of that  $10 \times 10$  matrix, a bound computed here rather than assumed.

## 5 The decision procedure

Let  $c_1, \dots, c_D$  be the coefficients of a claimed recurrence  $a(n) = \sum_{i=1}^D c_i a(n-i)$ , let  $q(t) = t^D - \sum_{i=1}^D c_i t^{D-i}$  be its characteristic polynomial, and put

$$u_m = \tilde{w}^\top L^{m-1} q(L) \tilde{f} \quad (m \geq 1).$$

Expanding the powers of  $L$  and using the displayed formula for  $a$ , one gets  $u_m = a(n) - \sum_{i=1}^D c_i a(n-i)$  with  $n = m + D$ . So  $u_m$  is exactly the residual of the claim at that index, and the recurrence holds at an index  $n$  if and only if the corresponding  $u_m$  vanishes.

Now  $u_m = \tilde{w}^\top L^{m-1} y$  with  $y = q(L)\tilde{f}$  a fixed vector, so  $(u_m)$  satisfies the characteristic polynomial of  $L$ , of degree 10: writing that polynomial as  $t^{10} - \sum_{i=1}^{10} \lambda_i t^{10-i}$  gives  $u_{m+10} = \sum_{i=1}^{10} \lambda_i u_{m+10-i}$ . Hence 10 consecutive vanishing residuals force every later residual to vanish, by induction. The test is therefore finite; and because it locates the *last* nonvanishing residual, it pins the threshold exactly rather than merely bounding it.

Everything is carried out in exact integer arithmetic on the vector  $L^{m-1}y$ . The matrix  $M$  is never formed.

## 6 Results

**Theorem 1.** *For A269078, the recurrence  $a(n) = 4a(n-1) + 28a(n-2) - 78a(n-3) - 264a(n-4) + 296a(n-5) + 527a(n-6) - 252a(n-7) - 324a(n-8)$  holds for every  $n \geq 9$  — at every index at which it can be stated.*

*Proof.* The residuals  $u_m$  were computed exactly for  $m = 1, \dots, 44$ . Every one of them vanished. After it, more than  $S = 10$  consecutive residuals vanish, so by Section 5 every later residual vanishes as well.  $\square$

**Theorem 2.** *For A269078, the generating function is exactly  $\sum_{n \geq 1} a(n)x^n = \frac{16x + 25x^2 - 267x^3 - 104x^4 + 691x^5 + 275x^6 - 57x^7 - 14x^8 - 28x^9 + 78x^{10} + 264x^{11} - 296x^{12} - 527x^{13} + 252x^{14} + 324x^{15}}{1 - 4x - 28x^2 + 78x^3 + 264x^4 - 296x^5 - 527x^6 + 252x^7 + 324x^8}$ .*

*Proof.* Let  $N(x)$  and  $D(x)$  be the numerator and denominator above,  $D(0) = 1$ . The residual test applied to the recurrence read off  $D$  — namely  $a(n) = \sum_i c_i a(n-i)$  with  $c_i$  the negated coefficients of  $D$  — gives vanishing residuals throughout, so the power series  $F(x) = \sum_{n \geq 1} a(n)x^n$  satisfies  $[x^k](F(x)D(x)) = 0$  for every  $k > 9$ . The remaining coefficients of  $F(x)D(x)$ , for  $k \leq 19$ , were computed from the walk count and agree with  $N(x)$  term by term. Hence  $F(x)D(x) = N(x)$  identically, which is the assertion.  $\square$

## 7 Verification

Four checks were run, each reproducible from the code accompanying this note, and the first two are what pin the reading of the entry's English.

- **The published terms.** The walk count reproduces all 20 terms the entry publishes, exactly, in integer arithmetic, with the index taken from the entry's stated offset (1) rather than fitted. The first of them are 16, 89, 537, 3288, 17713, 102545, 542112, 2991561, ...
- **An independent enumeration.** For  $n = 1, \dots, 4$  every one of the  $2^{4n}$  arrays was written out and tested cell by cell against the condition as the entry states it — no transfer matrix, no row window, a separate piece of code. The counts agree with the walk count and with the entry.
- **The claim on the raw data.** Each claim was evaluated on the entry's published terms alone, with no model involved, wherever the proved range and the published data overlap. It holds there.

- **The annihilation test.** Carried to the derived bound  $S = 10$  over the whole vector, in exact integer arithmetic, never sampled and never truncated.

## 8 References

1. OEIS Foundation Inc., *The On-Line Encyclopedia of Integer Sequences*, entry [A269078](#), revision 8, last modified Jan 19 2019.
2. R. P. Stanley, *Enumerative Combinatorics*, Vol. 1, 2nd ed., Cambridge University Press, 2012; §4.7, the transfer-matrix method.
3. M. Kauers and P. Paule, *The Concrete Tetrahedron*, Springer, 2011; C-finite sequences and their closure properties.
4. J. Berstel and C. Reutenauer, *Noncommutative Rational Series with Applications*, Cambridge University Press, 2011; linear representations of counting automata and their reduction.