

A proof of the conjectured linear recurrence for OEIS A274747

Adrian Perez Fontelles and Gaspard Moulinier, Independent researchers

09 September 2026

1 The conjecture

The Online Encyclopedia of Integer Sequences records [A274747](#) as

Number of $n \times 6$ 0..2 arrays with no element equal to any value at offset $(-1, -2)$ $(0, -1)$ or $(-1, 0)$ and new values introduced in order 0..2

and carries in its Formula section the line:

Empirical: $a(n) = 14*a(n-1) - 10*a(n-2) - 80*a(n-3) + 177*a(n-4) - 152*a(n-5) + 56*a(n-6) - a(n-7) - 5*a(n-8) + a(n-9)$ for $n > 11$

That line carries no separate signature; the entry itself is by R. H. Hardin (Jul 04 2016).

The word *Empirical* — equivalently *Conjecture* — is the entry's own: the statement is recorded as fitted to the published terms, and no proof is given there. As of revision 4, last modified Jul 04 2016, the entry records no proof and links to none, so the statement is open. This note settles it.

Written out, the claim is

$$a(n) = 14a(n-1) - 10a(n-2) - 80a(n-3) + 177a(n-4) - 152a(n-5) + 56a(n-6) - a(n-7) - 5a(n-8) + a(n-9).$$

stated for $n > 11$.

2 The object

The name is read as follows. The reading is not a guess: it is pinned against the entry's own published terms, and against an independent enumeration, before anything is proved (Section 7).

Let A be an $n' \times 6$ array over $\{0, \dots, 2\}$ with $n' = n$ rows.

The entry's offsets are $(-1, -2)$, $(0, -1)$, $(-1, 0)$. The condition is

$$A[i][j] \neq A[i+\delta][j+\varepsilon] \quad \text{for every listed } (\delta, \varepsilon) \text{ landing inside the array.}$$

An offset and its negative say the same thing, so the offsets are normalised here to point upwards or, in the same row, leftwards: $(-1,-2)$, $(-1,0)$, $(0,-1)$.

The entry's further clause, that the new values $0, \dots, 2$ are introduced in order, means that reading the array in row-major order the first occurrences of the values happen in increasing order: a value v may appear only once every smaller value has already appeared.

$a(n)$ is the number of such arrays. The entry's offset is 1, so its term of index n is the count for n rows.

3 The count is a walk count

Every offset reaches at most 1 row upwards, so the condition on a row is decided by that row together with the 1 rows above it. Take as state the last 1 rows, together with a counter recording how many of the values $0, \dots, 2$ have already been introduced — that counter is all the labelling clause needs, since a value may be used exactly when it is smaller than the counter, or equal to it, in which case the counter advances.

Appending a row is a transition, an array of n' rows is a walk of length n' from the empty state, and every array arises from exactly one walk. So $a(n) = w^\top M^{n'} f$ and the count is C-finite.

4 The degree bound, derived

The reachable part of the digraph has 99 states, 99 after trimming. Identifying states with the same number of completions of every length, and then the mirror identification on the edges coming in, leaves $S = 17$ blocks, and the count satisfies the linear recurrence given by the characteristic polynomial of the resulting 17×17 matrix. That bound is computed, not assumed.

5 The decision procedure

Let $c_1, \dots, c_D$ be the coefficients of a claimed recurrence $a(n) = \sum_{i=1}^D c_i a(n-i)$, let $q(t) = t^D - \sum_{i=1}^D c_i t^{D-i}$ be its characteristic polynomial, and put

$$u_m = \tilde{w}^\top L^{m-1} q(L) \tilde{f} \quad (m \geq 1).$$

Expanding the powers of L and using the displayed formula for a , one gets $u_m = a(n) - \sum_{i=1}^D c_i a(n-i)$ with $n = m + D$. So u_m is exactly the residual of the claim at that index, and the recurrence holds at an index n if and only if the corresponding u_m vanishes.

Now $u_m = \tilde{w}^\top L^{m-1} y$ with $y = q(L) \tilde{f}$ a fixed vector, so (u_m) satisfies the characteristic polynomial of L , of degree 17: writing that polynomial as $t^{17} - \sum_{i=1}^{17} \lambda_i t^{17-i}$ gives $u_{m+17} = \sum_{i=1}^{17} \lambda_i u_{m+17-i}$. Hence 17 consecutive vanishing residuals force every later residual to vanish, by induction. The test is therefore finite; and because it locates the *last* nonvanishing residual, it pins the threshold exactly rather than merely bounding it.

Everything is carried out in exact integer arithmetic on the vector $L^{m-1} y$. The matrix M is never formed.

6 Results

Theorem 1. *For A274747, the recurrence $a(n) = 14a(n-1) - 10a(n-2) - 80a(n-3) + 177a(n-4) - 152a(n-5) + 56a(n-6) - a(n-7) - 5a(n-8) + a(n-9)$ holds for every $n > 11$, and fails at $n = 11$.*

Proof. The residuals u_m were computed exactly for $m = 1, \dots, 48$. The last nonvanishing one is at $m = 2$, that is at $n = 11$. After it, more than $S = 17$ consecutive residuals vanish, so by Section 5 every later residual vanishes as well. $\square$

The entry claims the recurrence for $n > 11$. The theorem is at least as strong.

7 Verification

Four checks were run, each reproducible from the code accompanying this note, and the first two are what pin the reading of the entry's English.

- **The published terms.** The walk count reproduces all 16 terms the entry publishes, exactly, in integer arithmetic, with the index taken from the entry's stated offset (1) rather than fitted. The first of them are 16, 164, 2017, 25585, 327123, 4190533, 53680592, 687685512, $\dots$
- **An independent enumeration.** For $n = 1, \dots, 1$ every one of the 3^{6n} arrays was written out and tested cell by cell against the condition as the entry states it — no transfer matrix, no row window, a separate piece of code. The counts agree with the walk count and with the entry.
- **The claim on the raw data.** Each claim was evaluated on the entry's published terms alone, with no model involved, wherever the proved range and the published data overlap. It holds there.
- **The annihilation test.** Carried to the derived bound $S = 17$ over the whole vector, in exact integer arithmetic, never sampled and never truncated.

8 References

1. OEIS Foundation Inc., *The On-Line Encyclopedia of Integer Sequences*, entry [A274747](#), revision 4, last modified Jul 04 2016.
2. R. P. Stanley, *Enumerative Combinatorics*, Vol. 1, 2nd ed., Cambridge University Press, 2012; §4.7, the transfer-matrix method.
3. M. Kauers and P. Paule, *The Concrete Tetrahedron*, Springer, 2011; C-finite sequences and their closure properties.
4. J. Berstel and C. Reutenauer, *Noncommutative Rational Series with Applications*, Cambridge University Press, 2011; linear representations of counting automata and their reduction.