

A proof of the conjectured linear recurrence and generating function for OEIS A280309

Adrian Perez Fontelles and Gaspard Moulinier, Independent researchers

09 September 2026

1 The conjecture

The Online Encyclopedia of Integer Sequences records [A280309](#) as

Number of $n \times 1$ 0..2 arrays with no element equal to more than one of its horizontal and vertical neighbors, with the exception of exactly two elements, and with new values introduced in order 0 sequentially upwards

and carries in its Formula section the lines:

Empirical: $a(n) = 6a(n-1) - 6a(n-2) - 16a(n-3) + 12a(n-4) + 24a(n-5) + 8a(n-6)$ for $n > 10$.

Empirical g.f.: $x^4(1 - 4x + 3x^2 + 8x^3 - 6x^4 - 12x^5 - 4x^6) / (1 - 2x - 2x^2)^3$. - Colin Barker, Feb 19 2018

That line carries no separate signature; the entry itself is by R. H. Hardin (Dec 31 2016).

The line quoted ending in that signature is contributed by Colin Barker on Feb 19 2018.

The word *Empirical* — equivalently *Conjecture* — is the entry's own: the statement is recorded as fitted to the published terms, and no proof is given there. As of revision 12, last modified Feb 19 2018, the entry records no proof and links to none, so the statement is open. This note settles it.

Written out, the claim is

$$a(n) = 6a(n-1) - 6a(n-2) - 16a(n-3) + 12a(n-4) + 24a(n-5) + 8a(n-6).$$

stated for $n > 10$.

Written out, the claim is

$$\sum_{n \geq 1} a(n) x^n = \frac{x^4 - 4x^5 + 3x^6 + 8x^7 - 6x^8 - 12x^9 - 4x^{10}}{1 - 6x + 6x^2 + 16x^3 - 12x^4 - 24x^5 - 8x^6}.$$

2 The object

The name is read as follows. The reading is not a guess: it is pinned against the entry's own published terms, and against an independent enumeration, before anything is proved (Section 7).

Let A be an $n' \times 1$ array over $\{0, \dots, 2\}$ with $n' = n$ rows.

The entry's neighbours, *horizontal and vertical*, are the cells at the offsets

$$\mathcal{N} = \{(-1, 0), (0, -1), (0, 1), (1, 0)\}$$

that lie inside the array, and $c(i, j)$ is the number of them carrying the same value as (i, j) .

Call a cell *bad* when $c(i, j) > 1$.

The condition is that exactly 2 cells are bad.

The entry's further clause says that reading the array row by row the values $0, \dots, 2$ first occur in increasing order.

$a(n)$ is the number of such arrays. The entry's offset is 1, so its term of index n is the count for n rows.

3 The count is a walk count

Every neighbour offset moves at most one row, so whether a cell of row i is bad is decided by rows $i - 1$, i and $i + 1$. Take as state the last two rows, the number of bad cells so far capped at 3, and a counter for how many values have been introduced. Appending a row decides the row before it, and the accepting condition decides the last one.

An array of n' rows is then exactly a walk of length n' from the empty state to an accepting one, so $a(n) = w^T M^{n'} f$ and the count is C-finite.

4 The degree bound, derived

The reachable part of the digraph has 58 states, 44 after trimming; the forward identification of states with equal numbers of completions of every length, then the mirror identification on the edges coming in, leaves $S = 11$ blocks. The count satisfies the linear recurrence given by the characteristic polynomial of that 11×11 matrix; the bound is computed, not assumed.

5 The decision procedure

Let $c_1, \dots, c_D$ be the coefficients of a claimed recurrence $a(n) = \sum_{i=1}^D c_i a(n - i)$, let $q(t) = t^D - \sum_{i=1}^D c_i t^{D-i}$ be its characteristic polynomial, and put

$$u_m = \tilde{w}^T L^{m-1} q(L) \tilde{f} \quad (m \geq 1).$$

Expanding the powers of L and using the displayed formula for a , one gets $u_m = a(n) - \sum_{i=1}^D c_i a(n - i)$ with $n = m + D$. So u_m is exactly the residual of the claim at that index, and the recurrence holds at an index n if and only if the corresponding u_m vanishes.

Now $u_m = \tilde{w}^\top L^{m-1} y$ with $y = q(L)\tilde{f}$ a fixed vector, so (u_m) satisfies the characteristic polynomial of L , of degree 11: writing that polynomial as $t^{11} - \sum_{i=1}^{11} \lambda_i t^{11-i}$ gives $u_{m+11} = \sum_{i=1}^{11} \lambda_i u_{m+11-i}$. Hence 11 consecutive vanishing residuals force every later residual to vanish, by induction. The test is therefore finite; and because it locates the *last* nonvanishing residual, it pins the threshold exactly rather than merely bounding it.

Everything is carried out in exact integer arithmetic on the vector $L^{m-1}y$. The matrix M is never formed.

6 Results

Theorem 1. *For A280309, the recurrence $a(n) = 6a(n-1) - 6a(n-2) - 16a(n-3) + 12a(n-4) + 24a(n-5) + 8a(n-6)$ holds for every $n > 10$, and fails at $n = 10$.*

Proof. The residuals u_m were computed exactly for $m = 1, \dots, 51$. The last nonvanishing one is at $m = 4$, that is at $n = 10$. After it, more than $S = 11$ consecutive residuals vanish, so by Section 5 every later residual vanishes as well. $\square$

The entry claims the recurrence for $n > 10$. The theorem is at least as strong.

Theorem 2. *For A280309, the generating function is exactly $\sum_{n \geq 1} a(n)x^n = \frac{x^4 - 4x^5 + 3x^6 + 8x^7 - 6x^8 - 12x^9 - 4x^{10}}{1 - 6x + 6x^2 + 16x^3 - 12x^4 - 24x^5 - 8x^6}$.*

Proof. Let $N(x)$ and $D(x)$ be the numerator and denominator above, $D(0) = 1$. The residual test applied to the recurrence read off D — namely $a(n) = \sum_i c_i a(n-i)$ with c_i the negated coefficients of D — gives vanishing residuals for every $n > 10$, so the power series $F(x) = \sum_{n \geq 1} a(n)x^n$ satisfies $[x^k](F(x)D(x)) = 0$ for every $k > 11$. The remaining coefficients of $F(x)D(x)$, for $k \leq 19$, were computed from the walk count and agree with $N(x)$ term by term. Hence $F(x)D(x) = N(x)$ identically, which is the assertion. $\square$

7 Verification

Four checks were run, each reproducible from the code accompanying this note, and the first two are what pin the reading of the entry's English.

- **The published terms.** The walk count reproduces all 28 terms the entry publishes, exactly, in integer arithmetic, with the index taken from the entry's stated offset (1) rather than fitted. The first of them are 0, 0, 0, 1, 2, 9, 34, 124, ...
- **An independent enumeration.** For $n = 1, \dots, 11$ every one of the 3^{1n} arrays was written out and tested cell by cell against the condition as the entry states it — no transfer matrix, no row window, a separate piece of code. The counts agree with the walk count and with the entry.
- **The claim on the raw data.** Each claim was evaluated on the entry's published terms alone, with no model involved, wherever the proved range and the published data overlap. It holds there.
- **The annihilation test.** Carried to the derived bound $S = 11$ over the whole vector, in exact integer arithmetic, never sampled and never truncated.

8 References

1. OEIS Foundation Inc., *The On-Line Encyclopedia of Integer Sequences*, entry [A280309](#), revision 12, last modified Feb 19 2018.
2. R. P. Stanley, *Enumerative Combinatorics*, Vol. 1, 2nd ed., Cambridge University Press, 2012; §4.7, the transfer-matrix method.
3. M. Kauers and P. Paule, *The Concrete Tetrahedron*, Springer, 2011; C-finite sequences and their closure properties.
4. J. Berstel and C. Reutenauer, *Noncommutative Rational Series with Applications*, Cambridge University Press, 2011; linear representations of counting automata and their reduction.