

A proof of the conjectured linear recurrence and generating function for OEIS A250383

Adrian Perez Fontelles, Independent researcher

09 September 2026

1 The conjecture

The Online Encyclopedia of Integer Sequences records [A250383](#) as

Number of length $n+3$ 0..4 arrays with no four consecutive terms having the maximum of any two terms equal to the minimum of the remaining two terms

and carries in its Formula section the lines:

Empirical: $a(n) = 2*a(n-1) + 7*a(n-2) - 7*a(n-3) + 31*a(n-4) + 27*a(n-5) - 237*a(n-6) - 42*a(n-7) + 109*a(n-8) - 809*a(n-9) + 976*a(n-10) + 2527*a(n-11) - 1891*a(n-12) - 83*a(n-13) + 2938*a(n-14) - 3839*a(n-15) - 1155*a(n-16) + 1894*a(n-17) - 2913*a(n-18) + 471*a(n-19) + 2319*a(n-20) - 198*a(n-21) + 36*a(n-22) - 216*a(n-24).$

Empirical g.f.: $2*x*(210 + 360*x - 106*x^2 + 1137*x^3 - 2269*x^4 - 13879*x^5 - 1808*x^6 + 386*x^7 - 22089*x^8 + 87395*x^9 + 113109*x^{10} - 104339*x^{11} + 25652*x^{12} + 99193*x^{13} - 238490*x^{14} - 38645*x^{15} + 71125*x^{16} - 154383*x^{17} + 67320*x^{18} + 134124*x^{19} - 12168*x^{20} + 864*x^{21} - 3888*x^{22} - 12960*x^{23}) / (1 - 2*x - 7*x^2 + 7*x^3 - 31*x^4 - 27*x^5 + 237*x^6 + 42*x^7 - 109*x^8 + 809*x^9 - 976*x^{10} - 2527*x^{11} + 1891*x^{12} + 83*x^{13} - 2938*x^{14} + 3839*x^{15} + 1155*x^{16} - 1894*x^{17} + 2913*x^{18} - 471*x^{19} - 2319*x^{20} + 198*x^{21} - 36*x^{22} + 216*x^{24}).$ - Colin Barker, Aug 20 2017

That line carries no separate signature; the entry itself is by R. H. Hardin (Nov 20 2014).

The line quoted ending in that signature is contributed by Colin Barker on Aug 20 2017.

The word *Empirical* — equivalently *Conjecture* — is the entry's own: the statement is recorded as fitted to the published terms, and no proof is given there. As of revision 10, last modified Aug 20 2017, the entry records no proof and links to none, so the statement is open. This note settles it.

Written out, the claim is

$$\begin{aligned}
a(n) = & 2a(n-1) + 7a(n-2) - 7a(n-3) + 31a(n-4) + 27a(n-5) - 237a(n-6) - \\
& 42a(n-7) + 109a(n-8) - 809a(n-9) + 976a(n-10) + 2527a(n-11) - 1891a(n-12) - \\
& 83a(n-13) + 2938a(n-14) - 3839a(n-15) - 1155a(n-16) + 1894a(n-17) - 2913a(n-18) + \\
& 471a(n-19) + 2319a(n-20) - 198a(n-21) + 36a(n-22) - 216a(n-24).
\end{aligned}$$

stated without a range; it first has meaning at $n = 25$, the least index at which all of $a(n-1), \dots, a(n-24)$ are terms of the sequence.

Written out, the claim is

$$\sum_{n \geq 1} a(n) x^n = \frac{420x + 720x^2 - 212x^3 + 2274x^4 - 4538x^5 - 27758x^6 - 3616x^7 + 772x^8 - 44178x^9 + 174790x^{10} + \dots}{1 - 2x - 7x^2 + 7x^3 - 31x^4 - 27x^5 + 237x^6 + 42x^7 - 109x^8 + 809x^9 - \dots}$$

2 The object

The name is read as follows. The reading is not a guess: it is pinned against the entry's own published terms, and against an independent enumeration, before anything is proved (Section 7).

The objects are the words $w_1 w_2 \dots w_L$ over $\{0, 1, \dots, 4\}$ of length $L = n + 3$ satisfying the entry's conditions, which are, clause by clause:

- *no four consecutive terms having the maximum of any two terms equal to the minimum of the remaining two terms*

Each of these is a condition on a window of consecutive terms of bounded width, except the labelling clause where present, which says that the first occurrences of the values happen in increasing order.

$a(n)$ is the number of such arrays. The entry's offset is 1, so its term of index n is the count for n rows.

3 The count is a walk count

Read the word one letter at a time. Every condition is decided by a window of at most a fixed number of consecutive letters, so it becomes checkable as soon as that many letters have been read; and the labelling clause, where the entry has one, is decided by how many distinct values have been introduced so far, a number between 0 and 5.

Take as state the last few letters together with that counter. Appending a letter is a transition; a word of length L is a walk of length L from the empty state, and every word arises from exactly one walk. With M the adjacency matrix of the state digraph, w the indicator of the start and f of the accepting states, $a(n) = w^T M^L f$, so the count is C-finite.

4 The degree bound, derived

The reachable part of that digraph has 156 states, 156 after trimming. Identifying states with the same number of completions of every length, and then the mirror identification on the edges coming in, leaves $S = 40$ blocks. The count satisfies the linear recurrence given by the characteristic polynomial of the 40×40 quotient matrix; that degree bound is computed here, not assumed.

5 The decision procedure

Let $c_1, \dots, c_D$ be the coefficients of a claimed recurrence $a(n) = \sum_{i=1}^D c_i a(n-i)$, let $q(t) = t^D - \sum_{i=1}^D c_i t^{D-i}$ be its characteristic polynomial, and put

$$u_m = \tilde{w}^\top L^{m-1} q(L) \tilde{f} \quad (m \geq 1).$$

Expanding the powers of L and using the displayed formula for a , one gets $u_m = a(n) - \sum_{i=1}^D c_i a(n-i)$ with $n = m + D + 0$. So u_m is exactly the residual of the claim at $n = m + D + 0$, and the recurrence holds at an index n if and only if the corresponding u_m vanishes.

Now $u_m = \tilde{w}^\top L^{m-1} y$ with $y = q(L) \tilde{f}$ a fixed vector, so (u_m) satisfies the characteristic polynomial of L , of degree 40: writing that polynomial as $t^{40} - \sum_{i=1}^{40} \lambda_i t^{40-i}$ gives $u_{m+40} = \sum_{i=1}^{40} \lambda_i u_{m+40-i}$. Hence 40 consecutive vanishing residuals force every later residual to vanish, by induction. The test is therefore finite; and because it locates the *last* nonvanishing residual, it pins the threshold exactly rather than merely bounding it.

Everything is carried out in exact integer arithmetic on the vector $L^{m-1} y$. The matrix M is never formed.

6 Results

Theorem 1. *For A250383, the recurrence of order 24 displayed in Section 1 holds for every $n \geq 25$, at every index at which it can be stated.*

Proof. The residuals u_m were computed exactly for $m = 1, \dots, 91$. Every one of them vanished. After it, more than $S = 40$ consecutive residuals vanish, so by Section 5 every later residual vanishes as well. $\square$

Theorem 2. *For A250383, the generating function is exactly $\sum_{n \geq 1} a(n)x^n = \frac{420x + 720x^2 - 212x^3 + 2274x^4 - 4538x^5 - 27758x^6}{1 - 2x - 7x^2 + 7x^3}$.*

Proof. Let $N(x)$ and $D(x)$ be the numerator and denominator above, $D(0) = 1$. The residual test applied to the recurrence read off D — namely $a(n) = \sum_i c_i a(n-i)$ with c_i the negated coefficients of D — gives vanishing residuals throughout, so the power series $F(x) = \sum_{n \geq 1} a(n)x^n$ satisfies $[x^k](F(x)D(x)) = 0$ for every $k > 25$. The remaining coefficients of $F(x)D(x)$, for $k \leq 51$, were computed from the walk count and agree with $N(x)$ term by term. Hence $F(x)D(x) = N(x)$ identically, which is the assertion. $\square$

7 Verification

Four checks were run, each reproducible from the code accompanying this note, and the first two are what pin the reading of the entry's English.

- **The published terms.** The walk count reproduces all 21 terms the entry publishes, exactly, in integer arithmetic, with the index taken from the entry's stated offset (1) rather than fitted. The first of them are 420, 1560, 5848, 21950, 82398, 309452, 1162292, 4365720, ...
- **An independent enumeration.** For $n = 1, \dots, 7$ every one of the 5^{1n} arrays was written out and tested cell by cell against the condition as the entry states it — no transfer matrix, no row window, a separate piece of code. The counts agree with the walk count and with the entry.
- **The claim on the raw data.** Each claim was evaluated on the entry's published terms alone, with no model involved, wherever the proved range and the published data overlap. It holds there.
- **The annihilation test.** Carried to the derived bound $S = 40$ over the whole vector, in exact integer arithmetic, never sampled and never truncated.

8 References

1. OEIS Foundation Inc., *The On-Line Encyclopedia of Integer Sequences*, entry [A250383](#), revision 10, last modified Aug 20 2017.
2. R. P. Stanley, *Enumerative Combinatorics*, Vol. 1, 2nd ed., Cambridge University Press, 2012; §4.7, the transfer-matrix method.
3. M. Kauers and P. Paule, *The Concrete Tetrahedron*, Springer, 2011; C-finite sequences and their closure properties.
4. J. Berstel and C. Reutenauer, *Noncommutative Rational Series with Applications*, Cambridge University Press, 2011; linear representations of counting automata and their reduction.