

95-96	1	0	2	$x^6 - x^3 + 1$	3	24	4	11	5	29
Individual	6	$(-2, 2)$	7	216	8	$\frac{229}{99} = 2\frac{31}{99}$	9	40	10	$-(x-y)(y-z)(z-x)$

95-96	1	6	2	$3\sqrt{3}$ cm	3	91	4	$\frac{1}{5}$	5	$\frac{49}{100}$
Group	6	432	7	400	8	$\frac{365}{38}$	9	6	10	5

Individual Events

- I1**
- Find
- x
- if
- $4^{x-3} = 8^{x-2}$
- .

$$2^{2(x-3)} = 2^{3(x-2)}$$

$$2(x-3) = 3(x-2)$$

$$x = 0$$

- I2**
- If
- $f\left(\frac{1+x}{x}\right) = \frac{x^2+1}{x^2} + \frac{1}{x}$
- , find
- $f(x^3)$
- .

$$\text{Let } y = \frac{1+x}{x}, \text{ then } xy = 1+x$$

$$\Rightarrow x = \frac{1}{y-1}$$

$$f(y) = 1 + \frac{1}{x} + \frac{1}{x^2}$$

$$= 1 + y - 1 + (y-1)^2$$

$$= y + y^2 - 2y + 1$$

$$= y^2 - y + 1$$

$$f(y^3) = y^6 - y^3 + 1$$

$$\Rightarrow f(x^3) = x^6 - x^3 + 1$$

- I3**
- By considering
- $n! = n \times (n-1) \times (n-2) \times \dots \times 3 \times 2 \times 1$
- , find the number of trailing zeros of
- $100!$
- .

Reference: 1990 HG6, 1994 FG7.1, 2004 FG1.1, 2011 HG7, 2012 FI1.4, 2012 FG1.3

When each factor of 5 is multiplied by 2, a trailing zero will appear in the product.

The number of factors of 2 is clearly more than the number of factors of 5 in $100!$

It is sufficient to find the number of factors of 5.

5, 10, 15, ..., 100; altogether 20 numbers, have at least one factor of 5.

25, 50, 75, 100; altogether 4 numbers, have two factors of 5.

 \therefore Total number of factors of 5 is $20 + 4 = 24$ There are 24 trailing zeros of $100!$

- I4**
- What is the largest integral value
- n
- that satisfies the inequality
- $n^{200} < 5^{300}$
- ?

Reference: 1999 FG5.3, 2008 FI4.3, 2018FG2.4

$$n^2 < 5^3 = 125$$

$$n < \sqrt{125} < \sqrt{144} = 12$$

The largest integral $n = 11$

- I5** A set of 110 stamps of the denominations of \$0.1, \$3, \$5 worth \$100 in total.
Find the number of \$3 stamps in the set of stamps.
Suppose there are x \$0.1 stamps, y \$3 stamps, $(110 - x - y)$ \$5 stamps; where x and y are integers.

$$0.1x + 3y + 5(110 - x - y) = 100$$

$$x + 30y + 50(110 - x - y) = 1000$$

$$49x + 20y = 4500 \dots\dots (*)$$

$$49 = 20 \times 2 + 9 \dots\dots (1) \Rightarrow 9 = 49 - 20 \times 2 \dots\dots (1)'$$

$$20 = 9 \times 2 + 2 \dots\dots (2) \Rightarrow 2 = 20 - 9 \times 2 \dots\dots (2)'$$

$$9 = 2 \times 4 + 1 \dots\dots (3) \Rightarrow 1 = 9 - 2 \times 4 \dots\dots (3)'$$

$$\text{Sub. (1)'} \text{ into (2)'}: 2 = 20 - (49 - 20 \times 2) \times 2 \dots\dots (4)'$$

$$2 = 20 \times 5 - 49 \times 2 \dots\dots (5)'$$

$$\text{Sub. (1)'} \text{ and (5)'} \text{ into (3)'}: 1 = (49 - 20 \times 2) - (20 \times 5 - 49 \times 2) \times 4$$

$$1 = 49 \times 9 - 20 \times 22$$

$$\text{Multiply by 4500: } 49 \times 40500 + 20 \times (-99000) = 4500$$

$$\therefore x = 40500, y = -99000 \text{ is a particular solution to } (*)$$

The general solution is $x = 40500 - 20m$, $y = -99000 + 49m$, where m is an integer.

$$x > 0 \text{ and } y > 0 \text{ and } 110 - x - y > 0$$

$$40500 - 20m > 0 \text{ and } -99000 + 49m > 0 \text{ and } 110 - (40500 - 20m) - (-99000 + 49m) > 0$$

$$2025 > m > 2020 \frac{20}{49} \text{ and } 58610 > 29m$$

$$2025 > m > 2020 \frac{20}{49} \text{ and } 2021 \frac{1}{29} > m$$

$$\therefore m = 2021; y = -99000 + 49 \times 2021 = 29$$

Method 2 (*) can be written as $49x = 20(225 - y)$

$\therefore 49$ and 20 are relatively prime

$\therefore x$ is divisible by 20 and $225 - y$ is divisible by 49

x	$225 - y$	y	$110 - x - y$
20	49	176	no solution
40	98	127	no solution
60	147	78	no solution
80	196	29	1

$$\therefore y = 29$$

- I6** For any value of m , a straight line $y = mx + 2m + 2$ passes through a fixed point P . Find the coordinates of P . **Reference: 1990 HI5, 1991 HI6**

$$\text{Put } m = 0, y = 2$$

$$\text{Put } m = 1 \text{ and } y = 2 \Rightarrow x = -2$$

The coordinates of P is $(-2, 2)$.

- I7** How many 3-digit numbers can be made from the figures 4, 5, 6, 7, 8, 9 when repetitions are allowed?

$$\text{The number of 3-digit numbers} = 6^3 = 216$$

- I8** Express $2.\dot{3}\dot{1}$ as a fraction.

$$\text{Let } a = 2.\dot{3}\dot{1}$$

$$100a = 231.\dot{3}\dot{1}$$

$$100a - a = 229$$

$$a = \frac{229}{99} = 2\frac{31}{99}$$

- I9** If x and y are positive integers and $x - y = 5$, find the least value of $x^2 - y^2 + 5$.

$$x - y = 5 \Rightarrow x = 5 + y \dots\dots (1)$$

$$\begin{aligned} \text{Sub. (1) into } x^2 - y^2 + 5 &= (5 + y)^2 - y^2 + 5 \\ &= 30 + 10y \end{aligned}$$

$$y \geq 1 \Rightarrow 30 + 10y \geq 40$$

\therefore The least value of $x^2 - y^2 + 5$ is 40.

I10 Factorize $x^2(y - z) + y^2(z - x) + z^2(x - y)$.

$$\text{Let } f(x, y, z) = x^2(y - z) + y^2(z - x) + z^2(x - y)$$

$$f(x, y, z) = f(y, z, x) = f(z, x, y)$$

$f(x, y, z)$ is a cyclic expression of order 3.

$$f(x, x, z) = x^2(x - z) + x^2(z - x) + z^2(x - x) = 0$$

$\therefore (x - y)$ is a factor

By symmetry, $(y - z)$ and $(z - x)$ are factors

$$f(x, y, z) = k(x - y)(y - z)(z - x)$$

Compare the coefficients of x^2y : $-k = 1$

$$\Rightarrow k = -1$$

$$\therefore x^2(y - z) + y^2(z - x) + z^2(x - y) = -(x - y)(y - z)(z - x)$$

Group Events

- G1** In the figure, the quadratic curve $y = f(x)$ cuts the x -axis at the two points $(1, 0)$ and $(5, 0)$ and the y -axis at the point $(0, -10)$. Find the value of p .


$$f(x) = k(x-1)(x-5)$$

$$\text{It passes through } (0, -10) \Rightarrow f(0) = -10$$

$$-10 = k(0-1)(0-5) \Rightarrow k = -2$$

$$f(x) = -2(x-1)(x-5)$$

$$f(4) = p = -2(4-1)(4-5) = 6$$



- G2** In the figure, O is the centre of the base circle of a cone and the points A, B, C and O lie in the same plane. An ant walks from A to B on the surface of the cone. Find the length of the shortest path from A to B .

Let the vertex of the cone be V .

If we cut the curved surface of the cone along OA , a sector

$VACA'$ is formed with C as the mid-point of $\widehat{AA'}$.

Let $\angle AVB = \theta$ (in degree), then $\angle A'VB = \theta$

$\widehat{ACA'}$ = circumference of the base = 4π cm


$$2\pi \cdot 6 \cdot \frac{2\theta}{360} = 4\pi$$

$$\theta = 60^\circ$$

The shortest distance = the line segment AB

$$\text{By cosine rule, } AB^2 = [3^2 + 6^2 - 2(3)(6) \cos 60^\circ] \text{ cm}^2$$

$$AB = 3\sqrt{3} \text{ cm}$$



- G3** When a sum of \$7020, in the form of ten-dollar notes, is divided equally among x persons, \$650 remains. When this sum \$650 is changed to five-dollar coins and then divided equally among the x persons, \$195 remains. Find x .

Suppose each person gets a pieces of ten-dollar notes and b five-dollar coins.

$$7020 = 10ax + 650 \dots\dots (1), a, x \text{ are positive integers and } x > 65$$

$$650 = 5bx + 195 \dots\dots (2), b, x \text{ are positive integers and } x > 195 \div 5 = 39$$

$$\text{From (1): } 637 = ax \dots\dots (3)$$

$$\text{From (2): } 91 = bx \dots\dots (4)$$

$$\text{From (3): } 637 = 7 \times 91 = 7 \times 7 \times 13$$

\therefore The only positive factor of 91 is 91 which is greater than 65.

$$\therefore x = 91$$

- G4** In a shooting competition, according to statistics, A misses one in every 5 shoots, B misses one in every 4 shoots and C misses one in every 3 shoots. Find the probability of obtaining successful shoots by A, B but not C .

$$\text{Probability} = \frac{4}{5} \times \frac{3}{4} \times \frac{1}{3} = \frac{1}{5}$$

- G5** Given that $\frac{1}{n(n+1)} = \frac{1}{n} - \frac{1}{n+1}$, find the value of $\frac{1}{2 \times 3} + \frac{1}{3 \times 4} + \dots + \frac{1}{99 \times 100}$.

$$\frac{1}{2 \times 3} + \frac{1}{3 \times 4} + \dots + \frac{1}{99 \times 100} = \left(\frac{1}{2} - \frac{1}{3} \right) + \left(\frac{1}{3} - \frac{1}{4} \right) + \dots + \left(\frac{1}{99} - \frac{1}{100} \right) = \frac{1}{2} - \frac{1}{100} = \frac{49}{100}$$

- G6** If 3 is added to a 3-digit number A , the sum of the digits of the new number is $\frac{1}{3}$ of the value of the sum of digits of the original number A . Find the sum of all such possible numbers A .

$$\text{Let } A = 100a + 10b + c$$

$$\text{New number} = 100a + 10b + c + 3$$

$$\text{If } c \leq 6, a + b + c + 3 = \frac{1}{3}(a + b + c)$$

$$\Rightarrow 2(a + b + c) + 9 = 0, \text{ no solution}$$

$$\text{If new number} = 100a + 10(b + 1) + c + 3 - 10 = 100a + 10(b + 1) + c - 7, c \geq 7$$

$$a + b + 1 + c - 7 = \frac{1}{3}(a + b + c)$$

$$\Rightarrow 2(a + b + c) - 18 = 0$$

$$a + b + c = 9$$

a	b	c	A
1	0	8	108
1	1	7	117
2	0	7	207

$$\text{If } a \leq 8, b = 9 \text{ and } c \geq 7, \text{ new number} = 100(a + 1) + c - 7$$

$$a + 1 + c - 7 = \frac{1}{3}(a + b + c)$$

$$\Rightarrow 2(a + c) - 18 = 9$$

$$\Rightarrow 2(a + c) = 27, \text{ no solution}$$

$$\text{If } a = 9, b = 9 \text{ and } c \geq 7, \text{ new number} = 1000 + c - 7$$

$$1 + c - 7 = \frac{1}{3}(a + b + c)$$

$$\Rightarrow 3c - 18 = 18 + c$$

$$\Rightarrow c = 18, \text{ rejected}$$

$$\text{Sum of all possible } A = 108 + 117 + 207 = 432$$

- G7** In the figure, the side of each smaller square is 1 unit long. Find the sum of the area of all possible rectangles (squares included) that can be formed in the figure.

$$\text{Number of rectangles with sides } 1 \times 1 = 16$$

$$\text{Number of rectangles with sides } 1 \times 2 = 2 \times 3 \times 4 = 24$$

$$\text{Number of rectangles with sides } 1 \times 3 = 2 \times 2 \times 4 = 16$$

$$\text{Number of rectangles with sides } 1 \times 4 = 2 \times 4 = 8$$

$$\text{Number of rectangles with sides } 2 \times 2 = 3 \times 3 = 9$$

$$\text{Number of rectangles with sides } 2 \times 3 = 2 \times 2 \times 3 = 12$$


$$\text{Number of rectangles with sides } 2 \times 4 = 2 \times 3 = 6$$

$$\text{Number of rectangles with sides } 3 \times 3 = 2 \times 2 = 4$$

$$\text{Number of rectangles with sides } 3 \times 4 = 2 \times 2 = 4$$

$$\text{Number of rectangles with sides } 4 \times 4 = 1$$

$$\text{Sum of areas} = 16 + 24 \times 2 + 16 \times 3 + 8 \times 4 + 9 \times 4 + 12 \times 6 + 6 \times 8 + 4 \times 9 + 4 \times 12 + 16 = 400$$



- G8** If prime numbers a, b are the roots of the quadratic equation $x^2 - 21x + t = 0$, find the value of $\left(\frac{b}{a} + \frac{a}{b}\right)$.

Reference: 1996FG7.1, 2001 FG4.4, 2005 FG1.2, 2012 HI6

$$a + b = 21; ab = t$$

$\therefore a, b$ are prime numbers and 21 is odd

$$\therefore a = 2, b = 19$$

$$\begin{aligned}\left(\frac{b}{a} + \frac{a}{b}\right) &= \frac{19}{2} + \frac{2}{19} \\ &= \frac{365}{38}\end{aligned}$$

- G9** Find the value of x such that the length of the path APB in the figure is the smallest.

Reference: 1983 FG8.1, 1991 HG9, 1993 HI1

Let the straight line segment DPE be as shown in the figure.

Reflect A along DE to A' .

Then $\triangle ADP \cong \triangle A'DP$

$$APB = A'P + PB$$

It is the shortest when A', P, B are collinear.

In this case, $\angle A'PD = \angle BPE$ (vert. opp. \angle s)



$\triangle A'PD \sim \triangle BPE$ (equiangular)

$DP : A'D = PE : BE$ (ratio of sides, $\sim \triangle$'s)

$$\frac{x}{3} = \frac{14-x}{4}$$

$$4x = 42 - 3x$$

$$x = 6$$



- G10** Find the units digit of the sum $1^2 + 2^2 + 3^2 + 4^2 + \dots + 123456789^2$.

Reference 2012 HI1

Sum of units digits from 1^2 to 10^2

$$\equiv 0 + 1 + 4 + 9 + 6 + 5 + 6 + 9 + 4 + 1 \pmod{10}$$

$$\equiv 5 \pmod{10}$$

\therefore Required units digit

$$= \text{units digit of } 12345679 \times 5$$

$$\equiv 5 \pmod{10}$$