

Locus Example 2

Created by Mr. Francis Hung on 20110422

Last updated: August 29, 2021

Given the circle $C: x^2 + y^2 - 4x + 6y + 4 = 0$ and a point $D(4, -2)$ lies inside the circle.

A variable chord passes through D meets the circle at A and B . Find the locus of the mid-point M .

Let the slope of AB be m .

$$AB: \frac{y+2}{x-4} = m \dots\dots\dots(1)$$

$$y = mx - 4m - 2 \dots\dots\dots(2)$$

$$\text{Sub. into } C: x^2 + (mx - 4m - 2)^2 - 4x + 6(mx - 4m - 2) + 4 = 0$$

$$x^2 + m^2x^2 - 2m(4m + 2)x + (4m + 2)^2 - 4x + 6mx - 6(4m + 2) + 4 = 0$$

$$(1 + m^2)x^2 + (-8m^2 + 2m - 4)x + (16m^2 + 16m + 4 - 24m - 12 + 4) = 0$$

$$(1 + m^2)x^2 - 2(4m^2 - m + 2)x + (16m^2 - 8m - 4) = 0 \dots\dots\dots(3)$$

If $A(x_1, y_1)$, $B(x_2, y_2)$, then x_1, x_2 are the solution of (3)

$$x_1 + x_2 = \text{sum of roots in (3)} = \frac{2(4m^2 - m + 2)}{1 + m^2}$$

$$\text{If } M(x_0, y_0), x_0 = \frac{x_1 + x_2}{2} = \frac{4m^2 - m + 2}{1 + m^2} = \frac{4m^2 + 4 - m - 2}{1 + m^2} = 4 - \frac{m + 2}{1 + m^2}$$

$$x_0 = 4 - \frac{m + 2}{1 + m^2} \dots\dots\dots(4)$$

$$\text{Sub. (1) into (4): } x = 4 - \frac{\frac{y+2}{x-4} + 2}{1 + \left(\frac{y+2}{x-4}\right)^2}$$

$$x - 4 = -\frac{(y+2)(x-4) + 2(x-4)^2}{(x-4)^2 + (y+2)^2}$$

$$1 = -\frac{y+2+2(x-4)}{x^2 + y^2 - 8x + 4y + 20}$$

$$x^2 + y^2 - 8x + 4y + 20 = -2x - y + 6$$

$$x^2 + y^2 - 6x + 5y + 14 = 0$$

Method 2

Let $M = (x, y)$, $G = \text{centre} = (2, -3)$

$GM \perp AB$ (line joining centre to mid point chord \perp chord)

$$\therefore m_{GM} \cdot m_{MD} = -1$$

$$\frac{y+3}{x-2} \cdot \frac{y+2}{x-4} = -1$$

$$(x-2)(x-4) + (y+3)(y+2) = 0$$

$$x^2 + y^2 - 6x + 5y + 14 = 0$$

