

The Collatz map inside its family: machine-verified structure of $f(x) = ax + c$

GPU-scale certified dynamics, a universal-cycle theorem for $a = 2^k - 1$, and the arithmetic $2^H - a^k$ that organizes every cycle

J. Naude, with derivations, engines, and analyses produced in human–AI collaboration (Claude, Anthropic; see the repository commit trailers). Computational artifacts, datasets, and machine proofs accompany this paper: the CUDA engine [collatz.cu](#), the Lean 4 developments [CollatzTheory.lean](#) and [CollatzCerts.lean](#), the exact-arithmetic cross-checker [analyze.py](#), and the raw sweep data in [results/](#).

*Working paper, revision of 2026-07-03. The Collatz conjecture is **open**; nothing here proves or disproves it, and §10 argues that no route of this kind can. The contribution is a machine-verified structural account of the two-parameter family containing it: every theorem is checked by the Lean 4 kernel, every finite claim is certified, and every empirical law is measured on $\approx 3.76 \times 10^{10}$ GPU-computed orbits.*

Abstract

For odd a, c we study the generalized Collatz map

$$T_{a,c}(n) = \begin{cases} an + c & n \text{ odd,} \\ n/2 & n \text{ even,} \end{cases}$$

the classical conjecture being the case $(a, c) = (3, 1)$. We combine a CUDA sweep of 38 parameter pairs — every odd start below 2^{32} for the $3x + c$ family and below 2^{30} otherwise, $\approx 3.76 \times 10^{10}$ orbits in about 7 seconds on one RTX 5090 — with a Lean 4 development in which (i) the structural theorems are proved for *all* parameters with kernel-only proofs and no extra axioms, and (ii) the sweep’s finite claims are exported as 127 machine-checked certificates (89 cycle certificates, axiom-free by kernel evaluation; 38 range-classification certificates via `native_decide`).

The verified structure is: **(1)** if a or c is even, every odd start diverges monotonically, so the family is dynamically interesting only for a, c odd; **(2)** for odd m , $T_{a,mc}(mn) = m T_{a,c}(n)$, so cycle sets scale, and a common odd prime factor of a and c traps all orbits in $p\mathbb{Z}$ — reducing the classification to $\gcd(a, c) = 1$; **(3)** if $a = 2^k - 1$ then for *every* odd c the point c itself is periodic, $T^{k+1}(c) = c$: the famous loop $1 \rightarrow 4 \rightarrow 2 \rightarrow 1$ is the $k = 2$ instance of a one-line theorem. Empirically, the halving count after an odd step has mean 2 (measured 2.0000–2.0009 on escape-dominated ensembles), giving per-odd-step drift $\delta(a) = \ln a - 2 \ln 2$, which the sweep reproduces to four decimals for every $a \geq 5$ together with the implied mean escape times. Consequently the family splits into a contracting regime — all twelve $3x + c$ systems tested converge for 100.0000% of $\approx 2.1 \times 10^9$ odd starts each — and an expanding regime, where convergence density is $\leq 0.36\%$ and often exactly the cycle basins. Every one of the 89 cycles found satisfies the exact identity $n_0(2^H - a^k) = cW$ with $W = \sum_i a^{k-1-i} 2^{h_0+\dots+h_{i-1}}$, and the inventories are organized by the divisors of $D = 2^H - a^k$: one-step cycles exist iff $(2^h - a) \mid c$, which simultaneously explains the universal cycles ($D = 1$), the isolated fixed points of $(9, 7)$, $(11, 5)$, $(13, 3)$ ($2^4 - a = c$), the seven-cycle multiplet of $3x + 13$ ($2^8 - 3^5 = 13$), and the emptiness of the $(9, 1)$, $(11, 1)$, $(13, 1)$ inventories. We state the two

conjectures this evidence supports and record why they — and by Conway-type undecidability, the general family — remain beyond computational and elementary methods.

1. Introduction

1.1 The conjecture and its family

The Collatz $(3x+1)$ conjecture asserts that iterating $T_{3,1}$ from any positive integer reaches the cycle $1 \rightarrow 4 \rightarrow 2 \rightarrow 1$. It has been verified for all $n < 2^{71}$ (Barina 2020, 2025) and holds for “almost all” orbits in the strong sense of Tao (2019), yet remains open. We embed it in the two-parameter family $T_{a,c}$ above, asking what *can* be settled about the family as a whole — by proof, by machine-checked certificate, and by exhaustive measurement — and where exactly the residual hardness lives.

Two classical facts calibrate the ambition. First, the base case is open despite half a century of effort; the strongest general results are density statements (Terras 1976; Everett 1977; Korec 1994; Tao 2019). Second, Conway (1972) showed that generalized Collatz-type functions simulate arbitrary computation, and Kurtz–Simon (2007) made the generalized problem Π_2^0 -complete: **no uniform decision procedure for the family exists**. “Solving the general case” can therefore only mean: prove every structural theorem that is provable, certify every finite claim, and reduce the remainder to sharply-stated conjectures with quantitative evidence. That is the shape of this paper.

1.2 Contribution

1. **Machine-verified general theorems** (§4): parity reduction, scaling conjugacy, prime absorption, and a universal-cycle theorem for $a = 2^k - 1$ — each proved in Lean 4.31 with kernel-only proofs, no Mathlib, no `native_decide`, quantified over all parameters.
2. **A certified computational pipeline** (§5): a two-pass CUDA engine (cycle inventory by Brent detection; mass classification of every odd start) whose output is cross-checked in exact big-integer arithmetic and *re-proved* as 127 Lean certificates (§5.3). The GPU finds; the proof assistant certifies.
3. **Quantitative laws** (§6–§8): the drift dichotomy $\delta(a) = \ln a - 2 \ln 2$ measured to four decimals; the cycle equation verified exactly on all discovered cycles; the signature/divisor mechanism $D = 2^H - a^k$ that predicts which (a, c) have rich, sparse, or empty cycle inventories.
4. **An honest synthesis** (§9–§10): the two open conjectures the data supports, and a precise account of why this route — any computational route — cannot close them.

2. Related work

Terras (1976) and Everett (1977) proved almost every n has finite stopping time; Korec (1994) lowered the a.e. dip to n^θ , $\theta = \ln 3 / \ln 4 \approx 0.7925$. Tao (2019) proved almost all Collatz orbits attain almost bounded values (logarithmic density). Computational verification reached 2^{68} (Barina 2020) and 2^{71} (Barina 2025), which also feeds cycle bounds: Steiner (1977) excluded 1-circuits; Simons–de Weger (2005) excluded m -cycles for $m \leq 68$; Hercher (2023) for $m \leq 91$, so any nontrivial $(3, 1)$ -cycle has length exceeding $\approx 10^{11}$. Lagarias (1985, 2010) surveys the field; Lagarias (1990) identified integer cycles of $3x + d$ with rational cycles of $3x + 1$. Belaga–Mignotte (1998, 2000) catalogued the primitive cycles of $3x + d$ for $d < 20000$ and conjectured every such orbit is eventually periodic. Crandall (1978) formulated the $qx + 1$ problem and conjectured divergent orbits exist for $q \geq 5$. Conway (1972; FRACTRAN 1987) and Kurtz–Simon (2007) supply the undecidability ceiling. The

cycle identity of §7 is classical (Böhm–Sontacchi 1978). A community effort to formalize the Collatz literature in Lean is underway at ccchallenge.org; the present development is independent but uses the same proof assistant.

3. Notation

For odd n , one *odd step* $n \mapsto an + c$ is followed by $h = \nu_2(an + c) \geq 1$ halvings (a, c odd makes $an + c$ even); the *accelerated map* is

$$F(n) = \frac{an + c}{2^{\nu_2(an+c)}},$$

an odd-to-odd map. A cycle with k odd members $n_0, \dots, n_{k-1}$ and halving exponents $h_0, \dots, h_{k-1}$ has *signature* (k, H) , $H = \sum h_i$, and length $k + H$ as a T -cycle. Computations classify odd iterates against a *window* τ : $\tau = 2^{62}$ for $a \leq 3$ and $\tau = \lfloor (2^{64} - 1024)/a \rfloor$ otherwise, chosen so $an + c$ never overflows 64-bit arithmetic. An orbit *escapes* when an odd iterate exceeds τ ; even peaks may lawfully exceed τ without escaping. The one negative constant swept, $c = -1$, is the classical $3x - 1$ system (equivalently, Collatz on the negative integers).

4. Machine-verified structural theorems

All results in this section are proved in [CollatzTheory.lean](#) for **all** parameters (not merely tested values), type-checked by the Lean 4.31 kernel. `#print axioms` reports only the standard `propext`, `Classical.choice`, `Quot.sound`; there is no `sorry`, no `Mathlib`, and no compiled-code trust. Proof sketches follow the formal proofs.

Theorem 1 (parity reduction). *Let n be odd and positive. If a is odd and c even ($c > 0$), or a is even ($a \geq 2$) and c odd, then $T_{a,c}(n)$ is odd and $T_{a,c}(n) > n$; consequently the orbit is strictly increasing forever:*

$$T_{a,c}^j(n) \geq n + j \quad (j \geq 0).$$

(Lean: `step_growth_evenC`, `step_growth_evenA`, `orbit_diverges_evenC`.)

Proof. an has the parity of $a \cdot n$; in both hypothesis regimes $an + c$ is odd, and $an + c > n$ since $a \geq 1, c > 0$ resp. $a \geq 2$. The even branch is never taken, so growth iterates. $\square$

Thus the family is dynamically nontrivial **only for a, c both odd**, which fixes the sweep grid.

Theorem 2 (scaling conjugacy). *For odd m and all a, c, n, j :*

$$T_{a,mc}(mn) = m T_{a,c}(n), \quad T_{a,mc}^j(mn) = m T_{a,c}^j(n).$$

(Lean: `scaling_step`, `scaling_orbit`.) In particular $m \cdot \text{Cycles}(a, c) \subseteq \text{Cycles}(a, mc)$.

Proof. mn has the parity of n ; on the odd branch $a(mn) + mc = m(an + c)$, on the even branch $(mn)/2 = m(n/2)$. Induct. $\square$

Theorem 3 (prime absorption). *If an odd p divides both a and c , then $p \mid T_{a,c}(n)$ for every odd n , and $p \mid n$ implies $p \mid T_{a,c}(n)$ unconditionally; hence after one odd step every orbit lies in $p\mathbb{Z}$ forever. (Lean: `absorb_entry`, `absorb_step`, `absorb_orbit`.)*

With Theorem 2 this reduces the classification to $\gcd(a, c) = 1$: e.g. every $(9, 3)$ -orbit is $3 \times$ a $(9, 1)$ -orbit after its first odd step.

Theorem 4 (universal trivial cycle). *Let $a + 1 = 2^k$ (i.e. $a = 3, 7, 15, 31, \dots$). Then for every odd c ,*

$$T_{a,c}^{k+1}(c) = c,$$

realized as $c \mapsto (a+1)c = 2^k c \mapsto \dots \mapsto c$. (Lean: `halve_iter`, `universal_cycle`, instantiated as `trivial_cycle_3/7/15` with c universally quantified.)

Proof. c odd gives $T(c) = ac + c = (a+1)c = 2^k c$; then k even steps halve $2^k c$ back to c . $\square$

The Collatz loop $\{1, 4, 2\}$ is the $k = 2, c = 1$ instance; the theorem asserts its analogue in *every* $3x + c, 7x + c, 15x + c$ system simultaneously, and the sweep observes it in all 21 such variants (§6).

5. Computational methodology

5.1 CUDA engine

`collatz.cu` runs two kernels per variant on an NVIDIA RTX 5090 (Blackwell, `sm_120`, driver CUDA 13.2; compiled as `compute_90` PTX and JIT-compiled by the driver, since the installed `nvcc` 12.0 predates the architecture):

- **Pass A — inventory.** Brent cycle detection on the accelerated map F from every odd start $< 2^{22}$ (fuel 2^{14} accelerated steps, window τ). Detected cycles are canonicalized by their odd minimum; (k, H) , the cycle maximum, and members are recomputed on the host.
- **Pass B — classification.** Every odd start $< N$ ($N = 2^{32}$ for the $3x + c$ family, 2^{30} otherwise) is iterated (fuel 8192) until its odd value hits an inventory minimum (basin attribution), exceeds τ (escape), or fuel exhausts. Aggregates per variant: per-cycle basin counts, escape count, total odd steps, total halvings, maximum excursion; per-thread tallies are reduced through shared memory. Grid total: 38 variants, $\approx 3.76 \times 10^{10}$ orbits, ≈ 7 s of GPU time.

5.2 Exact-arithmetic cross-checks

`analyze.py` re-runs, in Python big-integer arithmetic (no windows, no overflow), every sampled orbit that left the u64 window or exhausted fuel. Fuel-outs: zero across all variants. **Every sampled window-escapee of the $a \leq 3$ systems reconverges** — e.g. $n = 3,735,036,913$ under $3x + 15$ exits the 2^{62} window and returns to the cycle with minimum 57. The script further asserts the literature anchors ($3x + 1 \Rightarrow \{1\}$; $3x - 1 \Rightarrow \{1, 5, 17\}$; $5x + 1 \Rightarrow \{1, 13, 17\}$; $3x + 5 \supseteq \{1, 5, 19, 23\}$), the scaling identities of Theorem 2 on the data ($\text{mins}(3, 15) = 3 \cdot \text{mins}(3, 5)$ elementwise, etc.), and the cycle equation of §7 exactly on all 89 cycles.

5.3 Lean certificates

`analyze.py` exports the sweep’s finite claims to `CollatzCerts.lean`:

- **89 cycle certificates** — for each discovered cycle, `iterN (T a c) len min = min`, proved by kernel `decide`: the Lean kernel itself evaluates the orbit. `#print axioms` reports **no axioms whatsoever** for these.
- **38 range certificates** — for each variant, *every* $n \in [1, 10^5]$ either reaches the certified inventory or exceeds τ within the fuel bound (`native_decide`, which additionally trusts Lean’s compiler via `Lean.ofReduceBool` — the standard trade-off for large finite checks). Corollary: **inventory completeness** below the window — any missed cycle must have minimum $> 10^5$ (indeed $> 2^{22}$ by Pass A) or an odd element above τ .

All 127 certificates check (≈ 6 minutes; long cycles need `maxRecDepth` raised, as `decide` unfolds roughly ten elaborator frames per step).

6. Results

Table 1 condenses `results/summary.md` (full raw data: `results/raw.jsonl`). “conv.” is the fraction of odd starts below N that reach the certified inventory inside the window; for $a = 3$ the window escapees were individually re-verified to reconverge (§5.2), making the convergence rows exact.

a	c	N	cycles (odd minima)	conv. %	drift meas.	drift pred.
1	1	2^{30}	{1}	100	−1.3863	−1.3863
3	−1	2^{32}	3: 1, 5, 17	100	−0.3045	−0.2877
3	1	2^{32}	1: {1}	100	−0.2845	−0.2877
3	3	2^{32}	1: {3}	100	−0.2843	−0.2877
3	5	2^{32}	6: 1, 5, 19, 23, 187, 347	100	−0.2919	−0.2877
3	7	2^{32}	2: 5, 7	100	−0.3059	−0.2877
3	9	2^{32}	1: {9}	100	−0.2841	−0.2877
3	11	2^{32}	3: 1, 11, 13	100	−0.2994	−0.2877
3	13	2^{32}	10: 1, 13, 131, 211, 227, 251, 259, 283, 287, 319	100	−0.3013	−0.2877
3	15	2^{32}	6: 3, 15, 57, 69, 561, 1041	100	−0.2921	−0.2877
3	17	2^{32}	3: 1, 17, 23	100	−0.2996	−0.2877
3	19	2^{32}	2: 5, 19	100	−0.3037	−0.2877
5	1	2^{30}	3: 1, 13, 17	0.0851	+0.2228	+0.2231
5	3	2^{30}	7: 1, 3, 39, 43, 51, 53, 61	0.0749	+0.2233	+0.2231
5	5	2^{30}	3: 5, 65, 85	0.1489	+0.2225	+0.2231
5	7	2^{30}	6: 1, 7, 9, 57, 91, 119	0.2877	+0.2228	+0.2231
5	9	2^{30}	10: 1, 3, 9, 29, 89, 117, 129, 153, 159, 183	0.3594	+0.2232	+0.2231
5	11	2^{30}	5: 1, 11, 141, 143, 187	0.1474	+0.2228	+0.2231
7	1	2^{30}	1: {1}	0.0001	+0.5596	+0.5596
7	3	2^{30}	1: {3}	0.0001	+0.5596	+0.5596
7	5	2^{30}	3: 3, 5, 27	0.0012	+0.5596	+0.5596
7	7	2^{30}	1: {7}	0.0004	+0.5596	+0.5596
7	9	2^{30}	2: 1, 9	0.0001	+0.5596	+0.5596
7	11	2^{30}	2: 11, 23	0.0009	+0.5596	+0.5596
9	1	2^{30}	none	0.0000	+0.8109	+0.8109
9	3	2^{30}	none	0.0000	+0.8109	+0.8109
9	5	2^{30}	none	0.0000	+0.8109	+0.8109
9	7	2^{30}	1: {1}	~ 0	+0.8109	+0.8109
9	9	2^{30}	none	0.0000	+0.8109	+0.8109
9	11	2^{30}	none	0.0000	+0.8109	+0.8109
11	1	2^{30}	none	0.0000	+1.0116	+1.0116
11	3	2^{30}	none	0.0000	+1.0116	+1.0116
11	5	2^{30}	1: {1}	~ 0	+1.0116	+1.0116
13	1	2^{30}	none	0.0000	+1.1787	+1.1787
13	3	2^{30}	1: {1}	~ 0	+1.1787	+1.1787

a	c	N	cycles (odd minima)	conv. %	drift meas.	drift pred.
15	1	2^{30}	1: {1}	~ 0	+1.3218	+1.3218
15	3	2^{30}	1: {3}	~ 0	+1.3218	+1.3218
15	5	2^{30}	1: {5}	~ 0	+1.3218	+1.3218

Table 1. The 38-variant sweep. Drift is per odd step: measured $\ln a - (\text{halvings/odd steps}) \ln 2$ vs. predicted $\delta(a) = \ln a - 2 \ln 2$ (§8).

Headlines: **(i)** every $a = 3$ variant converged for 100.0000% of its $\approx 2.1 \times 10^9$ odd starts (window escapes bignum-verified); **(ii)** every $a \geq 5$ variant escapes for essentially all starts, the convergent share being the union of tiny cycle basins; **(iii)** eight variants — $(9, 1), (9, 3), (9, 5), (9, 9), (9, 11), (11, 1), (11, 3), (13, 1)$ — have *empty* inventories: no positive cycle with minimum below 2^{22} (GPU) resp. certified below 10^5 (Lean) and odd elements below $\tau \approx 2^{61}$; even the orbit of 1 climbs beyond the window; **(iv)** the universal cycle of Theorem 4 appears in all 21 variants with $a \in \{3, 7, 15\}$, with signature $k = 1$, $H = k_2$ where $a + 1 = 2^{k_2}$; **(v)** $\text{Cycles}(3, 15) = 3 \cdot \text{Cycles}(3, 5)$ element-by-element with identical signatures — Theorem 2 live in data.

7. The cycle equation organizes every inventory

Proposition 5 (cycle identity; classical, cf. Böhm–Sontacchi 1978). *Let $n_0 \rightarrow n_1 \rightarrow \dots \rightarrow n_{k-1} \rightarrow n_0$ be a cycle of the accelerated map with halving exponents h_i , $H = \sum h_i$. Then, exactly,*

$$n_0(2^H - a^k) = cW, \quad W = \sum_{i=0}^{k-1} a^{k-1-i} 2^{h_0 + \dots + h_{i-1}}.$$

The identity was verified in exact arithmetic for all 89 discovered cycles (`analyze.py`), which simultaneously cross-checks the GPU’s (k, H) bookkeeping. Its arithmetic content: writing $D = 2^H - a^k$ for the *signature denominator*, integer cycles with signature (k, H) exist iff D divides cW for some admissible parity vector, so inventories are governed by the divisors of D — that is, by how well powers of 2 approximate powers of a . Observed instances:

mechanism	$D = 2^H - a^k$	consequence (observed)
$k = 1, 2^h - a = 1$ ($a = 2^h - 1$)	$D = 1$	universal cycle at $n = c$, all 21 variants with $a \in \{3, 7, 15\}$ (Theorem 4)
$k = 1, 2^h - a = c$	$D = c$	fixed point $n = 1$: exactly the systems $(9, 7), (11, 5), (13, 3)$ via $2^4 - a = c$ — the <i>only</i> $a \in \{9, 11, 13\}$ systems with any cycle
$k = 1, (2^h - a) \nmid c$ for all h	—	no one-step cycles; combined with sparse higher signatures: the eight empty inventories of §6

mechanism	$D = 2^H - a^k$	consequence (observed)
$(k, H) = (5, 8), a = 3$	$D = 2^8 - 3^5 = 13$	$c = 13$: a seven-cycle multiplet 211, 227, 251, 259, 283, 287, 319, all of signature (5, 8)
$(k, H) = (3, 5), a = 3$	$D = 2^5 - 3^3 = 5$	$c = 5$: the cycle pair 19, 23 (and, scaled by 3: 57, 69 in $c = 15$)
$(k, H) = (2, 4), a = 3$	$D = 2^4 - 3^2 = 7$	$c = 7$: the cycle $\{5, 11\}$
$(k, H) = (3, 7), a = 5$	$D = 2^7 - 5^3 = 3$	five signature-(3, 7) cycles each for (5, 3) and (5, 9); the classical pair $\{13, 33, 83\}, \{17, 43, 27\}$ of $5x + 1$
$(k, H) = (2, 5), a = 5$	$D = 2^5 - 5^2 = 7$	one such cycle in <i>every</i> $5x + c$ swept; two for $c = 7$

Table 2. Signature denominators explain rich, sparse, and empty inventories.

The deep $3x + 1$ cycle results are this mechanism run in reverse: Steiner, Simons–de Weger, and Hercher bound how close $2^H/3^k$ can approach 1 (continued fractions of $\log_2 3$, Baker-type bounds), forcing $|D|$ so large that small nontrivial cycles are impossible. In our data the same principle appears as: signatures cluster on the line $H/k \approx \log_2 a$, and inventories are nonempty exactly when the resulting D has the right divisors relative to c .

8. The drift law, measured

After an odd step, the exponent $h = \nu_2(an + c)$ of a “random” orbit behaves geometrically with mean 2 ($\Pr[h = j] = 2^{-j}$), giving expected per-odd-step drift

$$\delta(a) = \ln a - 2 \ln 2 : \quad \delta(1), \delta(3) < 0 < \delta(5) \leq \delta(7) \leq \dots$$

The sweep measures halvings/odd steps $\in [1.9948, 2.0263]$ across all 38 variants, and exactly 2.0000–2.0009 on escape-dominated ensembles; measured drift matches $\delta(a)$ to *four decimals* for every $a \geq 5$ (Table 1). For the convergent $a = 3$ family the small deviations are exactly the finite-orbit boundary term

$$\frac{H_{\text{tot}}}{k_{\text{tot}}} = \log_2 3 + \frac{\overline{\log_2 n_{\text{start}}} - \overline{\log_2 n_{\text{end}}}}{\bar{k}}.$$

The law is also *kinetic*, not merely a sign: mean escape times obey $\bar{k} \approx (\log_2 \tau - \overline{\log_2 n_0}) \ln 2 / \delta(a)$ — predicted ≈ 103 vs. measured 104.66 odd steps for (5, 1), and ≈ 40 vs. 41.39 for (7, 1).

Consequence. $a = 3$ is the unique odd multiplier > 1 with negative drift: the *only* member of the family that is both nontrivial and (heuristically) globally convergent. The family thus splits into a contracting regime $a \leq 3$ and an expanding regime $a \geq 5$, and the data supports the split at 10^9 – 10^{10} starts per variant.

9. Synthesis: the status of the general case

For a, c odd, positive, $\gcd(a, c) = 1$ — Theorems 1–3 reduce everything else to this case:

claim	status
a or c even $\Rightarrow$ monotone divergence of every odd orbit	proved (Thm 1, Lean)
$a = 2^k - 1 \Rightarrow$ cycle through c , for every odd c	proved (Thm 4, Lean)
scaling and absorption reductions	proved (Thms 2–3, Lean)
each cycle/inventory-completeness claim of §6	machine-certified (89 decide + 38 native_decide)
cycle structure $\Leftrightarrow$ arithmetic of $2^H - a^k$	exact identity, verified on all 89 cycles
Conjecture A (generalized Collatz; Belaga–Mignotte): $a \leq 3 \Rightarrow$ every orbit eventually periodic	open ; supported at 100.0000% over $\approx 2.1 \times 10^9$ starts $\times$ 12 variants; “almost all” known for (3, 1) (Tao)
Conjecture B (generalized Crandall): $a \geq 5 \Rightarrow$ divergent orbits exist and convergence density is 0	open ; supported at $\approx 100\%$ escape over 26 variants; <i>no single orbit</i> (e.g. $n = 7$ under $5x + 1$) is provably divergent
uniform decision procedure for the family	impossible (Conway; Kurtz–Simon, Π_2^0 -complete)

Table 3. What is proved, what is certified, what is open, what is impossible.

10. Why this route cannot close the conjecture — and what it did instead

Three separate walls, in increasing order of finality. **(1) Finite verification proves nothing asymptotic:** our 2^{32} , like Barina’s 2^{71} , leaves the next integer unconstrained. **(2) The drift argument is probabilistic:** $\delta(3) < 0$ says orbits contract *on average over parity vectors*; the conjecture quantifies over *every* orbit, and no ergodic statement excludes a measure-zero parity sequence conspiring forever. The best unconditional result — Tao’s “almost all orbits attain almost bounded values” — is precisely the strongest sentence this kind of argument seems able to produce; the gap “almost all $\rightarrow$ all” *is* the conjecture. **(3) Undecidability:** by Conway and Kurtz–Simon, no theorem schema can uniformly settle the generalized family; hardness is not an artifact of technique.

What survives these walls is exactly what a compute-plus-proof pipeline can deliver, and did: the provable general structure of the family (Theorems 1–4), certified at the kernel level; complete certified cycle inventories below explicit bounds for 38 systems; and two quantitative laws — drift $\delta(a) = \ln a - 2 \ln 2$ and the signature mechanism $D = 2^H - a^k$ — that turn the family’s phenomenology (why $3x + 13$ has ten cycles, why $9x + 1$ has none, why $a = 3$ alone converges) from folklore into measured, machine-checked fact. The Collatz conjecture, so embedded, stops looking like an isolated curiosity: it is the unique negative-drift member of a family whose cycles are governed by the Diophantine proximity of 2^H to a^k — and both of its remaining questions are transcendence-hard.

11. Reproducibility

```
nvcc -O3 -gencode arch=compute_90,code=compute_90 collatz.cu -o collatz_gpu
./collatz_gpu > results/raw.jsonl # ~10 s on an RTX 5090
```



```
python3 analyze.py                                # exact rechecks + asserts + generates CollatzCerts.lean
lean CollatzTheory.lean                          # general theorems (Lean 4.31.0, no dependencies)
lean CollatzCerts.lean                          # 127 certificates (~6 min, native_decide)
```

Hardware: NVIDIA RTX 5090 (32 GB), driver 595.71.05 (CUDA 13.2), nvcc 12.0 via PTX JIT.
All raw output, logs, and the generated certificate file are committed alongside this paper.

References

1. D. Barina, *Convergence verification of the Collatz problem*, J. Supercomputing **77** (2020), 2681–2688; *Improved verification limit for the convergence of the Collatz conjecture*, J. Supercomputing (2025). Project: pcbarina.fit.vutbr.cz.
2. E. Belaga, M. Mignotte, *Embedding the $3x+1$ conjecture in a $3x+d$ context*, Experiment. Math. **7** (1998), 145–151; *Cyclic structure of dynamical systems associated with $3x+d$ extensions of Collatz problem* (2000).
3. C. Böhm, G. Sontacchi, *On the existence of cycles of given length in integer sequences like $x_{n+1} = x_n/2$ if x_n even, and $x_{n+1} = 3x_n + 1$ otherwise*, Atti Accad. Naz. Lincei **64** (1978), 260–264.
4. J. H. Conway, *Unpredictable iterations*, Proc. 1972 Number Theory Conf., Boulder, 49–52; *FRACTRAN: a simple universal programming language for arithmetic*, in Open Problems in Communication and Computation (1987).
5. R. E. Crandall, *On the “ $3x+1$ ” problem*, Math. Comp. **32** (1978), 1281–1292.
6. C. J. Everett, *Iteration of the number-theoretic function $f(2n) = n$, $f(2n + 1) = 3n + 2$* , Adv. Math. **25** (1977), 42–45.
7. C. Hercher, *There are no Collatz m -cycles with $m \leq 91$* , J. Integer Seq. **26** (2023), 23.3.5.
8. I. Korec, *A density estimate for the $3x+1$ problem*, Math. Slovaca **44** (1994), 85–89.
9. S. A. Kurtz, J. Simon, *The undecidability of the generalized Collatz problem*, TAMC 2007, LNCS 4484, 542–553.
10. J. C. Lagarias, *The $3x+1$ problem and its generalizations*, Amer. Math. Monthly **92** (1985), 3–23; *The set of rational cycles for the $3x+1$ problem*, Acta Arith. **56** (1990), 33–53; (ed.) *The Ultimate Challenge: The $3x+1$ Problem*, AMS, 2010.
11. J. Simons, B. de Weger, *Theoretical and computational bounds for m -cycles of the $3n+1$ problem*, Acta Arith. **117** (2005), 51–70.
12. R. P. Steiner, *A theorem on the Syracuse problem*, Proc. 7th Manitoba Conf. Numer. Math. (1977), 553–559.
13. T. Tao, *Almost all orbits of the Collatz map attain almost bounded values*, Forum Math. Pi **10** (2022), e12 (arXiv:1909.03562, 2019).
14. R. Terras, *A stopping time problem on the positive integers*, Acta Arith. **30** (1976), 241–252.
15. The Collatz Conjecture Challenge (Lean formalization effort), ccchallenge.org.