

Notes-From Face Ring to Partition Complex

Mingzhi Zhang

0 Background

This note is constantly evolving. The ultimate goal is to develop an understanding of the field **combinatorial commutative algebra**.

This note is very personal. The use of diagrams, analogies, stories, and other means is only a shadow of how the author thinks or likes to think/understand.

1 Sets and Multisets

1.1 Sets and subsets

Let

$$E = \{x_1, x_2, \dots, x_n\}$$

We will abbreviate or identify it with

$$[n] := \{1, 2, \dots, n\}.$$

Definition 1.1.1. For a subset $I \subseteq E$, the **characteristic function** of I is defined as:

$$\delta_I : E \rightarrow \{0, 1\}$$

where

$$\delta_I(i) = \begin{cases} 1, & \text{if } i \in I, \\ 0, & \text{if } i \notin I. \end{cases}$$

Identify I with the monomial $\prod_{i \in I} x_i$ or its characteristic vector $v_I = (\delta_I(1), \delta_I(2), \dots, \delta_I(n))$.

Definition 1.1.2. A **k-subset** of E is a subset $I \subseteq E$ whose cardinality is k .

1.2 Multisets

Fix $E = [n]$.

Definition 1.2.1. A **multiset** M is a pair (E, δ_E) , where

$$\delta_E(i) = \begin{cases} k_i & \text{if } i \in E, \\ 0 & \text{if } i \notin E, \end{cases}$$

where $k_i \in \mathbb{Z}_{>0}$ is the **multiplicity** of i .

We call $S = \{i \in E \mid \delta_E(i) \neq 0\}$ the **support** of the multiset, written as $\text{supp}(M) = S$.

Example 1.2.2. Let $E = \{1, 2, 3\}$, and

$$\delta_E(1) = 4, \quad \delta_E(2) = 0, \quad \delta_E(3) = 1.$$

The multiset is represented as: $M = \{1, 1, 1, 1, 3\}$, and $\text{supp}(M) = (1, 3)$.

Remark 1.2.3. A submultiset is subset of a multiset, and a **k-submultiset** is a submultiset whose cardinality is k .

Given $I \subseteq E$, we want to characterize the k -submultisets with support I .

Take $I = \{1, 3\} \subseteq [4]$ as an example. What are the 4-submultisets with support I ? Let's list them all: $\{1, 1, 1, 3\}$, $\{1, 1, 3, 3\}$, $\{1, 3, 3, 3\}$. We have 3 4-submultisets with support $\{1, 3\}$.

Lemma 1.2.4. Let $I \subseteq E = [n]$ and $m = |I|$. The set of k -submultisets with support I are in bijection with the set $\{(z_1, z_2, \dots, z_m) \in \mathbb{Z}_{\geq 1}^m \mid \sum_{i=1}^m z_i = k\}$.


Corollary 1.2.5. Let $I \subseteq E = [n]$ and $m = |I|$. The number of k -submultisets with support I is $\binom{k-1}{m-1}$.

1.3 Graded structures and reciprocity

The set of subsets of E forms a finite **graded Boolean lattice** B_n with the rank function:

$$\text{rk} : 2^E \rightarrow \mathbb{Z}_{\geq 0}, \quad I \mapsto |I| \quad (\text{cardinality}).$$

Example 1.3.1. Let $E = [3]$ and we have the Hasse diagram of B_3 :



The generating function of the Boolean lattice B_n is

$$F(x_1, x_2, \dots, x_n) = 1 + \sum_{i=1}^n x_i + \sum_{i < j} x_i x_j + \dots + x_1 x_2 \dots x_n = \prod_{i=1}^n (1 + x_i)$$

Let $x_1 = x_2 = \dots = x_n = x$. The generating function becomes:

$$F(x, x, \dots, x) = (1 + x)^n := \binom{n}{0} + \binom{n}{1}x + \binom{n}{2}x^2 + \dots + \binom{n}{n}x^n.$$

and we get the definition of **binomial coefficients**, the most familiar counting function.

Definition 1.3.2. $\binom{n}{k}$ is the number of k -subsets in $[n]$.

Remark 1.3.3. The vector $(\binom{n}{0}, \binom{n}{1}, \binom{n}{2}, \dots, \binom{n}{n})$ is the rank vector (or face vector) of the Boolean lattice (or simplicial complex) B_n .

All submultisets of a multiset form a graded lattice \mathcal{L}^∞ . The generating function of the Boolean lattice \mathcal{L}^∞ is

$$F(x_1, x_2, \dots, x_n) = 1 + \sum_{i=1}^{\infty} x_i + \sum_{i < j} x_i x_j + \dots = \prod_{i=1}^n \frac{1}{1 - x_i}$$

Let $x_1 = x_2 = \dots = x_n = x$. The generating function becomes:

$$F(x, x, \dots, x) = \frac{1}{(1 - x)^n} := \binom{n}{0} + \binom{n}{1}x + \binom{n}{2}x^2 + \dots + \binom{n}{n}x^n + \dots$$

In what follows we assume that each element in S has multiplicity infinity. Denote this multiset as S^∞ .

1.4 Down-closed property

For a family \mathcal{F} of subsets of $[n]$, we define the **down-closed property** as follows:

For any $\sigma \subseteq \tau \in \mathcal{F}$, then $\sigma \in \mathcal{F}$.

1.5 Euler characteristic

2 Stanley-Reisner Ring

2.1 Simplicial complex

2.2 Hilbert function

2.3 Cohen-Macaulay property

2.4 Upper Bound Theorem

2.5 Lefschetz property

2.6 G-theorem

References

- [Adi18] Adiprasito, Karim. "Combinatorial Lefschetz theorems beyond positivity." arXiv preprint arXiv:1812.10454 (2018).
- [AHK18] Adiprasito, Karim, June Huh, and Eric Katz. "Hodge theory for combinatorial geometries." *Annals of Mathematics* 188.2 (2018): 381-452.
- [AY21] Adiprasito, Karim, and Geva Yashfe. "The partition complex: an invitation to combinatorial." *Surveys in combinatorics 2021* 470 (2021): 1.
- [Sta75] Stanley RP. The upper bound conjecture and CohenMacaulay rings. *Studies in Applied Mathematics*. 1975 Jun;54(2):135-42.