

2D Exact Coherent State equations

November 6, 2013

1 The full equations

$$T_{ij} = \frac{1}{\text{Wi}}(C_{ij} - \mathbf{I}_{ij}) \quad (1)$$

$$u = -\partial_y \psi \quad (2)$$

$$v = \partial_x \psi \quad (3)$$

1.1 ψ

$$\begin{aligned} 0 = & \text{Re } \nu \partial_x \nabla^2 \psi - \text{Re } u \partial_x \nabla^2 \psi - \text{Re } v \partial_y \nabla^2 \psi \\ & + \beta \nabla^4 \psi - \frac{(1-\beta)}{\text{Wi}} (\partial_{xx} C_{xy} + \partial_{xy} (C_{yy} - C_{xx}) - \partial_{yy} C_{xy}) \end{aligned} \quad (4)$$

1.2 C_{xx}

$$\begin{aligned} 0 = & \text{Re } \nu \partial_x C_{xx} - (u \partial_x + v \partial_y) C_{xx} \\ & + 2C_{xx} \partial_x u + 2C_{xy} \partial_x v - T_{xx} \end{aligned} \quad (5)$$

1.3 C_{yy}

$$\begin{aligned} 0 = & \text{Re } \nu \partial_x C_{yy} - (u \partial_x + v \partial_y) C_{yy} \\ & + 2C_{xy} \partial_y u + 2C_{yy} \partial_y v - T_{yy} \end{aligned} \quad (6)$$

1.4 C_{xy}

$$\begin{aligned} 0 = & \text{Re } \nu \partial_x C_{xy} - (u \partial_x + v \partial_y) C_{xy} \\ & + C_{xx} \partial_y u + C_{yy} \partial_x v - T_{xy} \end{aligned} \quad (7)$$

1.5 ψ_0

$$\begin{aligned} 0 = & -(\overline{\partial_y \psi})(\overline{\partial_x \psi}) - \overline{(\partial_x \psi)}(\overline{\partial_y \psi}) \\ & + \beta \overline{\partial_{yy} \psi} + \frac{(1-\beta)}{\text{Wi}} \overline{\partial_y C_{xy}} \end{aligned} \quad (8)$$

2 The 1st order terms in a Taylor expansion

These are the equations as they will need to be entered into the Jacobian matrix.

$$\psi = \Psi + \delta\psi C_{ij} = C_{ij} + \delta c_{ij} \quad (9)$$

2.1 ψ

$$\begin{aligned} 0 = & \nu \text{Re } \partial_x \nabla^2 \delta\psi + \text{Re } (\nabla^2 \Psi) \partial_y \delta\psi - \text{Re } U \partial_x \nabla^2 \delta\psi \\ & - \text{Re } (\partial_y \nabla^2 \Psi) \partial_x \delta\psi - \Re V \partial_y \nabla^2 \delta\psi + \beta \nabla^4 \delta\psi \\ & - \frac{(1-\beta)}{\text{Wi}} (\partial_{xx} \delta c_{xy} + \partial_{yy} (\delta c_{yy} - \delta c_{xx}) - \partial_{yy} \delta c_{xy}) \end{aligned} \quad (10)$$

2.2 C_{xx}

$$\begin{aligned} 0 = & \nu \partial_x \delta c_{xx} - (\mathbf{V} \cdot \nabla) \delta c_{xx} \\ & + (\partial_x C_{xx}) \partial_y \delta\psi - (\partial_y C_{xx}) \partial_x \delta\psi \\ & + 2(\partial_x U) \delta c_{xx} + 2(\partial_x V) \delta c_{xy} \\ & - 2C_{xx} \partial_{xy} \delta\psi + 2C_{xy} \partial_{xx} \delta\psi - \frac{1}{\text{Wi}} \delta c_{xx} \end{aligned} \quad (11)$$

2.3 C_{yy}

$$\begin{aligned} 0 = & \nu \partial_x \delta c_{yy} - (\mathbf{V} \cdot \nabla) \delta c_{yy} \\ & + (\partial_x C_{yy}) \partial_y \delta\psi - (\partial_y C_{yy}) \partial_x \delta\psi \\ & + 2(\partial_y U) \delta c_{xy} + 2(\partial_y V) \delta c_{yy} - (\partial_x C_{yy}) \partial_y \delta\psi \\ & - C_{xy} \partial_{yy} \delta\psi + c_{yy} \partial_{xy} \delta\psi - \frac{\delta c_{yy}}{\text{Wi}} \end{aligned} \quad (12)$$

2.4 C_{xy}

$$\begin{aligned} 0 = & \nu \partial_x \delta c_{xy} - (\mathbf{V} \cdot \nabla) \delta c_{xy} \\ & + (\partial_x C_{xy}) \partial_y \delta\psi - (\partial_y C_{xy}) \partial_x \delta\psi \\ & + (\partial_x C_{xy}) \partial_y \delta\psi - (\partial_y C_{xy}) \partial_x \delta\psi \\ & - C_{xx} \partial_{yy} \delta\psi + C_{yy} \partial_{xx} \delta\psi - \frac{\delta c_{xy}}{\text{Wi}} \end{aligned} \quad (13)$$

2.5 ψ_0

$$\begin{aligned} 0 = & -\overline{(\partial_{xy} \Psi) \delta\psi} - \overline{(\partial_x \Psi) \delta\psi} \\ & + \beta \overline{\partial_{yy} \delta\psi} + \frac{1-\beta}{\text{Wi}} \overline{\delta c_{xy}} \end{aligned} \quad (14)$$