


Homework #2

Yunting Chiu

2021-02-02

- (1.2) The members of a health spa pay annual membership dues of \$\$\$300 plus a charge of \$\$\$2 for each visit to the spa. Let Y denote the dollar cost for the year for a member and X the number of visits by the member during the year. Express the relation between X and Y mathematically. Is it a functional relation or a statistical relation (that is, is the relation deterministic or stochastic)?
 - deterministic: the output of the model is entirely determined by the values of the parameters and the initial conditions; stochastic: random, unpredictable.
 - The association between X and Y is: $Y = 300 + 2X$ (dollars).
This is a functional relationship because a effect in the value of the X will cause the corresponding change in the value of the Y . X and Y will not have a uncertain effect.
- (1.6) Suppose the regression parameters are $\beta_0 = 200$ and $\beta_1 = 5.0$.

(a) Plot the regression equation.



(b) Predict the response for $X = 10$, 20 , and 40 .

(b)

$$Y_i = \beta_0 + \beta_1 X_i + \epsilon_i$$

$$Y = 200 + 5X_i + \epsilon_i$$

$$\epsilon_i \sim N(0, 1)$$

$$E\{Y_i\} = 200 + 5X_i + E\{\epsilon_i\} \xrightarrow{0}$$

$$E\{Y_i\} = 200 + 5 \times 10 = 250 \$ \text{ (if } X=10)$$

$$E\{Y_i\} = 200 + 5 \times 20 = 300 \$ \text{ (if } X=20)$$

$$E\{Y_i\} = 200 + 5 \times 40 = 400 \$ \text{ (if } X=40)$$

(c) Explain the meaning of parameters β_0 and β_1 .

- β_0 = Y intercept of regression line.
 - β_1 : one unit change in X, generates a β_1 unit change in Y.
3. (1.10) An analyst in a large corporation studied the relation between current annual salary (Y) and age (X) for the 46 computer programmers presently employed in the company. The analyst concluded that the relation is curvilinear, reaching a maximum at 47 years. Does this imply that the salary for a programmer increases until age 47 and then decreases? Explain.
- **curvilinear** only explain that X and Y are not linear relation. It is not true because it reaches its maximum at a point and then increasing at a decreasing rate meaning that wage first increases to a max at year 47 and then the increasing rate slows down. In reality, decreasing salary as people age in any company is not make sense, we also can find a bunch of examples in real world.
4. The time it takes to transmit a file always depends on the file size. Suppose you transmitted 30 files, with the average size of 126 Kbytes and the standard deviation of 35 Kbytes. The average transmittance time was 0.04 seconds with the standard deviation of 0.01 seconds. The correlation coefficient between the time and the size was 0.86. Based on this data, fit a linear regression model and predict the time it will take to transmit a 400 Kbyte file.
- Based on the model above, it will take 0.1085 sec to transmit a 400 Kbyte file.

30 samples $\left\{ \begin{array}{l} \text{time} = y \quad \text{average} : 0.04 \text{ sec} \quad \text{sd} : 0.01 \text{ sec} \\ \text{filesize} = x \quad \text{average} = 126 \text{ K} \quad \text{sd} = 35 \text{ K} \end{array} \right.$
 correlation coefficient (r) = 0.86

Q: find predict /m take to transmit a 400K byte file.

$$r = \frac{S_{xy}}{S_x S_y} \Rightarrow \beta_1 = \frac{S_{xy}}{S_x^2} = r \frac{S_y}{S_x}$$

$$r = 0.86 \Rightarrow \beta_1 = 0.86 \frac{0.01}{35} = 0.00025$$

$$\begin{aligned} \beta_0 &= \bar{y} - \beta_1 \bar{x} \Rightarrow 0.04 - 0.00025 \times 126 \\ &= 0.04 - 0.0315 = 0.0085 \end{aligned}$$

Model

$$y_i = \beta_0 + \beta_1 x_i + \epsilon_i$$

$$\begin{aligned} E\{y_i\} &= 0.0085 + E\{0.00025 \times 400\} \\ &= 0.0085 + E\{0.1\} \\ &= 0.1085 (\text{sec}) \end{aligned}$$

5. At a gas station, 180 drivers were asked to record the mileage of their cars and the number of miles per gallon. The results are summarized in the table.

(a) Compute the least squares regression line which describes how the number of miles per gallon depends on the mileage.

- X is Mileage, y is Miles per gallon. The y-intercept that I computed is 24.77155. That is, if mileage add one unit, the Miles per gallon will have 24.77155 increment.
- reference: https://www.google.com/search?q=least+squares+regression+line+mean+standard+deviation&rlz=1C5CHFA_enUS920US920&oq=east+squares+regression+mean+de&aqs=chrome.1.69i57j0i22i30l4.8645j1j9&sourceid=chrome&ie=UTF-8#kpvalbx=_lJMZYKK_EIqw5NoPqMOIuAg21

```
# slope(b1) = r*sd(y)/sd(x)
slope <- ((-0.17*3.4)/14634)
slope
```

```
## [1] -3.949706e-05
```

```
# y intercept(b0) = sample mean of y - slope* sample mean of x
yIntercept <- 23.8-(slope*24598)
yIntercept
```

```
## [1] 24.77155
```

(b) What do the obtained slope and intercept mean in this situation?

- The slope implies a line's steepness and the intercept shows the position where an y-axis intersects. That is, y intercept is on 24.77155 of y-axis.

(c) You purchase a used car with 35,000 miles on it. Predict the number of miles per gallon.

- 23.38915 Miles per gallon

```
# y = b0+b1*X
```

```
predict35000 <- yIntercept + slope * 35000
predict35000
```

```
## [1] 23.38915
```

6. (Stat-615 only) Show that the sample intercept b_0 is a linear and unbiased estimator of the population intercept β_0 .

7. (Computer project - 1.19, 1.24). Grade point average. The director of admissions of a small college selected 120 students at random from the new freshman class in a study to determine whether a students grade point average (GPA) at the end of the freshman year (Y) can be predicted from the ACT test score (X). The results of the study follow.

- Obtain the least squares estimates of β_0 and β_1 and state the estimated regression function.
- Plot the estimated regression function and the data. Does the estimated regression function appear to fit the data well?
- Obtain a point estimate of the mean freshman GPA for students with ACT test score $X = 30$.
- What is the point estimate of the change in the mean response when the entrance test score increases by one point?
- Obtain the residuals e_i and the sum of the squared residuals.
- Obtain point estimates of σ^2 and σ . In what units is each of them expressed?

```
library(tidyverse)
```

```
## -- Attaching packages --- tidyverse 1.3.0 --
```

```
## v ggplot2 3.3.2      v purrr   0.3.4
## v tibble  3.0.3      v dplyr   1.0.2
## v tidyr   1.1.2      v stringr 1.4.0
## v readr   1.3.1      v forcats 0.5.0
```

```
## -- Conflicts ----- tidyverse_conflicts() --
```

```
## x dplyr::filter() masks stats::filter()
## x dplyr::lag()    masks stats::lag()
```

```
aaa <- read_tsv("./data/CH01PR19.txt")
```

```
## Parsed with column specification:  
## cols(  
##   `3.897`    21` = col_character()  
## )
```

```
aaa
```

```
## # A tibble: 119 x 1  
##   `3.897`    21`  
##   <chr>  
## 1 3.885    14  
## 2 3.778    28  
## 3 2.540    22  
## 4 3.028    21  
## 5 3.865    31  
## 6 2.962    32  
## 7 3.961    27  
## 8 0.500    29  
## 9 3.178    26  
## 10 3.310    24  
## # ... with 109 more rows
```