

HIV: DIAGNOSIS

- Encourage patient and his/her partner/s and children to test for HIV.
- If HIV self-screening test done, confirm results with routine tests below.

Obtain informed consent

- Educate about HIV, methods of HIV transmission, risk factors, treatment and benefits of knowing one's HIV status.
- Explain test procedure and that it is completely voluntary.
- Obtain consent (children < 12 years need parental, guardian or caregiver consent). If consent is granted, test using the 'three-test kit HIV test' approach:

Do first HIV rapid screening test on fingerprick blood.

Screening test reactive

Using a different rapid test, do confirmatory HIV test 1 on fingerprick blood.

Screening test non-reactive

Confirmatory test 1 reactive
Do confirmatory HIV test 2 on fingerprick blood.

Confirmatory test 1 non-reactive
Immediately repeat the screening rapid HIV test only:

Confirmatory HIV test 2 reactive

Confirmatory HIV test 2 non-reactive

Second screening test reactive

Second screening test non-reactive

Discrepant results. Report as HIV inconclusive¹.
Send blood for an HIV ELISA test and advise patient to return for result within 7 days.

ELISA Reactive

ELISA results inconclusive

ELISA non-reactive

Report HIV test result as positive. Patient has HIV.

- Give routine HIV care at this visit ➤ 111.
- Encourage HIV testing for partner/s and children.
- Use HIV index testing forms, if available.

- HIV cannot be confirmed or excluded at this time.
- Advise patient to repeat rapid HIV tests in 6 weeks.

Report HIV test result as negative. Patient does not have HIV.

Has patient been at risk of HIV infection in past 6 weeks (had new/multiple partners or unprotected sex with partner who has HIV/is HIV unknown)?

Yes

- Explain that s/he may still be in window period². Advise to repeat HIV test after 6 weeks.
- Consider need for PrEP ➤ 106.

No

- Encourage patient to remain negative and advise when to re-test:
 - If sexually active: 6-12 monthly
 - If pregnant: at every antenatal visit.
 - If on PrEP or breastfeeding, retest 3 monthly.

Offer referral for male circumcision to decrease risk of HIV infection.

Support

- Ensure patient understands test result and knows where and when to access further care.
- Encourage patient to follow safe sex practices. Demonstrate and give male/female condoms.

¹If pregnant in labour, manage baby as high-risk until mother's status confirmed. ²The window period is the time between HIV infection and the point when a test can accurately pick up HIV in the blood.