

Manage ART medication interactions

- Ask patient if s/he is taking any over-the-counter or herbal/traditional medications.
- If on or needing contraception: reassure that there are no interactions with DTG. If on other ART regimen, assess eligibility to switch to DTG ↗ 117, otherwise manage below.
- For other interactions: check SAMF, EMGuidance app, Liverpool HIV iChart app, use web-based interaction checker (see QR code) or discuss with HIV expert/hotline ↗ 178.



Check for HIV
medication
interactions

Assess and manage common medication interactions:

If on this ART medication	Check for interacting medications	Adjust medications and/or doses
Dolutegravir (DTG)	Rifampicin	• Increase DTG dose to 50mg 12 hourly. If on TLD or ALD fixed dose combination tablet, add DTG 50mg 12 hours after TLD or ALD dose. Continue this dose until 2 weeks after TB treatment completed, <i>or</i> • If already on TB treatment and starting ART (patient has never been on ART): consider TEE (TDF + FTC + EFV). Switch to DTG-based regimen 2 weeks after TB treatment complete.
	Anticonvulsants: • Carbamazepine • Phenytoin	• Avoid giving carbamazepine or phenytoin together with DTG: - Switch to lamotrigine ↗ 149. If unable to use lamotrigine, consider valproate ↗ 149. Avoid valproate if woman of child-bearing potential. - If unable to switch anticonvulsant and patient to remain on carbamazepine, increase dose of DTG to give 50mg 12 hourly.
	Iron and/or calcium	• If taking iron only, advise to take iron and DTG together with food. • If taking calcium only, advise to take calcium and DTG together with food. • If taking iron <i>and</i> calcium, advise to take DTG and calcium together with food, then to take iron at least 4 hours later.
	Zinc	Advise to take zinc at least 6 hours before <i>or</i> 2 hours after DTG.
	Magnesium/aluminium (antacids)	Advise to take antacid at least 6 hours before <i>or</i> 2 hours after DTG.
	Metformin	Avoid giving more than 500mg metformin 12 hourly. If diabetes uncontrolled, move to step 2 (start glimepiride) ↗ 131.
Lopinavir/ritonavir (LPVr)	Rifampicin	• Assess eligibility to switch to DTG ↗ 117. If not eligible for DTG switch, gradually increase dose of LPVr according to ALT ↗ 97.
	Anticonvulsants: • Carbamazepine • Phenytoin	• Assess eligibility to switch to DTG ↗ 117. If switching to DTG, continue to manage as above (see dolutegravir row). • If not eligible for DTG switch, avoid giving carbamazepine or phenytoin together with LPVr: - Switch anticonvulsant to lamotrigine and double lamotrigine dose ↗ 149. If unable to use lamotrigine, consider valproate ↗ 149. Avoid valproate if woman of child-bearing potential. If unsure, discuss with HIV expert/hotline ↗ 178.
	Oral contraceptive	Avoid with LPVr. Assess eligibility to switch to DTG ↗ 117. If not eligible to switch to DTG, use instead an IUD, subdermal implant or injectable <i>and</i> condoms.
	Fluticasone/budesonide	Avoid with LPVr. Assess eligibility to switch to DTG ↗ 117. If not eligible to switch to DTG, use instead beclomethasone 12 hourly.
Atazanavir/ritonavir (ATVr)	Rifampicin	• Avoid ATVr. Assess eligibility for switch to DTG ↗ 117. • If not eligible for DTG switch, discuss with TB expert/hotline to switch rifampicin to rifabutin <i>or</i> switch ATVr to LPVr ↗ 178.
	Carbamazepine or phenytoin	• Assess eligibility to switch to DTG ↗ 117. If switching to DTG, also adjust choice of anticonvulsant as for LPVr/r above. • If unable to switch to DTG, discuss with HIV hotline ↗ 178
	Oral contraceptive	Avoid with ATVr. Assess eligibility to switch to DTG ↗ 117. If not eligible to switch to DTG, use instead an IUD, subdermal implant or injectable <i>and</i> condoms.
	Fluticasone/budesonide	Avoid with ATVr. Assess eligibility to switch to DTG ↗ 117. If not eligible to switch to DTG, use instead beclomethasone 12 hourly.
Efavirenz (EFV)	Bedaquiline	Avoid EFV. Switch to DTG ↗ 117.
	• Oral contraceptive • Subdermal implant	• Avoid giving these contraceptives together with EFV. Assess eligibility to switch to DTG ↗ 117. • If not eligible to switch to DTG, use instead IUD or injectable <i>and</i> condoms.
Zidovudine (AZT)	Linezolid	Avoid AZT. Discuss with HIV expert/hotline ↗ 178.

ATVr - atazanavir/ritonavir; AZT - zidovudine; EFV - efavirenz; FTC - emtricitabine; LPVr - lopinavir/ritonavir; TDF - tenofovir; TEE - TDF + FTC + EFV; TLD - TDF + 3TC + DTG; 3TC - lamivudine