

SCREEN ALL PATIENTS FOR COVID-19 AND TB

- Health care workers need to wear a surgical or N95 mask. Patients need to wear cloth or surgical masks and keep 1-2m apart from each other. Ensure queues are distanced.
- Have 70% alcohol-based hand sanitiser or soap and water handwashing stations available for all patients entering facility.
- Ensure a separate patient pathway for patients suspected of having COVID-19. All waiting areas need to be well-ventilated (open doors and all windows) or outside.
- Ensure triage station has a supply of surgical masks to give to symptomatic patients and patient information leaflets for close contacts¹.

If patient known with COVID-19 and returning with worsening symptoms, fast track this patient:
Give surgical mask and send patient to separate area identified for emergency care of COVID-19 patients for urgent attention → 40.

If patient is not known to be COVID-19 positive, screen for the following symptoms:

Ask each patient if s/he has had **new onset** of any of the following in the last 14 days:

- | | | |
|---|--|---|
| • Shortness of breath or difficulty breathing | • Sore throat | • Loss of sense of smell or change in sense of taste |
| • Cough | • Headache with blocked/runny nose or sneezing | • Is/he is known with asthma or COPD with chronic symptoms: worsening of cough or breathing |

Yes to any

No to all

Consider as patient with suspected COVID-19

- Give patient a surgical mask to wear.
- Does patient have shortness of breath or difficulty breathing?

Yes

No

Send patient to separate area identified for emergency care of COVID-19 patients for urgent attention → 40.

- Ensure patients sit 1-2m apart.
- Advise on cough and hand hygiene, and if available, patient to have a rapid antigen test → 40.
- If cough present, arrange to collect 1 sputum sample for TB NAAT to exclude TB → 92.

- Send patient to attend normal waiting area.
- Ask patients to sit 1-2m apart if possible.

- If TB symptoms other than cough (unexplained weight loss > 1.5kg in a month, drenching night sweats or fever), arrange to collect 1 sputum sample for TB NAAT → 92.
- If no TB symptoms, assess for TB preventive treatment (TPT) → 89 if any of:
 - TB contact² (repeat course of TPT for each new TB contact)
 - HIV positive (if not had TPT before), including HIV positive pregnant patient
 - Silicosis

Manage symptoms as on symptom pages.

¹Close COVID-19 contact is when a person has had face-to-face (within 1 metre) contact with someone with COVID-19, or has been in a closed environment (like room or vehicle) with someone with COVID-19 for at least 15 minutes. ²A TB contact refers to a patient who shared an enclosed space (at work, socially, in a hostel, or in a household setting), for ≥ 1 night or for frequent/extended daytime periods, with an adult/adolescent with pulmonary TB ("index patient"), during the 3-month period before the index patient started their TB treatment.