

ADDRESS THE PATIENT'S GENERAL HEALTH

Assess the patient's general health at every visit.

Assess	When to assess	Note
Symptoms	Every visit	Manage symptoms as on symptom pages.
TB	Every visit	If current cough (any duration), weight loss, night sweats or fever, exclude TB ↗ 92. Also assess need for TB preventive treatment (TPT) ↗ 89.
Family planning	Every visit	- Assess patient's contraceptive needs ↗ 154 and pregnancy plans. If pregnant, give antenatal care ↗ 161. - If HIV positive and planning pregnancy, advise patient to use contraception until viral load lower is suppressed¹.
Sexual health	Every visit	- Ask about genital symptoms ↗ 49 and sexual problems ↗ 58. - If risky sexual behaviour: new or multiple partner/s, uses condoms unreliably, has sex under influence of alcohol/drugs, give safe sex advice.
Depression	Every visit	In the past month, has patient: 1) felt down, depressed, hopeless or 2) felt little interest or pleasure in doing things? If yes to either ↗ 143.
Alcohol/drug use	Every visit	In the past year, has patient: 1) drunk ≥ 4 drinks ² /session, 2) used illegal drugs or 3) misused prescription or over-the-counter medications? If yes to any ↗ 142.
Smoking	Every visit	If patient smokes, encourage to stop ↗ 141.
Older person risk	If > 65 years: at every visit	- If patient has a change in function, check for symptoms suggesting a cause: fever ↗ 24, urinary symptoms ↗ 59, confusion ↗ 85. - Consider using lower medication doses (give full doses of antibiotics and ART). Avoid unnecessary medications. Discuss with doctor if patient on diazepam, amitriptyline, theophylline, codeine, ibuprofen, amlodipine or fluoxetine or is using ≥ 5 medications. - If memory problems and disorientation for at least 6 months, consider dementia ↗ 148.
Weight (BMI)	Yearly	- BMI = weight (kg) ÷ height (m) ÷ height (m). - If BMI > 25 ↗ 127. If BMI < 18.5, refer for nutritional support.
BP	First visit, then depending on result	Check BP: if ≥ 140/90 ↗ 132. If pregnant and BP ≥ 140/90 ↗ 159.
CVD risk	If ≥ 40 years or ≥ 2 risk factors	- Assess CVD risk ↗ 127 at first visit, then depending on risk. - Risk factors: smoking, BMI > 25, waist circumference > 80cm (woman) or 94cm (man), hypertension, diabetes, cholesterol > 5.2, parent/sibling with early onset CVD³ (man < 55 years or woman < 65 years).
Diabetes risk	At first visit if: - If ≥ 45 years or - If BMI ≥ 25 and ≥ 1 other risk factor	- If not known diabetic, check glucose ↗ 17. - Risk factors: physical inactivity, hypertension, parent or sibling with diabetes, polycystic ovarian disease, Indian ethnicity, cardiovascular disease, diabetes during pregnancy or previous big baby > 4000g, previous impaired glucose tolerance or impaired fasting glucose or TB in past year.
HIV	- If status unknown - If sexually active: 6-12 monthly - If pregnant: every antenatal visit - If breastfeeding: 3 monthly	Test for HIV ↗ 110.
Cervical screen (if woman)	When needed	- HIV negative: do 3 cervical screens, each 10 years apart from age 30 ↗ 55. - HIV positive: do cervical screen at HIV diagnosis (regardless of age), then 3-yearly ↗ 55.
Breast check (if woman)	- First visit - On contraceptive or hormone therapy: yearly - If > 40 years: 6 monthly	- Check for lumps in breasts ↗ 43 and axillae ↗ 25. - If on hormone therapy, refer for mammogram at initiation if available.

Continue to manage the patient's general health → 11.

¹Viral load < 50. ²One drink is 1 tot of spirits, or 1 small glass (125mL) of wine or 1 can/bottle (330mL) of beer. ³Cardiovascular disease (CVD) includes ischaemic heart disease, peripheral vascular disease and stroke/TIA.