



Decide when to change intensive phase to continuation phase

• If patient on shorter regimen:



• If patient on longer regimen:



Decide on treatment outcome

Record treatment outcome based on culture conversion, number of consecutive negative cultures, duration of minimum treatment and clinical status.

Cure

- TB culture converted
- ≥ 3 consecutive² negative cultures in continuation phase
- If shorter course: ≥ 9 months of treatment
- If longer course: ≥ 18 months of treatment
- Not clinically worsening

Treatment completed (success)

- TB culture converted
- < 3 consecutive² negative cultures in continuation phase
- If shorter course: ≥ 9 months of treatment
- If longer course: ≥ 18 months of treatment
- Not clinically worsening

Treatment failure

- Failure of month 4 TB culture to convert by month 6
- ≥ 2 cultures positive in continuation phase and clinically worsening
- Treatment stopped on clinical grounds or by instruction from PCAC
- ≥ 2 new drugs added to regimen due to poor clinical response

• **Other outcomes include:** 'Loss to follow up' (treatment interruption for ≥ 2 months), 'Moved', 'Transferred out', 'Died', 'Still on treatment', 'Not evaluated'.

¹Slow clinical response: poor weight gain, ongoing TB symptoms, poor improvement on chest x-ray or delayed smear/culture conversion. ²Cultures need to have been done in a row, at least 30 days apart.