

REVIEW THE PATIENT ON POST-EXPOSURE PROPHYLAXIS (PEP)

Review patient within 3 days, at 4 weeks and 4 months.


- Check adherence and ask about side effects from HIV PEP ↗ 116. Advise patient of side effects and to return promptly if they occur. Advise patient to use condoms for 4 months until results confirmed.
- If sexual assault ↗ 88. If case of human bite: repeat only HBsAg (at 4 months) from table below, use HBsAbs results to continue to give only hepatitis B prophylaxis below.
- Check bloods according to table and review results as below:

Assess	When to assess	Note
HIV rapid test	Repeat HIV rapid test at 4 weeks and 4 months	• Encourage to test for HIV ↗ 110. - If HIV negative, assess the need for PrEP ↗ 106. - If HIV positive, give routine HIV care ↗ 111.
Hepatitis B surface antigen (HBsAg)	At 4 months	If positive ↗ 120.
Hepatitis C antibody	Do only if source hepatitis C antibody positive: first visit	If positive, refer. If negative, do hepatitis C PCR at 6 weeks.
Hepatitis C PCR	If exposed hepatitis C antibody negative and source positive: at 6 weeks	If positive, refer.
Syphilis (if sexual exposure)	Do only if source syphilis positive/unknown: first visit, 4 months	If positive ↗ 53.
Creatinine (eGFR) result	Check baseline results at 3-day follow up	If eGFR ≤ 50, stop TDF + 3TC (or TDF + FTC), give instead AZT 300mg + 3TC 150mg 12 hourly and check FBC and differential count.
Full blood count	If on AZT: repeat at 2 weeks	If Hb ≤ 8 or neutrophils ≤ 1.0, discuss with HIV hotline ↗ 178 or specialist.
Source blood results (if done)	-	If source HBsAg or hepatitis C antibody positive, refer source patient. If syphilis positive ↗ 53.

AZT – zidovudine; FTC – emtricitabine; TDF – tenofovir; 3TC – lamivudine.

Continue to give hepatitis B prophylaxis according to vaccination status

Has exposed patient/health worker previously received 3 doses of hepatitis B vaccine?



¹If giving both hepatitis B vaccine and immunoglobulin, give at different sites. If immunoglobulin not available, refer to secondary care within 7 days but ideally within 24-72 hours after exposure. ²If health worker, repeat HBsAb titre 1-2 months after the last vaccine dose to ensure HBsAb ≥ 10.