

CVD risk: diagnosis if cholesterol not available

Step 3 If no established CVD and cholesterol *not* available, calculate the patient's 10-year CVD risk using the chart based on BMI instead of cholesterol:


- Use the patient's sex, age, BMI, systolic BP and smoking status to work out what colour block they fall into.

Step 4 Explain to the patient what his/her risk of heart attack or stroke might be over next 10 years:

- CVD risk is < 5%:** this means there is less than a 1 in 20 chance that s/he may have a heart attack or stroke over the next 10 years.
- CVD risk is 5-10%:** this means there is between a 1 in 20 and a 1 in 10 chance that s/he may have a heart attack or stroke over the next 10 years.
- CVD risk is 10-20%:** this means there is between a 1 in 10 and a 1 in 5 chance that s/he may have a heart attack or stroke over the next 10 years.
- CVD risk is > 20%:** this means there is more than 1 in 5 chance that s/he may have a heart attack or stroke over the next 10 years.

Step 5 Use the patient's CVD risk to decide treatment and frequency of follow-up:

- If CVD risk factor or a CVD risk $\geq 10\%$, manage the CVD risk $\rightarrow 129$.
- If no CVD risk factors and CVD risk < 10%, reassess CVD risk after 5 years.



¹Adapted from WHO cardiovascular disease risk non-laboratory-based Southern Sub-Saharan Africa. From: HEARTS technical package for cardiovascular disease management in primary health care: risk based CVD management. World Health Organization. Geneva, 2020.