

OTHER GENITAL SYMPTOMS

- First assess and advise the patient ➔ 49.
- Then manage according to main symptom:

Lumps or warts

Painless, raised skin coloured growths with round/cauliflower-like surface (skin around genitals, anus or cervix)



© University of Cape Town

Genital warts likely

- If warts atypical/fleshy/wet, test for syphilis. If positive ➔ 53.
- Arrange a cervical screen for patient/partner if needed ➔ 55.
- Offer to arrange medical male circumcision for patient/partner.
- Reassure that most warts resolve spontaneously within 2 years.
- Refer to gynaecology, urology or sexual health services if:
 - Warts > 10mm
 - Numerous lesions
 - Warts inside vagina, involving cervix or urethra
 - Pregnant with large warts
 - Bleeding or infected warts

Papules with central dent



© University of Cape Town

Molluscum contagiosum likely

- Reassure that most papules resolve spontaneously within 9-12 months.
- Apply **tincture of iodine BP** topically with an applicator to the core of the lesions.
- If no response to treatment, refer.

Itchy rash in pubic area

Intensely itchy bites
May see lice or nits (size of a pinhead) in pubic and peri-anal areas

Pubic lice (pediculosis) likely

- Apply **benzyl benzoate 25%** lotion to affected area for 24 hours. Avoid mucous membranes, face and eyes, urethral opening and raw areas. Repeat treatment after 1 week.
- Advise to shave genital area.
- Treat all sexual partners even if asymptomatic.
- Before treatment, wash and thoroughly dry clothing and linen that may have been contaminated within past 2 days.
- For itch, give **chlorphenamine** 4mg 8 hourly as needed for up to 10 days.

If eyelashes/eyebrows involved, **pediculosis of eyelashes/eyebrows** likely.

Apply yellow **petroleum jelly** to eyelid margins to (cover eyelashes) and eyebrows daily for 10 days to smother lice/nits. Advise patient to avoid getting petroleum in eye.

Itch worse at night, with red papules and nodules



© University of Cape Town

Genital scabies likely

- Apply **benzyl benzoate 25%** lotion from neck to soles of feet and rub in well:
 - Leave on for 24 hours, then wash off with soap and water.
 - If severe, repeat once after 24 hours or within 5 days.
- If no better, apply **permethrin 5%** lotion at night from neck to soles of feet. Wash off after 8-12 hours. Repeat after 1 week if needed.
- For itch, give **chlorphenamine** 4mg 8 hourly for up to 10 days. If mild itch, use only at night.
- Advise can return to work after first treatment.
- Treat all household contacts and sexual partners at the same time, even if asymptomatic.
- Wash recently used linen and clothing in very hot water and dry well. Expose to direct sunlight.

If scratch marks infected (pus/red/swollen/crusts), also treat for likely **impetigo** ➔ 78.

¹Avoid in pregnancy and breastfeeding.