

## INH MONO-RESISTANT TB: ROUTINE CARE

**TB tests changing from 'Xpert Ultra' to 'TB NAAT'**  
(NAAT = nucleic acid amplification test and includes Xpert as well as newer TB tests).

### Assess the patient with INH mono-resistant TB

| Assess                                         | When to assess                                                                                                                                              | Note                                                                                                                                                                                                                                                                                                                                                                                                                                                                          |
|------------------------------------------------|-------------------------------------------------------------------------------------------------------------------------------------------------------------|-------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
| Registration                                   | At diagnosis                                                                                                                                                | Ensure patient record completed and captured on TIER.net.                                                                                                                                                                                                                                                                                                                                                                                                                     |
| TB contacts                                    | At diagnosis                                                                                                                                                | Advise that all TB contacts <sup>1</sup> visit the clinic for TB screening and testing or ensure CHW does a home visit for TB screening and testing.                                                                                                                                                                                                                                                                                                                          |
| Alcohol/drug use                               | At diagnosis                                                                                                                                                | In the past year, has patient: 1) drunk $\geq 4$ drinks <sup>3</sup> /session, 2) used illegal drugs or 3) misused prescription or over-the-counter medications? If yes to any $\rightarrow$ 142.                                                                                                                                                                                                                                                                             |
| Symptoms                                       | Every visit                                                                                                                                                 | <ul style="list-style-type: none"> <li>If respiratory rate <math>\geq 30</math>, breathless at rest or while talking, prominent use of breathing muscles, drowsy/confused, coughs up <math>\geq 1</math> tablespoon fresh blood, neck stiffness, persistent vomiting or new weakness of arm/leg, give urgent attention <math>\rightarrow</math> 92.</li> <li>Expect gradual improvement on TB treatment. If symptoms worsen or do not get better, refer to doctor.</li> </ul> |
| Adherence                                      | Every visit                                                                                                                                                 | Request patient brings all medication to each visit. Check adherence on the TB card.                                                                                                                                                                                                                                                                                                                                                                                          |
| Side effects                                   | Every visit                                                                                                                                                 | Ask about side effects of treatment $\rightarrow$ 97.                                                                                                                                                                                                                                                                                                                                                                                                                         |
| Family planning                                | Every visit                                                                                                                                                 | <ul style="list-style-type: none"> <li>Encourage patient to avoid pregnancy during treatment, assess patient's contraceptive needs <math>\rightarrow</math> 154. If pregnant <math>\rightarrow</math> 159.</li> <li>Avoid oral contraceptive and subdermal implant<sup>2</sup> on TB treatment, use instead injectable or IUD <i>plus</i> condoms. No need to change interval between injectable.</li> </ul>                                                                  |
| Depression                                     | Every visit                                                                                                                                                 | In the past month, has patient: 1) felt down, depressed, hopeless or 2) felt little interest or pleasure in doing things? If yes to either $\rightarrow$ 143.                                                                                                                                                                                                                                                                                                                 |
| Palliative care                                | If advanced TB disease                                                                                                                                      | If patient is in bed or chair for 50% or more of the day or dependent on others for most care, also give palliative care $\rightarrow$ 170.                                                                                                                                                                                                                                                                                                                                   |
| Weight and BMI                                 | Every visit                                                                                                                                                 | <ul style="list-style-type: none"> <li>Expect weight gain on treatment and adjust TB treatment dose <math>\rightarrow</math> 96. If losing weight, refer to doctor.</li> <li>BMI = weight (kg) <math>\div</math> height (m) <math>\div</math> height (m). If <math>&lt; 18.5</math>, refer for nutritional support.</li> </ul>                                                                                                                                                |
| Glucose                                        | At diagnosis                                                                                                                                                | <ul style="list-style-type: none"> <li>If known diabetes, assess glucose control more often <math>\rightarrow</math> 130.</li> <li>If not known with diabetes, check glucose <math>\rightarrow</math> 17.</li> </ul>                                                                                                                                                                                                                                                          |
| HIV                                            | At diagnosis and every visit                                                                                                                                | If $> 3$ months since last HIV test, test for HIV $\rightarrow$ 110. If HIV positive, give routine HIV care and ART $\rightarrow$ 111. If on ART, adjust medication/dosing $\rightarrow$ 97.                                                                                                                                                                                                                                                                                  |
| TB microscopy (smear) and culture <sup>4</sup> | At diagnosis                                                                                                                                                | Register as smear negative or smear positive depending on result.                                                                                                                                                                                                                                                                                                                                                                                                             |
|                                                | Monthly                                                                                                                                                     | <ul style="list-style-type: none"> <li>If still culture positive at 3 months, request DST on that same positive specimen.</li> <li>If still culture positive at 4 months, discuss with specialist or refer to drug-resistant TB unit.</li> <li>If negative smear/culture becomes positive, request DST on that same positive specimen.</li> </ul>                                                                                                                             |
| DST                                            | <ul style="list-style-type: none"> <li>At diagnosis</li> <li>If culture positive at 3 months</li> <li>If negative smear/culture becomes positive</li> </ul> | <ul style="list-style-type: none"> <li>If resistant to INH only: if still culture positive at 4 months, discuss with specialist or refer to drug-resistant TB unit.</li> <li>If resistant to rifampicin, diagnose <b>rifampicin-resistant TB (RR-TB)</b> and give routine care <math>\rightarrow</math> 99.</li> </ul>                                                                                                                                                        |
| TB NAAT                                        | If needed                                                                                                                                                   | If INH resistance detected $> 28$ days after start of DS-TB treatment, send 1 further sputum sample for TB NAAT to confirm that there is no resistance to rifampicin.                                                                                                                                                                                                                                                                                                         |
| Treatment outcome                              | At completion of TB treatment                                                                                                                               | Decide on treatment outcome $\rightarrow$ 98.                                                                                                                                                                                                                                                                                                                                                                                                                                 |

### Advise and treat the patient with INH mono-resistant TB $\rightarrow$ 96.

<sup>1</sup>Close TB contact: any person who shared an enclosed space (social/work/congregate/household setting) with an adolescent or adult with pulmonary TB (index patient) for  $> 15$  minutes in 24 hours during the 3 months before index patient started TB treatment. <sup>2</sup>If patient already has subdermal implant, advise additional non-hormonal method (copper IUD or condoms) until 4 weeks after completing TB treatment. <sup>3</sup>One drink is 1 tot of spirits, or 1 small glass (125mL) of wine or 1 can/bottle (330mL) of beer. <sup>4</sup>Make every effort to obtain sputum. If difficulty, try in early morning or arrange for induced sputum.