# Cambridge International AS & A Level

CANDIDATE NAME					
CENTRE NUMBER			CANDIDATE NUMBER		

MATHEMATICS

October/November 2023

1 hour 50 minutes

9709/13

You must answer on the question paper.

You will need: List of formulae (MF19)

#### **INSTRUCTIONS**

Answer all questions.

Paper 1 Pure Mathematics 1

- Use a black or dark blue pen. You may use an HB pencil for any diagrams or graphs.
- Write your name, centre number and candidate number in the boxes at the top of the page.
- Write your answer to each question in the space provided.
- Do not use an erasable pen or correction fluid.
- Do not write on any bar codes.
- If additional space is needed, you should use the lined page at the end of this booklet; the question number or numbers must be clearly shown.
- You should use a calculator where appropriate.
- You must show all necessary working clearly; no marks will be given for unsupported answers from a calculator.
- Give non-exact numerical answers correct to 3 significant figures, or 1 decimal place for angles in degrees, unless a different level of accuracy is specified in the question.

#### **INFORMATION**

- The total mark for this paper is 75.
- The number of marks for each question or part question is shown in brackets [ ].

## **BLANK PAGE**

ŀ	Find the equation of the curve.
•	
•	
•	
•	
•	
•	
•	
•	

(a)	Find the y-coordinates of $A$ and $B$ , expressing your answers in terms of surds.	[2]
(b)	Find the equation of the circle which has $AB$ as its diameter.	[2]
<b>(b)</b>		[2]
(b)	Find the equation of the circle which has $AB$ as its diameter.	[2]
(b)	Find the equation of the circle which has $AB$ as its diameter.	[2]
(b)	Find the equation of the circle which has $AB$ as its diameter.	[2]
(b)	Find the equation of the circle which has $AB$ as its diameter.	[2]
(b)	Find the equation of the circle which has $AB$ as its diameter.	[2]
(b)	Find the equation of the circle which has $AB$ as its diameter.	[2]
(b)	Find the equation of the circle which has $AB$ as its diameter.	[2]
(b)	Find the equation of the circle which has $AB$ as its diameter.	[2]
(b)	Find the equation of the circle which has $AB$ as its diameter.	[2]
(b)	Find the equation of the circle which has $AB$ as its diameter.	[2]

3	(a)	Show	that	the	equa	ıtion
---	-----	------	------	-----	------	-------

**(b)** 

$5\cos\theta - \sin\theta\tan\theta + 1 = 0$
may be expressed in the form $a\cos^2\theta + b\cos\theta + c = 0$ , where a, b and c are constants to be found. [3]
Hence solve the equation $5\cos\theta - \sin\theta\tan\theta + 1 = 0$ for $0 < \theta < 2\pi$ . [4]


(i)	$(1+2x)^5.$	
		•••••
(ii)	$(1-ax)^6$ , where a is a constant.	
	pansion of $(1 + 2x)^5(1 - ax)^6$ , the coefficient of $x^2$ is $-5$ . If the possible values of $a$ .	

(a)	Find the possible values of the constant $p$ .	[3]
		•••••
		•••••
		•••••
		•••••
<b>(b)</b>	One of the values of $p$ found in $(a)$ is a negative fraction.	
<b>(b)</b>	One of the values of $p$ found in (a) is a negative fraction.  Use this value of $p$ to find the sum to infinity of this progression.	[4]
<b>(b)</b>		[4]
(b)		[4]

Find the possib	ble values of $c$ and the corresponding coordinates of $P$ .	[7]

	State the range of f.	[:
<b>(b)</b>	Obtain an expression for $f^{-1}(x)$ and state the domain of $f^{-1}$ .	[
	e function g is defined by $g(x) = 2x - 2$ for $x > 0$ .	
Γhe		
	Obtain a simplified expression for $gf(x)$ .	[2
	Obtain a simplified expression for $gf(x)$ .	[2
	Obtain a simplified expression for $gf(x)$ .	[2
	Obtain a simplified expression for gf(x).	[2
	Obtain a simplified expression for gf(x).	[2

8



The diagram shows part of the graph of  $y = \sin(a(x+b))$ , where a and b are positive constants.

(a)	State the value of $a$ and one possible value of $b$ .	[2]
		•••••
		• • • • • •

Another curve, with equation y = f(x), has a single stationary point at the point (p, q), where p and q are constants. This curve is transformed to a curve with equation

$$y = -3f\left(\frac{1}{4}(x+8)\right).$$


For the transformed curve, find the coordinates of the stationary point, giving your answer terms of $p$ and $q$ .	ir [3]
	••••
	••••
	••••
	• • • •

**(b)** 

A curve has equation  $y = 2x^{\frac{1}{2}} - 1$ .

	y = mx + c.	[3]
	point is moving along the curve $y = 2x^{\frac{1}{2}} - 1$ in such a way that at A the rate of increpordinate is $3  \mathrm{cm  s^{-1}}$ .	ease of the
<b>(b)</b>	Find the rate of increase of the <i>y</i> -coordinate at <i>A</i> .	[2]
way	A the moving point suddenly changes direction and speed, and moves down the norma	l in such a
way	A the moving point suddenly changes direction and speed, and moves down the normal that the rate of decrease of the y-coordinate is constant at 5 cm s <sup>-1</sup> .  As the point moves down the normal, find the rate of change of its x-coordinate.	al in such a
	At the moving point suddenly changes direction and speed, and moves down the normal that the rate of decrease of the y-coordinate is constant at $5 \mathrm{cm}\mathrm{s}^{-1}$ .  As the point moves down the normal, find the rate of change of its x-coordinate.	[3]
way	A the moving point suddenly changes direction and speed, and moves down the normal that the rate of decrease of the y-coordinate is constant at 5 cm s <sup>-1</sup> .  As the point moves down the normal, find the rate of change of its x-coordinate.	[3]
way	A the moving point suddenly changes direction and speed, and moves down the normal that the rate of decrease of the y-coordinate is constant at 5 cm s <sup>-1</sup> .  As the point moves down the normal, find the rate of change of its x-coordinate.	[3]

**10** 




The diagram shows points A, B and C lying on a circle with centre O and radius r. Angle AOB is 2.8 radians. The shaded region is bounded by two arcs. The upper arc is part of the circle with centre O and radius r. The lower arc is part of a circle with centre C and radius R.

(a)	State the size of angle ACO in radians.	[1]
		••••
		••••
		••••
<b>(b)</b>	Find $R$ in terms of $r$ .	[1]
		••••
		••••
		••••
		••••
		••••
		••••
		••••
		••••
		••••

)	Find the area of the shaded region in terms of $r$ .	[7]
		•••••
		••••••
		••••••

11



The diagram shows part of the curve with equation  $y = x + \frac{2}{(2x-1)^2}$ . The lines x = 1 and x = 2 intersect the curve at P and Q respectively and R is the stationary point on the curve.

Verify that the x-coordinate of R is $\frac{3}{2}$ and find the y-coordinate of R.	[4]

Find the exact value of the area of the shaded region.	
	•••••

### **Additional Page**

If you use the following lined page to complete the answer(s) to any question(s), the question number(s) must be clearly shown.

Permission to reproduce items where third-party owned material protected by copyright is included has been sought and cleared where possible. Every reasonable effort has been made by the publisher (UCLES) to trace copyright holders, but if any items requiring clearance have unwittingly been included, the publisher will be pleased to make amends at the earliest possible opportunity.

To avoid the issue of disclosure of answer-related information to candidates, all copyright acknowledgements are reproduced online in the Cambridge Assessment International Education Copyright Acknowledgements Booklet. This is produced for each series of examinations and is freely available to download at www.cambridgeinternational.org after the live examination series.

Cambridge Assessment International Education is part of Cambridge Assessment. Cambridge Assessment is the brand name of the University of Cambridge Local Examinations Syndicate (UCLES), which is a department of the University of Cambridge.