

[Turn over

BLANK PAGE

- 1 Two particles P and Q , of masses m kg and 0.3 kg respectively, are at rest on a smooth horizontal plane. P is projected at a speed of 5 m s^{-1} directly towards Q . After P and Q collide, P moves with a speed of 2 m s^{-1} in the same direction as it was originally moving.

(a) Find, in terms of m , the speed of Q after the collision. [2]

.....

.....

.....

.....

.....

.....

.....

.....

.....

After this collision, Q moves directly towards a third particle R , of mass 0.6 kg, which is at rest on the plane. Q is brought to rest in the collision with R , and R begins to move with a speed of 1.5 m s^{-1} .

(b) Find the value of m . [2]

.....

.....

.....

.....

.....

.....

.....

.....

.....

2 A particle P of mass 0.4 kg is projected vertically upwards from horizontal ground with speed 10 m s^{-1} .

(a) Find the greatest height above the ground reached by P . [2]

.....

.....

.....

.....

.....

.....

.....

When P reaches the ground again, it bounces vertically upwards. At the first instant that it hits the ground, P loses 7.2 J of energy.

(b) Find the time between the first and second instants at which P hits the ground. [4]

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

- $$s = t^{\frac{5}{2}} - \frac{15}{4}t^{\frac{3}{2}} + 6.$$

[4]

This image shows a single page of white paper with horizontal ruling lines. The lines are evenly spaced and run across the width of the page. There are no margins, text, or other markings on the paper.

A velocity-time graph for a car starting from rest. The vertical axis is labeled $v \text{ (m s}^{-1}\text{)}$ and the horizontal axis is labeled $t \text{ (s)}$. The graph consists of several segments: a straight line from $(0, 0)$ to $(3, 0.9)$; a horizontal line from $t = 3$ to $t = 9$ at $v = 0.9$; a straight line from $(9, 0.9)$ to $(10, 0)$; a straight line from $(10, 0)$ to a minimum velocity (indicated by a dashed line to the horizontal axis); and a straight line from the minimum velocity back to the horizontal axis at time T . Dashed lines indicate the coordinates $(3, 0.9)$ and $(9, 0.9)$.

(a) Find the distance travelled by the particle in the first 10 s of its motion. [2]

[illegible]

- (b) Given that $T = 12$, find the minimum velocity of the particle.

[2]

.....

.....

.....

.....

.....

.....

.....

.....

.....

- (c) Given instead that the greatest speed of the particle is 3 m s^{-1} , find the value of T and hence find the average speed of the particle for the whole of the motion.

[4]

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....


.....

.....

.....

.....

.....



Four coplanar forces act at a point. The magnitudes of the forces are F N, 10 N, 50 N and 40 N. The directions of the forces are as shown in the diagram.

- (a) Given that the forces are in equilibrium, find the value of F and the value of θ . [6]

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

- (b) Given instead that $F = 10\sqrt{2}$ and $\theta = 45$, find the direction and the exact magnitude the resultant force. [3]

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

(a) It is given that $\theta = 60^\circ$, the plane BC is rough and the coefficient of friction between Q and the plane BC is 0.7 . The particles are released from rest.

[4]

[illegible]

- Find the magnitude of the acceleration of P as it moves on the plane, and find the value of θ . [4]

This image shows a full page of primary-ruled paper. It features approximately 20 horizontal dashed lines spaced evenly down the page, providing a guide for handwriting practice. The paper is otherwise blank, with no margins, text, or other markings.

- 7 A car of mass 1200 kg is travelling along a straight horizontal road. The power of the car's engine is constant and is equal to 16 kW. There is a constant resistance to motion of magnitude 500 N.

(a) Find the acceleration of the car at an instant when its speed is 20 m s^{-1} . [3]

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

(b) Assuming that the power and the resistance forces remain unchanged, find the steady speed at which the car can travel. [2]

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

The car comes to the bottom of a straight hill of length 316 m, inclined at an angle to the horizontal of $\sin^{-1}(\frac{1}{60})$. The power remains constant at 16 kW, but the magnitude of the resistance force is no longer constant and changes such that the work done against the resistance force in ascending the hill is 128 400 J. The time taken to ascend the hill is 15 s.

- (c) Given that the car is travelling at a speed of 20 m s^{-1} at the bottom of the hill, find its speed at the top of the hill. [6]

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

Additional Page

If you use the following lined page to complete the answer(s) to any question(s), the question number(s) must be clearly shown.

This image shows a single sheet of white paper with horizontal ruling lines. The lines are evenly spaced and run across the width of the page. There are no margins, text, or other markings on the paper.

BLANK PAGE

BLANK PAGE

Permission to reproduce items where third-party owned material protected by copyright is included has been sought and cleared where possible. Every reasonable effort has been made by the publisher (UCLES) to trace copyright holders, but if any items requiring clearance have unwittingly been included, the publisher will be pleased to make amends at the earliest possible opportunity.

To avoid the issue of disclosure of answer-related information to candidates, all copyright acknowledgements are reproduced online in the Cambridge Assessment International Education Copyright Acknowledgements Booklet. This is produced for each series of examinations and is freely available to download at www.cambridgeinternational.org after the live examination series.

Cambridge Assessment International Education is part of Cambridge Assessment. Cambridge Assessment is the brand name of the University of Cambridge Local Examinations Syndicate (UCLES), which is a department of the University of Cambridge.