

AMATH 353: Homework 6

Due April, 20 2018

ID: 1064712

Trent YAROSEVICH

April 20, 2018

Instructor: Jeremy Upsal

Part 1.) With $u(x, 0) = \sin(x)$ and $u_t(x, 0) = xe^{-x^2}$ and $c = 1$ we arrive at the following using the D'Alembert solution:

$$\int_{x-ct}^{x+ct} xe^{-x^2} = \frac{1}{2}e^{-(x-ct)^2} - \frac{1}{2}e^{-(x+ct)^2}$$

Then plugging this into the solution we get (1)

$$u(x, t) = \frac{1}{2}(\sin(x-t) + \sin(x+t)) + \frac{1}{4}e^{-(x-ct)^2} - \frac{1}{4}e^{-(x+ct)^2}$$

This problem relates to the previous two insofar as the solution is composed of the sum of four traveling waves, two of which appear in 8.4, and the other two in 8.5. For me personally, the easiest way to visualize this is to consider the solutions from 8.4 and 8.5 independently, and to rewrite the solution from 8.4 as follows:

$$u(x, t) = \cos(t) \sin(x) \quad (2)$$

And then 8.5:

$$\frac{1}{4}e^{-(x-t)^2} - \frac{1}{4}e^{-(x+t)^2} \quad (3)$$

Equation (2) is a sin wave with amplitude varying over time from -1 to 1, and equation (3) is composed of two traveling wave pulses. The solution in equation (1) above is the sum of these, so there is a sin wave with varying amplitude over time, and two wave pulses are travelling 'through' this wave so to speak. Thus the solution in (3) will behave just like equation (1) except at those regions where it is being disrupted by the two wave pulses. The following figures show the initial sin wave with no disruption from the wave pulses (which are 0 for all x at $t = 0$) and three t values that show how it disrupts the wave.




Figure 1: $t = 0$




Figure 2: $t = 1.5$




Figure 3: $t = 2$




Figure 4: $t = 2.5$

A few observations can be made. In figure 2, we see the varying amplitude of the sin wave is very small, making the disruptions from the wave pulses more obvious than in figure 4, for example. Furthermore we can observe in figure two that the left-moving wave pulse has a negative value, and the right-moving a positive one, thus we can see in figures 2 and 3 that the left moving wave pulse disrupts the sin wave by 'pulling' it down, and the right-moving pulse does the opposite.

Part 2.) From the D'Alembert solution we get:

$$\begin{aligned}
 u(x,t) &= \frac{1}{2}(x-2t+x+2t) + \frac{1}{8} \int_{x-2t}^{x+2t} y^2 dy \\
 u(x,t) &= x + \frac{1}{8} \int_{x-2t}^{x+2t} y^2 dy \\
 u(x,t) &= x + \frac{1}{8} \left(\frac{1}{3}(x+2t)^3 - \frac{1}{3}(x-2t)^3 \right)
 \end{aligned} \tag{4}$$

I'm not sure I understand the question being asked, as we haven't really discussed physical solutions, and I only have the brief overview of it from the notes and a roughly one-sentence discussion of it from the Wednesday 3/28 lecture. Based on this limited discussion and the question, I can only speculate that these initial conditions are problematic for physical solutions because the initial

condition $u(x, t) = x$ becomes insignificant as t increases. So making estimates moving the initial conditions along tangent vectors would probably provide a highly inaccurate result? Other than that I'm not sure what the question is asking. The solution doesn't act like a traveling wave, and just becomes a polynomial when t isn't zero.