AMATH 353: Homework 12 Due May, 18 2018 ID: 1064712

Trent Yarosevich

May 22, 2018

Instructor: Jeremy Upsal

Part 1

a.) Because the PDE in question is homogeneous, we have $x = c(u_o(x_0))t + x_0$. For this equation, c(u(x,t)) = u(x,t). From this we get the following:

$$u(x,0) = u_0(x_0) = e^{-x^2}$$

$$\frac{du}{dt} = 0$$

$$u(x(t),t) = A = e^{-x_0^2}$$
(1)

And the characteristic lines:

$$x = x_0 + te^{-x^2}t = (x - x_0)e^{x^2}$$
 (2)





Figure 1: $x_0 \in [-1.8, 1.8]$

b.)

c.) I wrote a script in matlab to find the roots for $f(x_0) = x_0 - x + te^{-x^2}$, which is included in my HW submission.

d.) and **e.)** Below are the plots, using x_0 values from my rootfinder, for t = 0, 0.6, 0.9, 1.18 and $x \in [-1.8, 1.8]$ with a mesh of values. From the first figure, it is clear that at t = 0 the rootfinder is returning the correct values, yielding the profile of $u_0(x_0) = e^{-x^2}$. As the figures below show, as t increases, the top of the profile has a faster rate of change than the bottom, causeing a 'crashing wave' to form. This is to be expected, since the time velocity of the PDE is depending on both the spatial velocity and the value of u itself, thus the larger u values have larger time velocity.



f.) When my rootfinder is used to plot the values at t=1.5, it appears to get 'stuck' at the point of the wave crashing and falling over. This occurs at the point that the characteristics start to cross, which by closely examining (zooming in) on Figure 1, would seem to happen around t=1.2. Below are two plots using my rootfinder, of t=1.5 and t=2. In both, we see an abrupt change in the value of u, caused by crossing characteristic lines which result in the value of u 'jumping' between two x_0 values produced by my rootfinder.




Figure 6: t = 1.5

Figure 7: t=2