

**MS WRITTEN EXAMINATION IN BIOSTATISTICS,
PART I**

Friday, August 10, 2012: 9:00 AM - 3:00PM

Room: 1305

INSTRUCTIONS:

- This is a **CLOSED BOOK** examination.
- Submit answers to **exactly** 3 out of 4 questions. If you submit answers to more than 3 questions, then only questions 1-3 will be counted.
- Put the answers to different questions on **separate sets of paper**. Write on **one side** of the sheet only.
- Put your code letter, **not your name**, on each page, in the upper right corner.
- Return the examination with a **signed honor pledge form**, separate from your answers.
- You are required to answer **only what is asked** in the questions and not to tell all you know about the topics.

1. Suppose that the random variable X is distributed as Poisson with mean $\mu > 0$, and that given $X = x$, the random variable Y is distributed as Poisson with mean x .

In what follows, derive explicit expressions and simplify them as much as possible. Show *all* your derivations, not just the final answer.

Hint: Conditioning is very helpful in several parts of this problem.

- (a) Find $E[Y]$ and $\text{Var}(Y)$.
- (b) Find $\text{Corr}(X, Y)$.
- (c) Find the conditional probability mass function of X given $Y = 0$. Does it follow a recognizable form? If so, identify it.
- (d) Find $E[X|Y \geq 1]$. Hint: Make use of the result in (c) and of what is known about X .
- (e) Find $E[\frac{Y}{X+1}]$. Verify whether this expectation is less than 1 for all μ . Note: Evaluating an infinite sum is needed to get the final answer.
- (f) Define $T = wX + (1 - w)Y$, where w is a non-random constant, $w \in [0, 1]$. If T is to be used as an estimator of μ , what is the best choice of w ? Justify. Why does the answer make sense?

Points: (a) 3, (b) 3, (c) 5, (d) 5, (e) 5, (f) 4.

2. The glucose tolerance level in patients with diabetes is normally distributed with mean 6 and variance 9. The glucose tolerance level in people without diabetes has a normal distribution with mean 4 and variance 4. When glucose tolerance is used as a test for diabetes, individuals with glucose tolerance level of 5 and greater are classified as having diabetes (that is, the test is considered positive), while individuals with glucose tolerance level below 5 are classified as non-diabetic (that is, the test is considered negative). Let p denote the probability that a randomly chosen person actually has diabetes, $0 < p < 1$.

In what follows, derive explicit expressions and simplify them as much as possible. Show *all* your derivations, not just the final answer.

- (a) What is the probability that a person who has diabetes gets correctly classified as having diabetes according to the result of the glucose tolerance test?
- (b) What is the probability that a person who does not have diabetes gets correctly classified as diabetes free by the glucose tolerance test?
- (c) What is the probability of a positive test result for a randomly chosen person? The answer is a function of p .
- (d) A randomly chosen subject has tested positive. What is the probability that he/she actually has diabetes? The answer is a function of p .
- (e) A randomly chosen subject has a glucose tolerance level of 6. What is the probability that that subject has diabetes? The answer is a function of p .
- (f) Suppose that two glucose tolerance measurements, taken one month apart, on a random diabetic individual, are distributed as bivariate normal, and their correlation is 0.5 (and means and variances are as given above). A randomly chosen diabetic subject tested today has a glucose tolerance value of 6. If the same subject is tested again a month from now, what is the probability of a positive test? (Compute the numerical answer)

Points: (a) 4, (b) 4, (c) 4, (d) 4, (e) 4, (f) 5.

3. Let X_1, \dots, X_n be a random sample from the following probability density function

$$\frac{3}{\theta} x^2 e^{-\frac{x^3}{\theta}}, \quad \theta > 0, x > 0.$$

In what follows, derive explicit expressions and simplify them as much as possible. Show *all* your derivations, not just the final answer.

- (a) Find a sufficient statistic for θ .
- (b) Is the statistic in (a) minimal? If it is, provide a proof. If it is not, find a minimal sufficient statistic.
- (c) Find the maximum likelihood estimator of θ .
- (d) Find the expected value of the estimator in (c).
- (e) Is the estimator in (c) UMVUE of θ ? Justify.
- (f) Find the Cramer-Rao lower bound on the variance of unbiased estimators of θ .
- (g) Derive the rejection region for the most powerful size α test of $H_0 : \theta = \theta_0$ against $H_a : \theta = \theta_1$, where $\infty > \theta_1 > \theta_0 > 0$. Is the given test uniformly most powerful of its size against the alternative $H_1 : \theta > \theta_0$? Justify.

Points: (a) 2 , (b) 3 , (c) 4 , (d) 4 , (e) 2 , (f) 5 , (g) 5.

4. Let X_1, \dots, X_n be a random sample from the following probability density function

$$f(x|\theta, \beta) = \frac{1}{\theta} e^{-\frac{(x-\beta)}{\theta}}, \quad x > \beta > 0, \theta > 0.$$

In what follows, derive explicit expressions and simplify them as much as possible. Show *all* your derivations, not just the final answer.

- (a) Find the maximum likelihood estimators for θ and β .
- (b) Derive the likelihood ratio test for testing $H_0 : \theta = \theta_0$ against $H_a : \theta > \theta_0$, with β unknown, and θ_0 a given positive constant. Identify a distribution (approximate or exact, whichever is easier) that may be used in constructing the rejection region for the test.
- (c) Derive the likelihood ratio test for testing $H_0 : \beta = \beta_0$ against $H_a : \beta > \beta_0$, with θ unknown, and β_0 a given positive constant. Identify a distribution (approximate or exact, whichever is easier) that may be used in constructing the rejection region for the test.

Points: (a) 7 , (b) 9 , (c) 9.