Table 10. Away from Home¹: Percentages² of Selected Nutrients Contributed by Food and Beverages Consumed Away from Home, by Race/Ethnicity and Age, in the United States, 2011-2012

Race/ethnicity and age	repor	_		ergy		otein	hyd	rbo- rate	sug	otal gars	fit	etary	fa		fa		unsat fa	ono- urated at	unsat f	oly- urated at
(years)	%	(SE)	%	(SE)	%	(SE)	<u></u>	(SE)	%	(SE)	%	(SE)	%	(SE)	%	(SE)	%	(SE)	%	(SE)
Non-Hispanic White:		I																		
2 - 5		(3.4)	34	(2.4)	34	(2.4)	34	(2.5)	34	(3.5)	36	(2.3)	33	(2.7)	32	(2.9)	33	(2.9)	35	(2.2)
6 - 11	83	(2.7)	38	(2.6)	38	(2.8)	37	(2.6)	39	(2.8)	37	(2.1)	38	(2.7)	38	(2.8)	38	(2.9)	38	(2.6)
12 - 19	77	(3.3)	41	(1.2)	42	(2.0)	41	(1.4)	41	(3.1)	38	(2.3)	42	(1.4)	41	(1.5)	42	(1.6)	43	(1.5)
20 and over	67	(1.8)	33	(1.1)	33	(1.4)	33	(1.0)	33	(1.2)	30	(1.2)	34	(1.2)	34	(1.3)	35	(1.2)	35	(1.3)
2 and over	69	(1.6)	35	(1.0)	34	(1.2)	34	(0.9)	34	(1.1)	31	(1.1)	35	(1.1)	35	(1.1)	36	(1.1)	36	(1.1)
Non-Hispanic Black:																				
2 - 5	53	(5.3)	25	(2.4)	23	(2.4)	26	(2.5)	27	(2.9)	25	(2.4)	25	(2.4)	25	(2.6)	25	(2.4)	25	(2.4)
6 - 11	79	(4.2)	42	(2.7)	40	(2.8)	44	(3.0)	47	(3.1)	44	(3.5)	42	(2.7)	43	(2.7)	41	(2.9)	40	(2.8)
12 - 19	60	(4.9)	30	(3.4)	28	(3.3)	30	(3.6)	30	(3.7)	28	(4.0)	30	(3.3)	29	(3.1)	29	(3.6)	31	(3.8)
20 and over	62	(2.9)	34	(3.2)	31	(3.1)	34	(3.3)	35	(3.5)	30	(2.7)	35	(3.2)	34	(3.3)	35	(3.2)	37	(3.4)
2 and over	62	(2.6)	34	(2.9)	31	(2.8)	34	(2.9)	35	(3.0)	31	(2.6)	35	(2.9)	34	(2.8)	35	(2.9)	36	(3.1)
Non-Hispanic Asian4:																				
2 - 5	62	(6.5)	26	(4.1)	24	(4.3)	26	(4.2)	26	(4.6)	25	(5.3)	26	(4.3)	27	(6.1)	25	(3.7)	28	(3.6)
6 - 11	64	(4.7)	25	(2.7)	23	(3.4)	25	(2.6)	26	(2.8)	23	(3.4)	24	(3.0)	22	(3.0)	23	(3.1)	27	(3.6)
12 - 19	67	(7.5)	32	(4.8)	30	(4.9)	31	(4.8)	34	(6.2)	29	(3.9)	35	(5.4)	34	(5.1)	36	(5.8)	36	(5.5)
20 and over	69	(2.2)	34	(1.4)	32	(1.5)	32	(1.4)	36	(2.1)	29	(2.0)	37	(1.8)	39	(2.1)	37	(2.0)	36	(1.6)
2 and over	68	(1.9)	33	(1.1)	31	(1.2)	31	(1.1)	34	(2.0)	28	(1.6)	36	(1.4)	36	(1.9)	35	(1.5)	36	(1.1)
Hispanic:																				
2 - 5	60	(3.1)	25	(2.1)	23	(2.0)	25	(2.1)	24	(1.9)	23	(2.1)	26	(2.3)	25	(2.2)	26	(2.5)	28	(2.7)
6 - 11	78	(4.1)	35	(2.3)	34	(2.5)	35	(2.2)	38	(2.7)	32	(1.9)	36	(2.6)	37	(2.7)	35	(2.6)	36	(2.8)
12 - 19	69	(3.2)	34	(2.4)	32	(2.4)	33	(2.4)	35	(2.7)	31	(2.4)	36	(2.6)	35	(2.8)	36	(2.5)	37	(2.8)
20 and over	66	(2.8)	35	(1.5)	35	(1.7)	34	(1.7)	36	(2.2)	30	(2.0)	37	(1.5)	36	(1.3)	37	(1.5)	38	(2.1)
2 and over	68	(2.1)	34	(1.1)	33	(1.2)	33	(1.3)	35	(1.7)	30	(1.5)	36	(1.1)	35	(1.0)	36	(1.1)	37	(1.6)

Table 10. Away from Home¹: Percentages² of Selected Nutrients Contributed by Food and Beverages Consumed Away from Home, by Race/Ethnicity and Age, in the United States, 2011-2012 *(continued)*

Race/ethnicity and age (years)	Choles- terol % (SE)	Vitamin A (RAE) % (SE)	Beta- carotene % (SE)	Lycopene % (SE)	Thiamin % (SE)	Ribo- flavin % (SE)	Niacin % (SE)	Vitamin B6 % (SE)	Folate (DFE) % (SE)
Non-Hispanic White:									
2 - 5	34 (5.4)	29 (3.5)	33*(12.7)	28 (6.4)	33 (1.7)	30 (2.3)	34 (2.1)	30 (2.2)	29 (1.8)
6 - 11	38 (3.4)	29 (2.3)	35 (4.8)	38 (4.1)	32 (2.1)	32 (2.5)	34 (1.6)	32 (2.2)	29 (2.6)
12 - 19	46 (4.4)	32 (3.5)	49 (5.9)	50 (6.1)	38 (1.9)	32 (1.7)	37 (2.2)	34 (2.0)	32 (2.3)
20 and over	34 (1.5)	25 (1.9)	30 (2.8)	29 (2.0)	31 (1.2)	28 (1.1)	34 (1.3)	32 (1.3)	28 (1.2)
2 and over	36 (1.6)	26 (1.8)	31 (2.6)	32 (1.7)	32 (1.0)	29 (1.0)	34 (1.1)	32 (1.2)	29 (1.0)
Non-Hispanic Black:									
2 - 5	20 (2.6)	26 (3.0)	35 (7.7)	24 (3.7)	24 (2.8)	23 (2.3)	23 (2.7)	21 (2.5)	24 (2.9)
6 - 11	38 (3.0)	43 (3.7)	40 (5.1)	45 (5.5)	39 (3.0)	43 (3.8)	36 (2.7)	36 (3.6)	37 (3.0)
12 - 19	27 (3.2)	28 (4.7)	35 (6.5)	31 (8.5)	28 (3.5)	27 (3.5)	28 (3.8)	27 (4.4)	26 (4.0)
20 and over	30 (3.0)	26 (2.7)	26 (3.2)	25 (5.0)	30 (3.1)	30 (3.1)	33 (3.4)	31 (3.4)	29 (2.9)
2 and over	30 (2.6)	28 (2.5)	27 (3.0)	28 (4.8)	30 (2.8)	30 (2.7)	32 (3.1)	30 (3.2)	29 (2.7)
Non-Hispanic Asian4:									
2 - 5	19* (4.7)	22 (5.0)	25* (8.7)	31*(14.0)	24 (4.1)	23 (4.3)	26 (3.2)	22 (3.3)	24 (4.9)
6 - 11	19 (2.4)	19 (3.8)	22 (5.7)	28 (7.3)	22 (3.3)	21 (4.1)	24 (3.4)	20 (3.8)	19 (3.1)
12 - 19	36 (7.6)	21 (5.7)	18 (3.1)	38 (11.0)	26 (4.3)	25 (5.0)	30 (4.9)	27 (5.1)	20 (4.2)
20 and over	33 (2.0)	28 (2.2)	26 (3.3)	34 (3.7)	30 (1.6)	32 (1.9)	32 (1.2)	32 (1.5)	29 (1.5)
2 and over	32 (1.7)	27 (1.4)	26 (2.7)	34 (2.7)	29 (1.2)	30 (1.8)	31 (1.1)	31 (1.2)	27 (1.0)
Hispanic:									
2 - 5	20 (2.5)	23 (2.5)	36 (6.7)	27 (3.4)	23 (1.7)	21 (1.5)	24 (1.7)	20 (1.5)	21 (2.2)
6 - 11	33 (3.5)	35 (2.9)	38 (7.4)	39 (3.4)	30 (2.3)	33 (2.6)	30 (2.2)	27 (2.3)	26 (2.1)
12 - 19	33 (3.5)	33 (5.3)	36 (6.9)	33 (5.0)	31 (2.7)	30 (2.9)	31 (2.2)	27 (2.5)	28 (2.6)
20 and over	35 (2.4)	28 (2.1)	35 (4.7)	32 (3.6)	31 (1.6)	31 (1.8)	35 (1.7)	34 (1.8)	28 (1.7)
2 and over	33 (1.9)	29 (1.7)	36 (3.8)	33 (2.3)	30 (1.1)	30 (1.2)	33 (1.1)	31 (1.3)	28 (1.2)

Table 10. Away from Home¹: Percentages² of Selected Nutrients Contributed by Food and Beverages Consumed Away from Home, by Race/Ethnicity and Age, in the United States, 2011-2012 *(continued)*

Race/ethnicity					Vitamin E (alpha-				
and age	Choline	Vitamin E	12 Vitamin C	Vitamin D	tocopherol)	Vitamin K	Calcium	Phosphorus	Magnesium
(years)	% (SI			% (SE)	% (SE)	% (SE)	% (SE)	% (SE)	% (SE)
Non-Hispanic White:									
2 - 5	33 (3.9) 27 (2.0	5) 26 (5.8)	27 (2.7)	32 (2.8)	34 (3.9)	30 (2.6)	32 (2.3)	32 (1.9)
6 - 11	36 (2.7	(3.1) 31 (3.1)	5) 35 (5.3)	33 (3.0)	34 (2.4)	38 (3.2)	37 (2.9)	38 (2.5)	35 (2.1)
12 - 19	42 (3.4	31 (2.6)) 48 (9.4)	29 (5.0)	40 (2.7)	43 (4.1)	35 (2.8)	39 (2.1)	36 (2.2)
20 and over	32 (1	3) 29 (2.6)) 27 (1.2)	20 (2.2)	32 (1.2)	30 (3.0)	28 (1.2)	31 (1.2)	29 (1.1)
2 and over	33 (1	3) 29 (1.	5) 29 (1.2)	23 (2.1)	33 (1.1)	32 (2.7)	29 (1.1)	33 (1.1)	30 (1.0)
Non-Hispanic Black:									
2 - 5	22 (1.9) 25 (2.9	9) 22 (3.9)	20 (2.2)	27 (2.7)	31 (7.2)	23 (2.6)	24 (2.5)	23 (2.3)
6 - 11	41 (3.3	3) 40 (3.	42 (4.1)	47 (4.3)	39 (3.1)	43 (4.8)	48 (3.2)	45 (3.1)	41 (3.1)
12 - 19	28 (3.0	6) 29 (6.6)	30 (4.4)	24 (5.5)	29 (4.7)	27 (4.3)	28 (3.5)	29 (3.2)	27 (3.1)
20 and over	31 (3.0	30 (3.	7) 27 (2.9)	22 (2.7)	36 (3.1)	29 (3.7)	29 (2.9)	32 (3.1)	31 (2.6)
2 and over	31 (2.8	31 (3.	5) 28 (2.5)	25 (2.5)	35 (3.0)	30 (3.5)	31 (2.4)	33 (2.7)	31 (2.4)
Non-Hispanic Asian4:									
2 - 5	21 (4.0)) 21 (4	5) 30* (9.5)	20* (5.0)	26 (4.9)	37*(13.2)	23 (5.6)	24 (4.2)	25 (4.1)
6 - 11	20 (2.7	20 (4.9)	9) 14* (3.5)	17* (5.4)	23 (3.5)	15* (4.1)	20 (4.3)	22 (3.4)	22 (3.3)
12 - 19	31 (6.4) 27 (7.	22 (4.6)	21 (6.0)	33 (5.4)	21 (4.3)	25 (4.7)	29 (4.8)	28 (3.9)
20 and over	31 (1.7	31 (3.6)) 26 (2.3)	24 (2.9)	33 (2.1)	28 (3.6)	31 (1.6)	32 (1.6)	30 (1.7)
2 and over	30 (1	30 (2.3	7) 25 (1.8)	23 (2.6)	32 (1.6)	27 (2.8)	29 (1.3)	30 (1.2)	29 (1.3)
Hispanic:									
2 - 5	21 (2.2	2) 19 (1.6	5) 19 (1.3)	19 (1.5)	27 (1.9)	25 (2.7)	21 (2.2)	23 (2.0)	22 (1.8)
6 - 11	33 (2.9	33 (3.6	5) 32 (3.9)	34 (3.4)	35 (2.2)	37 (3.2)	37 (2.7)	35 (2.4)	33 (2.2)
12 - 19	32 (2.8			26 (4.5)	34 (2.2)	35 (3.6)	34 (3.4)	33 (2.9)	31 (2.3)
20 and over	33 (1.0	5) 32 (1.5)	32 (3.8)	24 (2.0)	36 (2.1)	42 (5.0)	30 (1.4)	32 (1.4)	31 (1.7)
2 and over	32 (1	31 (1.3	30 (2.8)	25 (1.5)	35 (1.5)	40 (4.1)	31 (0.9)	32 (1.0)	31 (1.2)

Table 10. Away from Home¹: Percentages² of Selected Nutrients Contributed by Food and Beverages Consumed Away from Home, by Race/Ethnicity and Age, in the United States, 2011-2012 *(continued)*

Race/ethnicity and age	Iro	on (SE)	Zi %	nc (SE)	Co _l	oper	Sele %	nium (SE)	Pota:	ssium (SE)	Soc %	lium (SE)	Caf	feine (SE)	Alc %	ohol (SE)
(years)	%	(SE)	70	(SE)	70	(SE)	70	(SE)	70	(SE)	%0	(SE)	90	(SE)	70	(SE)
Non-Hispanic White:																
2 - 5	30	(2.4)	31	(2.9)	34	(2.4)	34	(2.0)	33	(2.6)	33	(1.9)	41	(6.7)		
6 - 11	31	(2.3)	34	(2.7)	36	(2.2)	37	(2.5)	37	(2.4)	40	(2.2)	34	(7.8)		
12 - 19	35	(2.2)	37	(2.5)	40	(2.2)	43	(2.3)	39	(2.2)	41	(1.7)	37	(3.9)		
20 and over	29	(1.0)	31	(1.3)	29	(1.5)	32	(1.4)	30	(1.1)	34	(1.2)	30	(1.5)	35	(5.0)
2 and over	30	(1.0)	32	(1.2)	31	(1.4)	34	(1.3)	31	(1.1)	35	(1.0)	30	(1.5)		
Non-Hispanic Black:																
2 - 5	23	(2.8)	23	(2.8)	27	(3.4)	23	(2.4)	24	(2.4)	25	(2.8)	24	(4.3)		
6 - 11	38	(2.8)	40	(2.9)	43	(3.1)	38	(2.7)	44	(3.6)	39	(2.8)	28	(3.6)		
12 - 19	25	(3.7)	26	(3.4)	29	(3.5)	28	(3.3)	30	(3.7)	31	(3.9)	47	(8.2)		
20 and over	29	(2.8)	31	(2.8)	32	(2.3)	31	(2.8)	31	(3.1)	33	(3.4)	44	(3.4)	35	(5.8)
2 and over	29	(2.6)	31	(2.6)	32	(2.2)	31	(2.6)	32	(2.9)	33	(3.1)	43	(3.0)		
Non-Hispanic Asian4:																
2 - 5	25	(3.8)	22	(3.5)	26	(4.5)	26	(5.1)	23	(4.1)	25	(3.7)	12*	(7.5)		
6 - 11	22	(3.5)	20	(3.0)	23	(2.8)	22	(3.0)	22	(3.2)	24	(2.5)	40*	(12.6)		
12 - 19	26	(4.6)	29	(5.4)	30	(3.9)	29	(5.0)	28	(4.5)	30	(4.8)	50	(9.4)		
20 and over	30	(1.5)	32	(1.9)	29	(1.5)	31	(1.6)	31	(1.7)	32	(1.9)	35	(4.0)	32	(6.6)
2 and over	29	(1.3)	30	(1.3)	29	(1.2)	30	(1.3)	29	(1.3)	31	(1.3)	36	(3.7)		
Hispanic:																
2 - 5	22	(2.0)	21	(2.0)	23	(1.8)	23	(2.4)	23	(2.0)	24	(1.9)	28	(4.5)		
6 - 11	30	(1.8)	32	(2.6)	36	(2.5)	32	(2.6)	34	(2.5)	33	(2.2)	34	(5.2)		
12 - 19	28	(2.1)	31	(2.1)	36	(4.4)	32	(2.4)	31	(2.5)	35	(2.6)	42	(3.9)		
20 and over	30	(1.4)	33	(1.5)	32	(1.6)	34	(1.6)	32	(1.6)	36	(1.3)	35	(2.5)	35	(4.4)
2 and over	29	(1.0)	32	(1.1)	33	(1.2)	33	(1.2)	32	(1.3)	35	(1.0)	35	(2.4)		

Symbol Legend

* Indicates an estimate that may be less statistically reliable than estimates that are not flagged. The rules for flagging estimated percentages and ratios are as follows:

Percent reporting: An estimated percentage between 25 and 75 percent is flagged when based on a sample size of less than 30 times the variance inflation factor (VIF), where the VIF represents a broadly calculated average design effect, or when the relative standard error is greater than 30 percent. An estimated percentage less than or equal to 25 percent or greater than or equal to 75 percent is flagged when the smaller of np and n(1-p) is less than 8 times the VIF, where n is the sample size and p is the percentage expressed as a fraction. The VIF used in this table is 2.50.

Nutrient ratios expressed as percentages: An estimated ratio between 25 and 75 percent is flagged when based on a sample size n^* of less than 30 times the variance inflation factor (VIF), where the VIF represents a broadly calculated average design effect and n^* is the number of individuals in the sample reporting non-zero intake of the respective nutrient. An estimated ratio less than or equal to 25 percent or greater than or equal to 75 percent, is flagged when the smaller of n^*p and n^* (1-p) is less than 8 times the VIF, where p is the percentage expressed as a fraction. Additionally, an estimated ratio is flagged when either the relative standard error or p/(1-p) times the relative standard error is greater than 30 percent. The VIF used in this table is 2.50.

Footnotes

- Away from home includes any location other than home. Home is defined as an individual's dwelling unit and the surrounding areas that are used solely by the occupants of that dwelling unit.
- ² Percentages are estimated as a ratio of total nutrients from food and beverages consumed away from home for all individuals to total daily nutrient intakes for all individuals. Total daily nutrient intakes are available from: www.ars.usda.gov/nea/bhnrc/fsrg. See Table 2. Nutrient Intakes from Food and Beverages: Mean Amounts Consumed per Individual, by Race/Ethnicity and Age, in the United States, 2011-2012.
- ³ The percentage of respondents in the race/ethnicity/age group who reported consuming at least one item away from home.
- ⁴ A new sampling methodology was implemented for NHANES 2011-2012 to oversample non-Hispanic non-black Asians in addition to the ongoing oversample of Hispanics, non-Hispanic blacks, low-income whites/others, and whites/others 80+ years.

Abbreviations

SE = standard error; RAE = retinol activity equivalents; DFE = dietary folate equivalents.

Notes Applicable to All Tables in Series: What We Eat in America, NHANES 2011-2012

Sample weights designed for dietary analysis were used to allow estimates representative of the U. S. population for the years of collection.

The statistics in this table are estimated from Day 1 dietary recall interviews conducted in the *What We Eat in America*, National Health and Nutrition Examination Survey (NHANES) 2011-2012. The 24-hour dietary recalls were conducted in-person, by trained interviewers, using the USDA 5-step Automated Multiple-Pass Method. Food intakes were coded and nutrient values were determined using the USDA Food and Nutrient Database for Dietary Studies 2011-2012 www.ars.usda.gov/nea/bhnrc/fsrg which is based on nutrient values in the USDA National Nutrient Database for Standard Reference, Release 26 (Agricultural Research Service, Nutrient Data Laboratory, 2013).

Intakes of nutrients and other dietary components are based on the consumption of food and beverages, including water, and do not include intake from supplements or medications.

The table includes data from individuals 2 years and over. Breast-fed children were excluded because breast milk was not quantified in dietary recall interviews.

Although alcohol data are collected for all individuals, estimates are not presented for age groups under 20 years due to extreme variability and/or inadequate sample size.

Suggested Citation

U.S. Department of Agriculture, Agricultural Research Service. 2014. Away from Home: Percentages of Selected Nutrients Contributed by Food and Beverages Consumed Away from Home, by Race/Ethnicity and Age, What We Eat in America, NHANES 2011-2012.