

Evolution of resilience in protein interactomes across the tree of life

Marinka Zitnik^a, Rok Sosič^a, Marcus W. Feldman^{b,1}, and Jure Leskovec^{a,c,1}

^aDepartment of Computer Science, Stanford University, Stanford, CA 94305; ^bDepartment of Biology, Stanford University, Stanford, CA 94305; and ^cChan Zuckerberg Biohub, San Francisco, CA 94158

Contributed by Marcus W. Feldman, December 18, 2018 (sent for review October 19, 2018; reviewed by Edoardo Airoldi and Aviv Bergman)

Phenotype robustness to environmental fluctuations is a common biological phenomenon. Although most phenotypes involve multiple proteins that interact with each other, the basic principles of how such interactome networks respond to environmental unpredictability and change during evolution are largely unknown. Here we study interactomes of 1,840 species across the tree of life involving a total of 8,762,166 protein–protein interactions. Our study focuses on the resilience of interactomes to network failures and finds that interactomes become more resilient during evolution, meaning that interactomes become more robust to network failures over time. In bacteria, we find that a more resilient interactome is in turn associated with the greater ability of the organism to survive in a more complex, variable, and competitive environment. We find that at the protein family level proteins exhibit a coordinated rewiring of interactions over time and that a resilient interactome arises through gradual change of the network topology. Our findings have implications for understanding molecular network structure in the context of both evolution and environment.

protein–protein interaction networks | molecular evolution | ecology | network resilience | network rewiring

The enormous diversity of life shows a fundamental ability of organisms to adapt their phenotypes to changing environments (1). Most phenotypes are the result of an interplay of many molecular components that interact with each other and the environment (2–5). The study of life's diversity has a long history and extensive phylogenetic studies have demonstrated evolution at the DNA sequence level (6–8). While studies based on sequence data alone have demonstrated evolution of genomes, mechanistic insights into how evolution shapes interactions between proteins in an organism remain elusive (9, 10).

DNA sequence information has been used to associate genes with their functions (11), determine properties of ancestral life (12, 13), and understand how the environment affects genomes (14). Despite these advances in understanding DNA sequence evolution, little is known about basic principles that govern the evolution of interactions between proteins. In particular, evolution of DNA and amino acid sequences could lead to pervasive rewiring of protein–protein interactions and create or destroy the ability of the interactions to perform their biological functions.

The importance of protein–protein interactions has spurred experimental efforts to map all interactions between proteins in a particular organism, its interactome, namely the complex network of protein–protein interactions in that organism. A large number of high-throughput experiments have reported high-quality interactomes in a number of organisms (15–19). Because interactomes underlie all living organisms, it is critical to understand how these networks change during evolution (20, 21) and elucidate key principles of their structure.

Here, we use protein interactions measured by these large-scale interactome mapping experiments and study the evolutionary dynamics of the interactomes across the tree of life. Our protein interaction dataset contains a total of 8,762,166

physical interactions between 1,450,633 proteins from 1,840 species, encompassing all current protein interaction information at a cross-species scale (*SI Appendix, section S1 and Table S4*). We group these interactions by species and represent each species with a separate interactome network, in which nodes indicate a species' proteins and edges indicate experimentally documented physical interactions, including direct biophysical protein–protein interactions, regulatory protein–DNA interactions, metabolic pathway interactions, and kinase–substrate interactions measured in that species. We integrate into the dataset (22) the evolutionary history of species provided by the tree of life constructed from small subunit ribosomal RNA gene sequence information (12) (*SI Appendix, section S2*). Using network science, we study the network organization of each interactome, in particular its resilience to network failures, a critical factor determining the function of the interactome (23–26). We identify the relationship between the resilience of an interactome and evolution and use this resilience to uncover relationships with natural environments in which organisms live. Although the interactomes are incomplete and biased toward much-studied proteins and model species (*SI Appendix, section S1 and Fig. S7*), our analyses give results that are consistent across taxonomic groups, that are not sensitive to network data quality or network size change (*SI Appendix, section S8 and Fig. S8*), and indicate that our conclusions will still hold when more protein interaction data become available.

Significance

The interactome network of protein–protein interactions captures the structure of molecular machinery that underlies organismal complexity. The resilience to network failures is a critical property of the interactome as the breakdown of interactions may lead to cell death or disease. By studying interactomes from 1,840 species across the tree of life, we find that evolution leads to more resilient interactomes, providing evidence for a longstanding hypothesis that interactomes evolve favoring robustness against network failures. We find that a highly resilient interactome has a beneficial impact on the organism's survival in complex, variable, and competitive habitats. Our findings reveal how interactomes change through evolution and how these changes affect their response to environmental unpredictability.

Author contributions: M.Z., M.W.F., and J.L. designed research; M.Z., R.S., and J.L. performed research; M.Z., R.S., M.W.F., and J.L. analyzed data; and M.Z., R.S., M.W.F., and J.L. wrote the paper.

Reviewers: E.A., Harvard University; and A.B., Albert Einstein College Of Medicine.

The authors declare no conflict of interest.

This open access article is distributed under [Creative Commons Attribution-NonCommercial-NoDerivatives License 4.0 \(CC BY-NC-ND\)](https://creativecommons.org/licenses/by-nc-nd/4.0/).

¹To whom correspondence may be addressed. Email: mfeldman@stanford.edu or jure@cs.stanford.edu.

This article contains supporting information online at www.pnas.org/lookup/suppl/doi:10.1073/pnas.1818013116/-DCSupplemental.

Published online February 14, 2019.





Fig. 1. Protein interaction data of 1,840 species consisting of 8,762,166 interactions by 1,450,633 proteins reveal the resilience of interactomes across vast evolutionary distances. (A) The interactome of an organism consists of all physical interactions between proteins in the organism. When interactions involving a certain fraction ($f = 5/45$ in this example) of the proteins are removed from the interactome, the interactome fragments into a number of isolated network components. (B) Modified Shannon diversity H_{msh} (SI Appendix, section S5) measures how the interactome fragments into isolated components at a given network failure rate f . (C) The resilience of the interactome integrates modified Shannon diversity H_{msh} across all possible failure rates f (SI Appendix, section S5). Resilience value 1 indicates the most resilient interactome, and resilience value 0 indicates a complete loss of the connectivity of the interactome (SI Appendix, Fig. S3). *Homo sapiens* (*H. influenzae*) has the most (least) resilient interactome (their resilience is 0.461 and 0.267, respectively) among the three selected organisms. (D) A small neighborhood of the interactome in a eukaryotic and a bacterial species. As ancestral species have gone extinct, older interactomes have been lost, and only interactomes of present-day species are available to us. (E) Phylogenetic tree showing 1,539 bacteria, 111 archaea, and 190 eukarya (12). Evolution of a species is represented as the total branch length (nucleotide substitutions per site) from the root to the corresponding leaf in the tree (SI Appendix, section S2). The outside circle of bars shows the interactome resilience of every species. Current protein-protein interaction data might be prone to notable selection and investigative biases (SI Appendix, section S1). (F) This plot shows the interactome resilience for 171 species with at least 1,000 publications in the NCBI PubMed (SI Appendix, Fig. S7). Across all species, evolution of a species predicts resilience of the species' interactome to network failures (LOWESS fit; $R^2 = 0.36$); more genetic change implies a more resilient interactome. Three species with the most nucleotide substitutions per site (far right on the x axis) have on average a 20.4% more resilient interactome than the three species with the least substitutions (far left on the x axis).

Relationship Between Interactome Resilience and Ecology. We next ask whether there is a relationship between species' interactome resilience and aspects of species' ecology (*SI Appendix, section S4*). We examine the relationship between interactome resilience and the fraction of regulatory genes and find that bacteria with more resilient interactomes have significantly more regulatory genes in their genomes ($R^2 = 0.32$; Fig. 24). Bacteria with highly resilient interactomes can also survive in more diverse and competitive environments, as revealed by



Zitnik et al.




Fig. 3. Evolution mitigates local network structural changes in protein interactomes. (A) A hypothetical phylogenetic tree illustrates a speciation event that gives rise to two lineages according to the speciation-divergence model (11) and leads to present-day organisms "1" (green) and "2" (pink). In this example, a single ancestral protein A that was present in the ancestral species gives rise to proteins A' and A'' upon speciation; A' and A'' form an orthologous protein pair. As the two newly arising species diverge and protein sequences evolve, protein network neighborhoods (SI Appendix, Fig. S4) in their interactomes can rewire independently over time. Shown are also in-paralogs, proteins which arise through gene duplication events in species 1 and 2 after speciation. (B) A hypothetical protein family with three protein members (A', A'', A'''), each from a different organism. In the phylogenetic tree, organism 1 is located at the tip of the lineage with the shortest branch length, whereas organism 3 is in the lineage with the longest branch length in the tree. We represent the protein family by a sequence of orthologous proteins ordered by the branch length of proteins' originating species (SI Appendix, section S3). We then characterize the network neighborhood of each protein in the family by calculating two network metrics (SI Appendix, Fig. S5). Isolated components are given by the degree-adjusted number of connected components in the neighborhood that arise when the central protein is removed from the interactome (gray) (SI Appendix, section S6). The neighborhood size down-weighted by the redundancy of local interactions gives the effective size of the neighborhood (SI Appendix, section S6). (C and D) The number of isolated network components and the effective size of protein neighborhoods both decrease with evolution ($P = 3 \cdot 10^{-8}$ and $P = 0.03$, respectively; Spearman's ρ rank correlation), suggesting that local interaction neighborhoods rewire via a coordinated evolutionary mechanism. Lines in C and D show the LOWESS fit of median-aggregated network metric values for 81,673 proteins from 2,224 protein families; color bands indicate 95% confidence band for the LOWESS fit; gray lines show random expectation.

evolution. These structural changes in the neighborhoods suggest a molecular network model of evolution (Fig. 3B): For orthologous proteins in two species, as the evolutionary distance between the species increases, the proteins' local network neighborhoods become increasingly different and the neighborhood becomes more interconnected in the species that has undergone more genetic change.

Network Rewiring of Protein–Protein Interactions. To study evolutionary mechanisms of structural changes in the interactomes, we investigate network motifs (34, 35). We first identify orthologous protein pairs from evolutionarily close species (SI Appendix, section S3), resulting in 2,485,564 protein pairs, which we then use to calculate interaction rewiring rates (IRRs) for selected network motifs (Fig. 4A). We calculate the number of times each motif appears in each protein neighborhood and derive the IRR by comparing the motif occurrences between the interactomes of the older and the younger species of each protein pair (SI Appendix, section S7.1). We find strong statistical evidence that network motifs rewire during evolution ($P < 10^{-33}$ for all network motifs; Fig. 4B), suggesting that rewiring of interactions is an important mechanism for the evolution of interactomes. For example, proteins in evolutionarily older species on average participate in a factor of 0.861 fewer protein–protein interactions compared with proteins in evolutionarily younger species (IRR = -0.215 ; Fig. 4B). This significant negative correlation between a protein's number of interactions and the protein's evolutionary age confirms earlier studies of *Saccharomyces cerevisiae* (36). We also find that square motifs of interactions become more common in protein neighborhoods during evolution (IRR = 0.016 ; Fig. 4B). A range of biological evidence (18, 37, 38) supports this positive rate of change in the number of square motifs: From a structural perspective (38), protein–protein interactions often require complementary interfaces; hence two proteins with similar interfaces share many of their neighbors. However, they might not interact directly with each other, which manifests in the interactome as a square motif of interactions (see SI Appendix, Fig. S6 for an illustration of interaction interfaces recognizing the binding sites in proteins). Evolutionary arguments following gene duplication (18) reach the same conclusion; proteins with multiple shared interaction partners are likely to share even more partners and thereby produce new square motifs of interactions. To test the predictive power of our motif-based model of structural network changes, we estimate the size of the whole human interactome by extrapolating the *S. cerevisiae* interactome, using IRRs from Fig. 4B (SI Appendix, section S7.3). Assuming one splice isoform per gene, we predict the number of interactions in humans to be $\sim 160,000$. This prediction is in surprisingly good agreement with three previous estimates of the size of the human interactome, which range from 150,000 to 370,000 interactions (15, 17, 39) and have proved crucial in establishing the complexity of the human interactome (19).

Discussion

Our analyses reveal how protein–protein interaction networks change through evolution and how changes in these networks affect phenotypes and organismal response to environmental complexity. This systematic investigation of protein–protein interaction networks from an evolutionary perspective was enabled by a dataset of interactomes, consisting of protein–protein interaction networks from 1,840 species. To date, most evolutionary analyses of biological networks have focused on a small number of organisms with high-coverage protein–protein

interaction data, such as *S. cerevisiae*, *Mus musculus*, and humans. This is because interactomes mapped by unbiased tests of all possible pairwise combinations of proteins on the same platform remain scarce, an important limitation of the present study. Furthermore, experimentally documented protein interactions are currently subject to a high number of false positives and negatives. As more protein interaction data are collected, and more genomes become available, the generalizability of our findings can be further evaluated. However, our results are consistent across both different subsets of protein interaction data (SI Appendix, Table S2) and different phylogenetic lineages (SI Appendix, Fig. S10) and are not explained by many possible genomic and network confounders (SI Appendix, section S8, Fig. S8, and Table S1), thus providing confidence that our key findings cannot be attributed to biases in the datasets.

Interactome resilience is an important aspect of our study. The resilience measures fragmentation of the interactome into isolated components and thus represents a global measure of the interactome's topological stability. Beyond fragmentation, there are other possible modifications of the interactome that could alter the network's biological function without necessarily disconnecting the network (40–42). As more detailed information about functions of individual proteins in the interactome (43), as well as dynamic protein-expression data (44), becomes available, our measure of interactome resilience could be adapted to give a more complex definition of resiliency, which might yield more detailed evolutionary predictions. Additionally, information on how protein–protein interactions change dynamically both in time and space (45–47) might reveal how topological stability of the interactome depends on large-scale interactome connectivity as well as on the interactome's dynamic properties (40).

Our study presents an additional paradigm for evolutionary studies by demonstrating that interactomes reveal fundamental structural principles of molecular networks. Our findings highlight evolution as an important predictor of structural network change and show that evolution of a species predicts resilience of the species' interactome to protein failures. The findings offer quantitative evidence for the biological proposition that an organism that has undergone more genetic change has a more resilient interactome, which, in turn, is associated with the greater ability of the organism to survive in a more complex, variable, or competitive environment. Our findings can also help clarify the mechanisms of how interactomes change during evolution, why currently observed network structures exist, and how they may change in the future and facilitate the extrapolation of functional information from experimentally characterized proteins to their orthologous proteins in poorly studied organisms.

Materials and Methods

Detailed description of data, statistical methodology, and additional analyses are provided in SI Appendix.

Code and Data Availability. Software implementation of statistical methodology is publicly available at snap.stanford.edu/tree-of-life. All data used in this paper, including the processed interactomes, are shared with the community and available from snap.stanford.edu/tree-of-life.

ACKNOWLEDGMENTS. For helpful discussion and comments on the manuscript, we thank Aaron Goodman, Hunter B. Fraser, Alice Y. Ting, Emma Pierson, Michael An, and Pratyaksh Sharma. M.Z., R.S., and J.L. were supported in part by NSF, NIH, Defense Advanced Research Projects Agency, Boeing, Stanford Data Science Initiative, and Chan Zuckerberg Biohub. M.W.F. was supported in part by the Stanford Center for Computational, Evolutionary, and Human Genomics and a grant from the John Templeton Foundation.

1. Castelle CJ, Banfield JF (2018) Major new microbial groups expand diversity and alter our understanding of the tree of life. *Cell* 172:1181–1197.
2. Barabási AL, Oltvai ZN (2004) Network biology: Understanding the cell's functional organization. *Nat Rev Genet* 5:101–113.

3. Hu JX, Thomas CE, Brunak S (2016) Network biology concepts in complex disease comorbidities. *Nat Rev Genet* 17:615–629.
4. Huttlin EL, et al. (2017) Architecture of the human interactome defines protein communities and disease networks. *Nature* 545:505–509.

5. Chen S, et al. (2018) An interactome perturbation framework prioritizes damaging missense mutations for developmental disorders. *Nat Genet* 50:1032–1040.
6. Britten RJ (1986) Rates of DNA sequence evolution differ between taxonomic groups. *Science* 231:1393–1398.
7. Yang Z, Rannala B (2012) Molecular phylogenetics: Principles and practice. *Nat Rev Genet* 13:303–314.
8. Marsit S, et al. (2017) Evolutionary biology through the lens of budding yeast comparative genomics. *Nat Rev Genet* 18:581–598.
9. Studer RA, et al. (2016) Evolution of protein phosphorylation across 18 fungal species. *Science* 354:229–232.
10. Sorrells TR, Booth LN, Tuch BB, Johnson AD (2015) Intersecting transcription networks constrain gene regulatory evolution. *Nature* 523:361–365.
11. Tatusov RL, Koonin EV, Lipman DJ (1997) A genomic perspective on protein families. *Science* 278:631–637.
12. Hug LA, et al. (2016) A new view of the tree of life. *Nat Microbiol* 1:16048.
13. Parks DH, et al. (2017) Recovery of nearly 8,000 metagenome-assembled genomes substantially expands the tree of life. *Nat Microbiol* 2:1533–1542.
14. Mukherjee S, et al. (2017) 1,003 reference genomes of bacterial and archaeal isolates expand coverage of the tree of life. *Nat Biotechnol* 35:676–683.
15. Rual JF, et al. (2005) Towards a proteome-scale map of the human protein–protein interaction network. *Nature* 437:1173–1178.
16. Roguev A, et al. (2008) Conservation and rewiring of functional modules revealed by an epistasis map in fission yeast. *Science* 322:405–410.
17. Venkatesan K, et al. (2009) An empirical framework for binary interactome mapping. *Nat Methods* 6:83–90.
18. Arabidopsis Interactome Mapping Consortium, et al. (2011) Evidence for network evolution in an Arabidopsis interactome map. *Science* 333:601–607.
19. Rolland T, et al. (2014) A proteome-scale map of the human interactome network. *Cell* 159:1212–1226.
20. Leskovec J, Kleinberg J, Faloutsos C (2005) Graphs over time: Densification laws, shrinking diameters and possible explanations. *Proceedings of the Eleventh ACM SIGKDD International Conference on Knowledge Discovery in Data Mining*, ed Grossman RL (ACM, New York), pp 177–187.
21. Leskovec J, Kleinberg J, Faloutsos C (2007) Graph evolution: Densification and shrinking diameters. *ACM Trans Knowl Discovery Data* 1:1–41.
22. Zitnik M, et al. (2019) Machine learning for integrating data in biology and medicine: Principles, practice, and opportunities. *Inf Fusion* 50:71–91.
23. Albert R, Jeong H, Barabási AL (2000) Error and attack tolerance of complex networks. *Nature* 406:378–382.
24. Wagner A (2003) Does selection mold molecular networks? *Sci Signaling* 2003:pe41.
25. Wagner A (2005) Robustness, evolvability, and neutrality. *FEBS Lett* 579:1772–1778.
26. Wagner A (2013) *Robustness and Evolvability in Living Systems* (Princeton Univ Press, Princeton).
27. Vo TV, et al. (2016) A proteome-wide fission yeast interactome reveals network evolution principles from yeasts to human. *Cell* 164:310–323.
28. Sheldon AL (1969) Equitability indices: Dependence on the species count. *Ecology* 50:466–467.
29. MaGuarran A (1988) *Ecological Diversity and Its Measurement* (Princeton Univ Press, Princeton).
30. Goodwin SB, Spielman L, Matuszak J, Bergeron S, Fry W (1992) Clonal diversity and genetic differentiation of *Phytophthora infestans* populations in northern and central Mexico. *Phytopathology* 82:955–961.
31. Baczowski A, Joanes D, Shamia G (1997) Properties of a generalized diversity index. *J Theor Biol* 188:207–213.
32. Freilich S, et al. (2009) Metabolic-network-driven analysis of bacterial ecological strategies. *Genome Biol* 10:R61.
33. Barve A, Wagner A (2013) A latent capacity for evolutionary innovation through exaptation in metabolic systems. *Nature* 500:203–206.
34. Benson AR, Gleich DF, Leskovec J (2016) Higher-order organization of complex networks. *Science* 353:163–166.
35. Yin H, Benson AR, Leskovec J (2018) Higher-order clustering in networks. *Phys Rev E* 97:052306.
36. Fraser HB, Hirsh AE, Steinmetz LM, Scharfe C, Feldman MW (2002) Evolutionary rate in the protein interaction network. *Science* 296:750–752.
37. Ispolatov I, Yuryev A, Mazo I, Maslov S (2005) Binding properties and evolution of homodimers in protein–protein interaction networks. *Nucleic Acids Res* 33:3629–3635.
38. Keskin O, Tuncbag N, Gursoy A (2016) Predicting protein–protein interactions from the molecular to the proteome level. *Chem Rev* 116:4884–4909.
39. Hart GT, Ramani AK, Marcotte EM (2006) How complete are current yeast and human protein–interaction networks? *Genome Biol* 7:120.
40. Siegal ML, Promislow DE, Bergman A (2007) Functional and evolutionary inference in gene networks: Does topology matter? *Genetica* 129:83–103.
41. Leclerc RD (2008) Survival of the sparsest: Robust gene networks are parsimonious. *Mol Syst Biol* 4:213.
42. Masel J, Siegal ML (2009) Robustness: Mechanisms and consequences. *Trends Genet* 25:395–403.
43. Agrawal M, Zitnik M, Leskovec J (2018) Large-scale analysis of disease pathways in the human interactome. *Pacific Symposium on Biocomputing*, eds Altman RB, Dunker AK, Hunter L, Ritchie MD, Klein TE (World Scientific, Singapore), Vol 23, pp 111–122.
44. Deng Q, Ramsköld D, Reinisius B, Sandberg R (2014) Single-cell RNA-seq reveals dynamic, random monoallelic gene expression in mammalian cells. *Science* 343:193–196.
45. Han JDJ, et al. (2004) Evidence for dynamically organized modularity in the yeast protein–protein interaction network. *Nature* 430:88–93.
46. Smith C, Puzio RS, Bergman A (2014) Hierarchical network structure promotes dynamical robustness. arXiv:1412.0709. Preprint, posted December 1, 2014.
47. Smith C, Pechuan X, Puzio RS, Biro D, Bergman A (2015) Potential unsatisfiability of cyclic constraints on stochastic biological networks biases selection towards hierarchical architectures. *J R Soc Interface* 12:20150179.