

Measured polarization — 241 samples / ~60 s per condition

Arithmetic mean +/- sample SD; no fits. A+B externally driven. Values are in the recorded PAX frame.

Quantity	A only	B only	A+B (moving)
Samples	241	241	241
Span (s)	60.000389	60.000372	60.000504
theta / ellipse azimuth (deg)	1.667607 +/- 0.003453	-87.304329 +/- 0.002348	0.924560 +/- 39.573563
eta / signed ellipticity (deg)	2.135098 +/- 0.001049	1.870229 +/- 0.001680	-5.221094 +/- 27.682354
absolute ellipticity (deg)	2.135098 +/- 0.001049	1.870229 +/- 0.001680	24.543820 +/- 13.739615
minor/major ellipse axis ratio	0.037282 +/- 0.000018	0.032653 +/- 0.000029	0.486286 +/- 0.289690
S1 orientation	0.995535 +/- 0.000007	-0.993455 +/- 0.000011	0.078475 +/- 0.060394
S2 orientation	0.058016 +/- 0.000120	-0.093758 +/- 0.000081	-0.095625 +/- 0.640027
S3 orientation	0.074460 +/- 0.000037	0.065237 +/- 0.000059	-0.158644 +/- 0.741785
Raw DOP	1.016701 +/- 0.000093	1.000054 +/- 0.000025	0.849456 +/- 0.032666
Length of mean orientation vector	0.999999992	0.999999995	0.201172379

All values are mean +/- sample SD (N-1), not uncertainty of the mean or absolute accuracy.

External phi1 drive during A+B is operator-supplied context; no drive waveform, phase, or frequency is inferred.

A/B are H-like/V-like in the recorded PAX frame (+S1/-S1); no axis rotation or model transfer is applied.

A+B averages describe time occupancy of a moving state; signed ellipticity can cancel. No single ellipse represents it.

CSV Stokes are reconstructed unit directions. Their mean length is not the PAX DOP or an intensity-weighted DOP.

Raw DOP >1: A 241/241, B 239/241. Retained unchanged; absolute accuracy is not established.


Run metadata says PAX at C; catalog setup says final output. Location was not independently verified in this report.

No fitting, row rejection, smoothing, detrending, drive reconstruction, DOP clipping, or model updates.




Full precision, axis distances, block means and formulas: [summary.md](#) / [statistics.csv](#) / [axis-angles.csv](#) / [block-means.csv](#)

Static paths: measured orientation and ellipticity





A only (H-like)



B only (V-like)



A+B: sampled movement near the $S_1=0$ circle (external ϕ_1 drive)



Static mean-state axis distances

Poincare degrees; unit direction of the averaged Stokes vector. H/V reference = +S1/-S1.

Signed pole	A only	B only
+S1	5.416420	173.441322
-S1	174.583580	6.558678
+S2	86.674055	95.379823
-S2	93.325945	84.620177
+S3	85.729805	86.259542
-S3	94.270195	93.740458

A-B separation: 171.730100 deg; departure from antipodal: 8.269900 deg.

A+B latitude from S1=0: 4.509571 +/- 3.477757 deg (sample SD).

Sampled around-S1 covering arc: 302.969 deg; no rotation count inferred.

Mean A+B direction is intentionally not presented as a static state; see its trajectory and per-sample axis statistics.