

Efficient Implementation of Interpolation Algorithms for EUF and Octagonal Theories

by

José Abel Castellanos Joo

B.Tech., Universidad de las Américas Puebla, 2015

THESIS

Submitted in Partial Fulfillment of the
Requirements for the Degree of

Master of Science
Computer Science

The University of New Mexico

Albuquerque, New Mexico

January, 2019

Dedication

To my family and friends, and love.

“Lol” – Anonymous

Acknowledgments

I would like to thank my advisor, Professor Martin Sheen, for his support and some great action movies. I would also like to thank my dog, Spot, who only ate my homework two or three times. I have several other people I would like to thank, as well.¹

¹To my brother and sister, who are really cool.

Efficient Implementation of Interpolation Algorithms for EUF and Octagonal Theories

by

José Abel Castellanos Joo

B.Tech., Universidad de las Américas Puebla, 2015

M.S., Computer Science, University of New Mexico, 2019

Abstract

The theory of relativity is a real “toughie” to prove, but with the help of my family and my great grandpa Al, this paper presents the proof in its entirety. Most of the math is correct, and the part about “warp speed” and “parallel universe” sounds very high-tech.

Contents

| | |
|---|----------|
| List of Figures | vii |
| List of Tables | viii |
| Glossary | ix |
| 1 Introduction | 1 |
| 1.1 Overview | 1 |
| 1.2 Conclusions | 1 |
| 2 Related Work | 2 |
| 3 Background | 3 |
| 4 The Theory of EUF | 4 |
| 5 The Theory of Octagonal Formulas | 5 |
| 6 Combining Theories: EUF and Octagonal Formulas | 7 |

Contents

| | |
|----------------------|-----------|
| 7 Future Work | 8 |
| Appendices | 9 |
| References | 10 |

List of Figures

List of Tables

Glossary

| | |
|------------------|---|
| a_{lm} | Taylor series coefficients, where $l, m = \{0..2\}$ |
| $A_{\mathbf{p}}$ | Complex-valued scalar denoting the amplitude and phase. |
| A^T | Transpose of some relativity matrix. |

Chapter 1

Introduction

1.1 Overview

What???

1.2 Conclusions

I conclude that this is a really short thesis.

Chapter 2

Related Work

Chapter 3

Background

Chapter 4

The Theory of EUF

Theorem 4.0.1. *Let t be an uncommon term and let H be a collection of Horn equations. Assume $\text{antecedent}(H) \neq \{\}$. If $\text{consequent}(t) = \{\}$, then we cannot conditionally eliminate t from any Horn equation $h \in H$.*

Proof. Suppose, by contradiction, that we can conditionally eliminate the term t from H . Then, there exists $h' \in H$ such that t appears in the consequent of h' . But $\text{consequent}(t) = \{\}$, contradiction. \square

Corollary 4.0.1.1. *We cannot eliminate an uncommon term t unconditionally once $\text{consequent}(t) = \{\}$.*

Chapter 5

The Theory of Octagonal Formulas

Theorem 5.0.1. *Given a mutually contradictory pair (α, β) , where α, β are finite conjunctions of octagonal atoms, the above algorithm terminates with an interpolant I_α that is a finite conjunction of octagonal atoms and is equivalent to $\exists \vec{x}.\alpha$, where \vec{x} is the symbols in α which are not in β . Further I_α is the strongest interpolant for (α, β) .*

Proof. First we will prove that $\models \alpha \rightarrow I_\alpha$. The latter follows since the algorithm produces a conjunction of octagonal formulas using the Elim rule, which is a truth-preserving rule of inference, eliminating conjuncts with uncommon symbols.

Now, we will prove that $\not\models I_\alpha \wedge \beta$. We will prove the latter by induction on the number of variables to eliminate (k).

- Base case: $k = 0$. Then the algorithm outputs α . Then the statement holds since (α, β) is unsatisfiable.
- Inductive case: $k = n + 1$. Since the set S of variables to eliminate is non-empty, we just take any variable $x \in S$ and apply the above algorithm to

Chapter 5. The Theory of Octagonal Formulas

eliminate such variable. Let X be the set of octagonal inequalities of (α, β) and X' be the set of octagonal inequalities (α', β) where α' is the conjunct obtained after removing the variable x . We know X and X' are equisatisfiable using a similar argument as in the Fourier-Motzkin elimination method [1]. Let us suppose (α', β) is satisfiable, hence (α, β) is unsatisfiable as well. But the latter entails a contradiction since (α, β) is assumed to be unsatisfiable. Hence, (α', β) is unsatisfiable. Since (α', β) is an unsatisfiable formula with n variables to eliminate, using the Inductive Hypothesis we conclude that (I_α, β) is unsatisfiable as well.

□

Chapter 6

Combining Theories: EUF and Octagonal Formulas

Chapter 7

Future Work

I'm sure my future work will consist of lots of other famous stuff.

Appendices

Bibliography

- [1] Alexander Schrijver. *Theory of Linear and Integer Programming*. John Wiley & Sons, Inc., New York, NY, USA, 1986.