

# **IB Linear Algebra**

Ishan Nath, Michaelmas 2022

Based on Lectures by Prof. Pierre Raphael

October 30, 2022

## Contents

<b>1</b>	<b>Vector Spaces and Subspaces</b>	<b>2</b>
1.1	Subspaces and Quotients . . . . .	4
<b>2</b>	<b>Spans, Linear Independence and the Steinitz Exchange Lemma</b>	<b>5</b>
<b>3</b>	<b>Basis, Dimension and Direct Sums</b>	<b>9</b>
<b>4</b>	<b>Linear maps, Isomorphisms and the Rank-Nullity Theorem</b>	<b>13</b>
<b>5</b>	<b>Linear maps and Matrices</b>	<b>17</b>
<b>6</b>	<b>Change of Basis and Equivalent Matrices</b>	<b>20</b>
6.1	Change of Basis . . . . .	20
<b>7</b>	<b>Elementary operations and Elementary Matrices</b>	<b>24</b>
7.1	Representation of Square Invertible Matrix . . . . .	25
<b>8</b>	<b>Dual Spaces and Dual Maps</b>	<b>27</b>
<b>9</b>	<b>Properties of the Dual Map</b>	<b>31</b>
9.1	Properties of the Dual Map . . . . .	31
9.2	Double Dual . . . . .	32
<b>10</b>	<b>Bilinear Forms</b>	<b>35</b>
<b>11</b>	<b>Determinant and Traces</b>	<b>39</b>
	<b>Index</b>	<b>40</b>

# 1 Vector Spaces and Subspaces

Let  $F$  be an arbitrary field.

**Definition 1.1** ( $F$  vector space). A  $F$  vector space is an abelian group  $(V, +)$  equipped with a function

$$\begin{aligned} F \times V &\rightarrow V \\ (\lambda, v) &\mapsto \lambda v \end{aligned}$$

such that

- $\lambda(v_1 + v_2) = \lambda v_1 + \lambda v_2$ ,
- $(\lambda_1 + \lambda_2)v = \lambda_1 v + \lambda_2 v$ ,
- $\lambda(\mu v) = (\lambda\mu)v$ ,
- $1 \cdot v = v$ .

We know how to

- Sum two vectors
- Multiply a vector  $v \in V$  by a scalar  $\lambda \in F$ .

## Example 1.1.

- (i) Take  $n \in \mathbb{N}$ , then  $F^n$  is the set of column vectors of length  $n$  with elements in  $F$ . We have

$$\begin{aligned} v \in F^n, v &= \begin{pmatrix} x_1 \\ \vdots \\ x_n \end{pmatrix}, x_i \in F, \\ v + w &= \begin{pmatrix} v_1 \\ \vdots \\ v_n \end{pmatrix} + \begin{pmatrix} w_1 \\ \vdots \\ w_n \end{pmatrix} = \begin{pmatrix} v_1 + w_1 \\ \vdots \\ v_n + w_n \end{pmatrix}, \\ \lambda v &= \begin{pmatrix} \lambda v_1 \\ \vdots \\ \lambda v_n \end{pmatrix}. \end{aligned}$$

Then  $F^n$  is a  $F$  vector space.

(ii) For any set  $X$ , take

$$\mathbb{R}^X = \{f : X \rightarrow \mathbb{R}\}.$$

Then  $\mathbb{R}^X$  is an  $\mathbb{R}$  vector space.

(iii) Take  $M_{n,m}(F)$ , the set of  $n \times m$   $F$  valued matrices. Then  $M_{n,m}(F)$  is a  $F$  vector space.

*Remark.* The axiom of scalar multiplication implies that for all  $v \in V$ ,  $0 \cdot v = \mathbf{0}$ .

**Definition 1.2** (Subspace). Let  $V$  be a vector space over  $F$ . A subset  $U$  of  $V$  is a vector subspace of  $V$  (denoted  $U \leq V$ ) if

- $0 \in U$ ,
- $(u_1, u_2) \in U \times U$  implies  $u_1 + u_2 \in U$ ,
- $(\lambda, u) \in F \times U$  implies  $\lambda u \in U$ .

Note if  $V$  is an  $F$  vector space, and  $U \leq V$ , then  $U$  is an  $F$  vector space.

**Example 1.2.**

- (i) Take  $V = \mathbb{R}^{\mathbb{R}}$ , the space of functions  $f : \mathbb{R} \rightarrow \mathbb{R}$ . Let  $\mathcal{C}(\mathbb{R})$  be the space of continuous function  $f : \mathbb{R} \rightarrow \mathbb{R}$ . Then  $\mathcal{C}(\mathbb{R}) \leq \mathbb{R}^{\mathbb{R}}$ .
- (ii) Take the elements of  $\mathbb{R}^3$  which sum up to  $t$ . This is a subspace if and only if  $t = 0$ .

Note that the union of two subspaces is generally not a subspace, as it is usually not closed under addition.

**Proposition 1.1.** Let  $V$  be an  $F$  vector space, and  $U, W \leq V$ . Then  $U \cap W \leq V$ .

**Proof:** Since  $0 \in U, 0 \in W$ ,  $0 \in U \cap W$ . Now consider  $(\lambda, \mu) \in F^2$ , and  $(v_1, v_2) \in (U \cap W)^2$ . Take  $\lambda_1 v_1 + \lambda_2 v_2$ . Since  $u_1, v_1 \in U$ , this is in  $U$ . Similarly, it is in  $W$ . So it is in  $U \cap W$ , and  $U \cap W \leq V$ .

**Definition 1.3** (Sum of subspaces). Let  $V$  be an  $F$  vector space. Let  $U, W \leq V$ . Then the **sum** of  $U$  and  $W$  is the set

$$U + W = \{u + w \mid (u, w) \in U \times W\}.$$

**Proof:** Note  $0 = 0 + 0 \in U + W$ . Take  $\lambda_1 f + \lambda_2 g$ , where  $f, g \in U + W$ . Then we can write  $f = f_1 + f_2, g = g_1 + g_2$ , where  $f_1, g_1 \in U, f_2, g_2 \in W$ . Then

$$\lambda_1 f + \lambda_2 g = \lambda_1(f_1 + f_2) + \lambda_2(g_1 + g_2) = (\lambda_1 f_1 + \lambda_2 g_1) + (\lambda_1 f_2 + \lambda_2 g_2) \in U + W.$$

*Remark.*  $U + W$  is the smallest subspace of  $V$  which contains both  $U$  and  $W$ .

## 1.1 Subspaces and Quotients

**Definition 1.4** (Quotient). Let  $V$  be an  $F$  vector space. Let  $U \leq V$ . The quotient space  $V/U$  is the abelian group  $V/U$  equipped with the scalar product multiplication

$$\begin{aligned} F \times V/U &\rightarrow V/U \\ (\lambda, v + U) &\mapsto \lambda v + U \end{aligned}$$

**Proposition 1.2.**  $V/U$  is an  $F$  vector space.

## 2 Spans, Linear Independence and the Steinitz Exchange Lemma

**Definition 2.1** (Span of a family of vectors). Let  $V$  be a  $F$  vector space. Let  $S \subset V$  be a subset. We define

$$\begin{aligned}\langle S \rangle &= \{\text{finite linear combinations of elements of } S\} \\ &= \left\{ \sum_{\delta \in J} \lambda_{\delta} v_{\delta}, v_{\delta} \in S, \lambda_{\delta} \in F, J \text{ finite} \right\}.\end{aligned}$$

By convention, we let  $\langle \emptyset \rangle = \{0\}$ .

*Remark.*  $\langle S' \rangle$  is the smallest vector subspace which contains  $S$ .

**Example 2.1.** Take  $V = \mathbb{R}^3$ , and

$$S = \left\{ \begin{pmatrix} 1 \\ 0 \\ 0 \end{pmatrix}, \begin{pmatrix} 0 \\ 1 \\ 2 \end{pmatrix}, \begin{pmatrix} 3 \\ -2 \\ 4 \end{pmatrix} \right\}.$$

Then we have

$$\langle S' \rangle = \left\{ \begin{pmatrix} a \\ b \\ 2b \end{pmatrix}, (a, b) \in \mathbb{R}^2 \right\}.$$

Take  $V = \mathbb{R}^n$ , and let  $e_i$  be the  $i$ 'th basis vector. Then  $V = \langle e_1, \dots, e_n \rangle$ .

Take  $X$  a set, and  $V = \mathbb{R}^X$ . Let  $S_x : X \rightarrow \mathbb{R}$ , such that  $y \mapsto 1$  if  $x = y$ , otherwise  $y \mapsto 0$ . Then

$$\langle (S_x)_{x \in X} \rangle = \{f \in \mathbb{R}^X \mid f \text{ has finite support}\}.$$

**Definition 2.2.** Let  $V$  be a  $F$  vector space. Let  $S'$  be a subset of  $V$ . We may say that  $S$  **spans**  $V$  if  $\langle S \rangle = V$ .

**Definition 2.3** (Finite dimension). Let  $V$  be a  $F$  vector space. We say that  $V$  is **finite dimensional** if it is spanned by a finite set.

**Example 2.2.** Consider  $P[x]$ , the polynomials over  $\mathbb{R}$ , and  $P_n[x]$ , the polynomials over  $\mathbb{R}$  with degree  $\leq n$ . Then since

$$\langle 1, x, \dots, x^n \rangle = P_n[x],$$

$P_n[x]$  is finite dimensional, however  $P[x]$  is not.

**Definition 2.4** (Independence). We say that  $(v_1, \dots, v_n)$ , elements of  $V$  are **linearly independent** if

$$\sum_{i=1}^n \lambda_i v_i = 0 \implies \lambda_i = 0 \forall i.$$

*Remark.*

1. We also say that the family  $(v_1, \dots, v_n)$  is **free**.
2. Equivalently,  $(v_1, \dots, v_n)$  are not linearly independent if one of these vectors is a linear combination of the remaining  $(n-1)$ .
3. If  $(v_i)$  is free, then  $v_i = 0$  for all  $i$ .

**Definition 2.5** (Basis). A subset  $S$  of  $V$  is a **basis** of  $V$  if and only if

- (i)  $\langle S \rangle = V$ ,
- (ii)  $S$  is linearly independent.

*Remark.* A subset  $S$  that generates  $V$  is a generating family, so a basis  $S$  is a free generating family.

**Example 2.3.** For  $V = \mathbb{R}^n$ , then  $(e_i)$  is a basis of  $V$ .

If  $V = \mathbb{C}$ , then for  $F = \mathbb{C}$ ,  $\{1\}$  is a basis.

If  $V = P[x]$ , then  $S = \{x^n, n \geq 0\}$  is a basis for  $V$ .

**Lemma 2.1.**  $V$  is a  $F$  vector space. Then  $(v_1, \dots, v_n)$  is a basis of  $V$  if and only if any vector  $v \in V$  has a unique decomposition

$$v = \sum_{i=1}^n \lambda_i v_i.$$

*Remark.* We call  $(\lambda_1, \dots, \lambda_n)$  the coordinates of  $v$  in the basis  $(v_1, \dots, v_n)$ .

**Proof:** Since  $\langle v_1, \dots, v_n \rangle = V$ , we must have

$$v = \sum_{i=1}^n \lambda_i v_i$$

for some  $\lambda_i$ . Now assume

$$\begin{aligned} v &= \sum_{i=1}^n \lambda_i v_i = \sum_{i=1}^n \lambda'_i v_i, \\ \implies \sum_{i=1}^n (\lambda_i - \lambda'_i) v_i &= 0. \end{aligned}$$

Since  $v_i$  are free,  $\lambda_i = \lambda'_i$ .

**Lemma 2.2.** *If  $(v_1, \dots, v_n)$  spans  $V$ , then some subset of this family is a basis of  $V$ .*

**Proof:** If  $(v_1, \dots, v_n)$  are linearly independent, we are done. Otherwise assume they are not independent, then by possibly reordering the vectors, we have

$$v_n \in \langle v_1, \dots, v_{n-1} \rangle.$$

Then we have  $V = \langle v_1, \dots, v_n \rangle = \langle v_1, \dots, v_{n-1} \rangle$ . By iterating, we must eventually get to an independent set.

**Theorem 2.1** (Steinitz Exchange Lemma). *Let  $V$  be a finite dimensional vector space over  $F$ . Take*

- (i)  $(v_1, \dots, v_m)$  free,
- (ii)  $(w_1, \dots, w_n)$  generating.

*Then  $m \leq n$ , and up to reordering,  $(v_1, \dots, v_m, w_{m+1}, \dots, w_n)$  spans  $V$ .*

**Proof:** Induction. Suppose that we have replaced  $l$  of the  $w_i$ , reordering if necessary, so

$$\langle v_1, \dots, v_l, w_{l+1}, \dots, w_n \rangle = V.$$

If  $m = l$ , we are done. Otherwise,  $l < m$ . Then since these vectors span  $V$ , we have



$$v_{l+1} = \sum_{i \leq l} a_i v_i + \sum_{i > l} \beta_i w_i.$$

Since  $(v_1, \dots, v_{l+1})$  is free, some of the  $\beta_i$  are non-zero. Upon reordering, we may let  $\beta_{l+1} \neq 0$ . Then,

$$w_{l+1} = \frac{1}{\beta_{l+1}} \left[ v_{l+1} - \sum_{i \leq l} \alpha_i v_i - \sum_{i > l+1} \beta_i w_i \right].$$

Hence,  $V = \langle v_1, \dots, v_l, w_{l+1}, \dots, w_n \rangle = \langle v_1, \dots, v_l, v_{l+1}, w_{l+1}, \dots, w_n \rangle = \langle v_1, \dots, v_{l+1}, w_{l+2}, \dots, w_n \rangle$ . Iterating this process, we eventually get  $l = m$ , which then proves  $m \leq n$ .

### 3 Basis, Dimension and Direct Sums

**Corollary 3.1.** *Let  $V$  be a finite dimensional vector space over  $F$ . Then any two bases of  $V$  have the same number of vectors, called the **dimension** of  $V$ .*

**Proof:** take  $(v_1, \dots, v_n), (w_1, \dots, w_m)$  bases of  $V$ .

(i) As  $(v_i)$  is free and  $(w_i)$  is generating,  $n \leq m$ .

(ii) As  $(w_i)$  is free and  $(v_i)$  is generating,  $m \leq n$ .

So  $m = n$ .

**Corollary 3.2.** *Let  $V$  be a vector space over  $F$  with dimension  $n \in \mathbb{N}$ .*

- (i) *Any set of independent vectors has at most  $n$  elements, with equality if and only if it is a basis.*
- (ii) *Any spanning set of vectors has at least  $n$  elements, with equality if and only if it is a basis.*

**Proof:** Exercise (fill this in).

**Proposition 3.1.** *Let  $U, W$  be finite dimensional subspaces of  $V$ . If  $U$  and  $W$  are finite dimensional, then so is  $U + W$ , and*

$$\dim(U + W) = \dim U + \dim W - \dim(U \cap W).$$

**Proof:** Pick  $(v_1, \dots, v_l)$  a basis of  $U \cap W$ . Extend this to a basis  $(v_1, \dots, v_l, u_1, \dots, u_m)$  of  $U$ , and a basis  $(v_1, \dots, v_l, w_1, \dots, w_n)$  of  $W$ . Then we show  $(v_1, \dots, v_l, u_1, \dots, u_m, w_1, \dots, w_n)$  is a basis of  $U + W$ .

It is clearly a generating family, so we will show it is free. Suppose

$$\sum_{i=1}^l \alpha_i v_i + \sum_{i=1}^m \beta_i u_i + \sum_{i=1}^n \gamma_i w_i = 0.$$

Then we get

$$\sum_{i=1}^n \gamma_i w_i \in U \cap W,$$

implying that

$$\sum_{i=1}^l s_i v_i = \sum_{i=1}^n \gamma_i w_i.$$

But since  $(v_1, \dots, w_n)$  is a basis of  $W$ , we get  $\gamma_i = 0$ . Similarly,  $\beta_i = 0$ . Thus,

$$\sum_{i=1}^l \alpha_i v_i = 0.$$

Since  $(v_i)$  is a basis of  $U \cap W$ ,  $\alpha_i = 0$ .

**Proposition 3.2.** *Let  $V$  be a finite dimensional vector space over  $F$ . Let  $U \leq V$ . Then  $U$  and  $V/U$  are both finite dimensional and*

$$\dim V = \dim U + \dim(V/U).$$

**Proof:** Let  $(u_1, u_2, \dots, u_l)$  be a basis of  $U$ . Extend this to a basis  $(u_1, \dots, u_l, w_{l+1}, \dots, w_n)$  of  $V$ . Then we show that  $(w_{l+1} + U, \dots, w_n + U)$  is a basis of  $V/U$ . (Fill this in).

*Remark.* If  $U \leq V$ , then we say  $U$  is proper if  $U \neq V$ . Then for finite dimensions,  $U$  proper implies  $\dim U < \dim V$ , as  $\dim(V/U) > 0$ .

**Definition 3.1** (Direct sum). Let  $V$  be a vector space over  $F$ , and  $U, W \leq V$ . We say  $V = U \oplus W$  if and only if any element of  $v \in V$  can be uniquely decomposed as  $v = u + w$  for  $u \in U, w \in W$ .

*Remark.* If  $V = U \oplus W$ , we say that  $W$  is a complement of  $U$  in  $V$ . There is no uniqueness of such a complement.

In the sequel, we use the following notation. Let  $\mathcal{B}_1 = \{u_1, \dots, u_l\}$  and  $\mathcal{B}_2 = \{w_1, \dots, w_m\}$  be collections of vectors. Then

$$\mathcal{B}_1 \cup \mathcal{B}_2 = \{u_1, \dots, u_l, w_1, \dots, w_m\}$$

with the convention that  $\{v\} \cup \{v\} = \{v, v\}$ .

**Lemma 3.1.** *Let  $U, W \leq V$ . Then the following are equivalent:*

- (i)  $V = U \oplus W$ ;
- (ii)  $V = U + W$  and  $U \cap W = \{0\}$ ;
- (iii) For any basis  $\mathcal{B}_1$  of  $U$ ,  $\mathcal{B}_2$  of  $W$ , the union  $\mathcal{B} = \mathcal{B}_1 \cup \mathcal{B}_2$  is a basis of  $V$ .

**Proof:** We show (ii) implies (i). Let  $V = U + W$ , then clearly  $U, W$  generate  $V$ . We only need to show uniqueness. Suppose  $u_1 + w_1 = u_2 + w_2$ . Then

$$u_1 - u_2 = w_2 - w_1 \in U \cap W = \{0\}.$$

Hence  $u_1 = u_2$  and  $w_1 = w_2$ , as required.

Now we show (i) implies (iii). Let  $\mathcal{B}_1$  be a basis of  $U$ , and  $\mathcal{B}_2$  a basis of  $W$ . Then  $\mathcal{B} = \mathcal{B}_1 \cup \mathcal{B}_2$  generates  $U + W = V$ , and  $\mathcal{B}$  is free, as if  $\sum \lambda_i v_i = u + w = 0$ , then  $0 = 0 + 0$  uniquely, so  $u = 0, w = 0$ , giving  $\lambda_i = 0$  for all  $i$ .

Finally, we show (iii) implies (ii). Let  $\mathcal{B} = \mathcal{B}_1 \cup \mathcal{B}_2$ . Then since  $\mathcal{B}$  is a basis of  $V$ ,

$$v = \sum_{u_i \in \mathcal{B}_1} \lambda_i u_i + \sum_{w_i \in \mathcal{B}_2} \lambda_i w_i = u + w.$$

Now if  $v \in U \cap W$ ,

$$v = \sum_{u \in \mathcal{B}_1} \lambda_u u = \sum_{w \in \mathcal{B}_2} \lambda_w w.$$

This gives

$$\sum_{u \in \mathcal{B}_1} \lambda_u u - \sum_{w \in \mathcal{B}_2} \lambda_w w = 0.$$

Since  $\mathcal{B}_1 \cup \mathcal{B}_2$  is free, we get  $\lambda_u = \lambda_w = 0$ , so  $U \cap W = \{0\}$ .

**Definition 3.2.** Let  $V$  be a vector space over  $F$ , and  $V_1, \dots, V_l \leq V$ . Then

(i) The sum of the subspaces is

$$\sum_{i=1}^l V_i = \{v_1 + \cdots + v_l \mid v_j \in V_j, 1 \leq j \leq l\}.$$

(ii) The sum is direct:

$$\sum_{i=1}^l V_i = \bigoplus_{i=1}^l V_i$$

if and only if

$$v_1 + \cdots + v_l = v'_1 + \cdots + v'_l \implies v_1 = v'_1, \dots, v_l = v'_l.$$

**Proof:** Exercise.

**Proposition 3.3.** *The following are equivalent:*

(i)

$$\sum_{i=1}^l V_i = \bigoplus_{i=1}^l V_i,$$

(ii)

$$\forall i, V_i \cap \left( \sum_{j < i} V_j \right) = \{0\},$$

(iii) *For any basis  $\mathcal{B}_i$  of  $V_i$ ,*

$$\mathcal{B} = \bigcup_{i=1}^l \mathcal{B}_i \text{ is a basis of } \sum_{i=1}^l V_i.$$

## 4 Linear maps, Isomorphisms and the Rank-Nullity Theorem

**Definition 4.1** (Linear map). Let  $V, W$  be vector spaces over  $F$ . A map  $\alpha : V \rightarrow W$  is **linear** if and only if for all  $\lambda_1, \lambda_2 \in F$  and  $v_1, v_2 \in V$ , we have

$$\alpha(\lambda_1 v_1 + \lambda_2 v_2) = \lambda_1 \alpha(v_1) + \lambda_2 \alpha(v_2).$$

**Example 4.1.**

- (i) Take an  $m \times n$  matrix  $M$ , Then we can take the linear map  $\alpha : \mathbb{R}^m \rightarrow \mathbb{R}^n$  defined by  $X \mapsto MX$ .
- (ii) Take the linear map  $\alpha : \mathcal{C}[0, 1] \rightarrow \mathcal{C}[0, 1]$  by

$$f \mapsto \alpha(f)(x) = \int_0^x f(t) dt.$$

- (iii) Fix  $x \in [a, b]$ . Then we can take a linear map  $\mathcal{C}[a, b] \rightarrow \mathbb{R}$  by  $f \mapsto f(x)$ .

*Remark.* Let  $U, V, W$  be  $F$ -vector spaces.

- (i) The identity map  $\text{id}_V : V \rightarrow V$  by  $x \mapsto x$  is a linear map.
- (ii) If  $U \rightarrow V$  is  $\beta$  linear, and  $V \rightarrow W$  is  $\alpha$  linear, then  $U \rightarrow W$  is linear by  $\alpha \circ \beta$ .

**Lemma 4.1.** Let  $V, W$  be  $F$ -vector spaces, and  $\mathcal{B}$  a basis of  $V$ . Let  $\alpha_0 : \mathcal{B} \rightarrow W$  be any map, then there is a unique linear map  $\alpha : V \rightarrow W$  extending  $\alpha_0$ .

**Proof:** For  $v \in V$ , we can write

$$v = \sum_{i=1}^n \lambda_i v_i,$$

where  $\mathcal{B} = (v_1, \dots, v_n)$ . Then by linearity, we must have

$$\alpha(v) = \alpha \left( \sum_{i=1}^n \lambda_i v_i \right) = \sum_{i=1}^n \lambda_i \alpha_0(v_i).$$

This is unique as  $\mathcal{B}$  is a basis.

*Remark.* This is true in infinite dimensions as well.

Often, to define a linear map, we define its value on a basis and extend by linearity. As a corollary, if  $\alpha_1, \alpha_2 : V \rightarrow W$  are linear and agree on a basis of  $V$ , they are equal.

**Definition 4.2** (Isomorphism). Let  $V, W$  be vector spaces over  $F$ . A map  $\alpha : V \rightarrow W$  is called an **isomorphism** if and only if  $\alpha$  is linear and bijective. If such an  $\alpha$  exists, we say  $V \cong W$ .

*Remark.* If  $\alpha : V \rightarrow W$  is an isomorphism, then  $\alpha^{-1} : W \rightarrow V$  is linear. Indeed, for  $w_1, w_2 \in W \times W$ , let  $w_1 = \alpha(v_1), w_2 = \alpha(v_2)$ . Then,

$$\begin{aligned} \alpha^{-1}(\lambda_1 w_1 + \lambda_2 w_2) &= \alpha^{-1}(\lambda_1 \alpha(v_1) + \lambda_2 \alpha(v_2)) \\ &= \alpha^{-1}(\alpha(\lambda_1 v_1 + \lambda_2 v_2)) \\ &= \lambda_1 v_1 + \lambda_2 v_2 \\ &= \lambda_1 \alpha^{-1}(w_1) + \lambda_2 \alpha^{-1}(w_2). \end{aligned}$$

**Lemma 4.2.** *Congruence is an equivalence relation on the class of all vector spaces of  $F$ :*

- (i)  $\text{id}_V : V \rightarrow V$  is an isomorphism.
- (ii)  $\alpha : V \rightarrow W$  is an isomorphism implies  $\alpha^{-1} : W \rightarrow V$  is an isomorphism.
- (iii) If  $\alpha : U \rightarrow V$  is an isomorphism,  $\beta : V \rightarrow W$  is an isomorphism, then  $\beta \circ \alpha : U \rightarrow W$  is an isomorphism.

**Proof:** Exercise.

**Theorem 4.1.** *If  $V$  is a vector space over  $F$  of dimension  $n$ , then  $V \cong F^n$ .*

**Proof:** Let  $\mathcal{B} = (v_1, \dots, v_n)$  be a basis of  $V$ . Then take

$$\alpha : V \rightarrow F^n$$

$$v = \sum_{i=1}^n \lambda_i v_i \mapsto \begin{pmatrix} \lambda_1 \\ \vdots \\ \lambda_n \end{pmatrix}$$

as an isomorphism.

*Remark.* In this way, choosing a basis of  $V$  is like choosing an isomorphism from  $V$  to  $F^n$ .

**Theorem 4.2.** *Let  $V, W$  be vector spaces over  $F$  with finite dimension. Then  $V \cong W$  if and only if  $\dim V = \dim W$ .*

**Proof:** If  $\dim V = \dim W$ , then  $V \cong F^n \cong W$ , so  $V \cong W$ .

Otherwise, let  $\alpha : V \rightarrow W$  be an isomorphism, and  $\mathcal{B}$  a basis of  $V$ . Then we show  $\alpha(\mathcal{B})$  is a basis of  $W$ .

- $\alpha(\mathcal{B})$  spans  $W$  from the surjectivity of  $\alpha$ .
- $\alpha(\mathcal{B})$  is free from the injectivity of  $\alpha$ .

Hence  $\dim V = \dim W$ .

**Definition 4.3** (Kernal and Image). Let  $V, W$  be vector spaces over  $F$ . Let  $\alpha : V \rightarrow W$  be a linear map. We define

- (i)  $\text{Ker } \alpha = \{v \in V \mid \alpha(v) = 0\}$ , the kernel of  $\alpha$ .
- (ii)  $\text{Im}(\alpha) = \{w \in W \mid \exists v \in V, \alpha(v) = w\}$ , the image of  $\alpha$ .

**Lemma 4.3.**  *$\text{Ker } \alpha$  is a subspace of  $V$ , and  $\text{Im } \alpha$  is a subspace of  $W$ .*

**Proof:** Let  $\lambda_1, \lambda_2 \in F$ , and  $v_1, v_2 \in \text{Ker } \alpha$ . Then

$$\alpha(\lambda_1 v_1 + \lambda_2 v_2) = \lambda_1 \alpha(v_1) + \lambda_2 \alpha(v_2) = 0.$$

So  $\lambda_1 v_1 + \lambda_2 v_2 \in \text{Ker } \alpha$ .

Now if  $w_1 = \alpha(v_1), w_2 = \alpha(v_2)$ , then

$$\lambda_1 w_1 + \lambda_2 w_2 = \lambda_1 \alpha(v_1) + \lambda_2 \alpha(v_2) = \alpha(\lambda_1 v_1 + \lambda_2 v_2).$$

Hence  $\lambda_1 w_1 + \lambda_2 w_2 \in \text{Im } \alpha$ .

**Example 4.2.** Consider  $\alpha : \mathcal{C}^\infty(\mathbb{R}) \rightarrow \mathcal{C}^\infty(\mathbb{R})$ , given by

$$f \mapsto \alpha(f) = f'' + f.$$

Then  $\alpha$  is linear, and

$$\text{Ker } \alpha = \{f \in \mathcal{C}^\infty(\mathbb{R}) \mid f'' + f = 0\} = \langle \sin t, \cos t \rangle.$$



*Remark.* If  $\alpha : V \rightarrow W$  is linear, then  $\alpha$  is injective if and only if  $\text{Ker } \alpha = \{0\}$ , as

$$\alpha(v_1) = \alpha(v_2) \iff \alpha(v_1 - v_2) = 0.$$

**Theorem 4.3.** *Let  $V, W$  be vector spaces over  $F$ , and  $\alpha : V \rightarrow W$  linear. Then*

$$\begin{aligned} V / \text{Ker } \alpha &\rightarrow \text{Im } \alpha \\ v + \text{Ker } \alpha &\mapsto \alpha(v) \end{aligned}$$

*is an isomorphism.*

**Proof:** We proceed in steps.

- $\bar{\alpha}$  is well defined: Note if  $v + \text{Ker } \alpha = v' + \text{Ker } \alpha$ , then  $v - v' \in \text{Ker } \alpha$ , so  $\alpha(v - v') = 0$ . Hence  $\alpha(v) = \alpha(v')$ .
- $\bar{\alpha}$  is linear: This follows from linearity of  $\alpha$ .
- $\bar{\alpha}$  is a bijection: First, if  $\bar{\alpha}(v + \text{Ker } \alpha) = 0$ , then  $\alpha(v) = 0$ , so  $v \in \text{Ker } \alpha$ , hence  $v + \text{Ker } \alpha = 0 + \text{Ker } \alpha$ , so  $\alpha$  is injective. Then  $\bar{\alpha}$  is surjective from the definition of the image.

**Definition 4.4** (Rank and Nullity). We define the rank  $r(\alpha) = \text{rank}(\alpha) = \dim \text{Im } \alpha$ , and the nullity  $n(\alpha) = \text{null}(\alpha) = \dim \text{Ker } \alpha$ .

**Theorem 4.4** (Rank-nullity theorem). *Let  $U, V$  be vector spaces over  $F$ , with  $\dim U < \infty$ , and let  $\alpha : U \rightarrow V$  be a linear map. Then,*

$$\dim U = r(\alpha) + n(\alpha).$$

**Proof:** We have proven that  $U / \text{Ker } \alpha \cong \text{Im } \alpha$ , but we have already proven  $\dim U / \text{Ker } \alpha = \dim U - r(\alpha)$ , which proves the theorem.

**Lemma 4.4.** *Let  $V, W$  be vector spaces over  $F$  of equal finite dimension. Let  $\alpha : V \rightarrow W$  be a linear map. Then the following are equivalent:*

- $\alpha$  is injective,
- $\alpha$  is surjective,
- $\alpha$  is an isomorphism.

This follows immediately from the rank-nullity theorem.

## 5 Linear maps and Matrices

**Definition 5.1.** If  $V, W$  are vector spaces over  $F$ , then

$$L(V, W) = \{\alpha : V \rightarrow W \text{ linear}\}.$$

**Proposition 5.1.**  $L(V, W)$  is a vector space over  $F$  with

$$(\alpha_1 + \alpha_2)(v) = \alpha_1(v) + \alpha_2(v),$$

$$(\lambda\alpha)(v) = \lambda\alpha(v).$$

Moreover, if  $V$  and  $W$  are finite dimensional, then so is  $L(V, W)$ , and

$$\dim L(V, W) = \dim V \dim W.$$

**Definition 5.2.** An  $m \times n$  matrix over  $F$  is an array with  $m$  rows and  $n$  columns with entries in  $F$ ,  $A = (a_{ij})$ . Define

$$M_{m,n}(F) = \{\text{set of } m \times n \text{ matrices over } F\}.$$

**Proposition 5.2.**  $M_{m,n}(F)$  is a vector space over  $F$ , and  $\dim M_{m,n}(F) = mn$

**Proof:** Let  $E_{ij}$  be the matrix with  $a_{xy} = \delta_{xi}\delta_{yj}$ . Then  $(E_{ij})$  is a basis of  $M_{m,n}(F)$ , as

$$N = (a_{ij}) = \sum_{i,j} a_{ij} E_{ij},$$

and  $(E_{ij})$  is free.

If  $V, W$  are vector spaces over  $F$ , and  $\alpha : V \rightarrow W$  is a linear map, we take a basis  $\mathcal{B} = (v_1, \dots, v_n)$  of  $V$ , and  $\mathcal{C} = (w_1, \dots, w_m)$  of  $W$ . Let  $v \in V$ , then

$$v = \sum_{i=1}^n \lambda_i v_i \sim \begin{pmatrix} \lambda_1 \\ \vdots \\ \lambda_n \end{pmatrix} \in F^n.$$

We let this isomorphism from  $V$  to  $F^n$  be  $[v]_{\mathcal{B}}$ . Similarly, we can obtain  $[w]_{\mathcal{C}}$  for  $w \in W$ .

**Definition 5.3.** We define a matrix of  $\alpha$  with respect to a basis  $\mathcal{B}, \mathcal{C}$  as

$$[\alpha]_{\mathcal{B}, \mathcal{C}} = ([\alpha(v_1)]_{\mathcal{C}}, [\alpha(v_2)]_{\mathcal{C}}, \dots, [\alpha(v_n)]_{\mathcal{C}}).$$

By definition, if  $[\alpha]_{\mathcal{B},\mathcal{C}} = (a_{ij})$ , then

$$\alpha(v_j) = \sum_{i=1}^m a_{ij} w_i.$$

**Lemma 5.1.** *If  $v \in V$ , then*

$$[\alpha(v)]_{\mathcal{C}} = [\alpha]_{\mathcal{B},\mathcal{C}} \cdot [v]_{\mathcal{B}},$$

*or equivalently,*

$$(\alpha(v))_i = \sum_{j=1}^n a_{ij} \lambda_j.$$

**Proof:** Let  $v \in V$ , then

$$v = \sum_{j=1}^n \lambda_j v_j.$$

Then

$$\begin{aligned} \alpha(v) &= \alpha\left(\sum_{j=1}^n \lambda_j v_j\right) = \sum_{j=1}^n \lambda_j \alpha(v_j) \\ &= \sum_{j=1}^n \lambda_j \sum_{i=1}^m a_{ij} w_i = \sum_{i=1}^m \left(\sum_{j=1}^n a_{ij} \lambda_j\right) w_i. \end{aligned}$$

**Lemma 5.2.** *If  $U \rightarrow V$  is linear under  $\beta$ ,  $V \rightarrow W$  linear under  $\alpha$ , then  $U \rightarrow W$  is linear under  $\alpha \circ \beta$ . Let  $\mathcal{A}$  be a basis of  $U$ ,  $\mathcal{B}$  a basis of  $V$ , and  $\mathcal{C}$  a basis of  $W$ . Then*

$$[\alpha \circ \beta]_{\mathcal{A},\mathcal{C}} = [\alpha]_{\mathcal{B},\mathcal{C}} \cdot [\beta]_{\mathcal{A},\mathcal{B}}.$$

**Proof:** Let  $A = [\alpha]_{\mathcal{B},\mathcal{C}}$ ,  $B = [\beta]_{\mathcal{A},\mathcal{B}}$ . Pick  $u_l \in A$ . Then

$$\begin{aligned} (\alpha \circ \beta)(u_l) &= \alpha(\beta(u_l)) = \alpha\left(\sum_j b_{jl} v_j\right) \\ &= \sum_j b_{jl} \alpha(v_j) = \sum_j b_{jl} \sum_i a_{ij} w_i \\ &= \sum_i \left(\sum_j a_{ij} b_{jl}\right) w_i. \end{aligned}$$

**Proposition 5.3.** *If  $V$  and  $W$  are vector spaces over  $F$ , and  $\dim V = n$ ,  $\dim W = m$ , then  $L(V, W) \cong M_{m,n}(F)$ , so  $\dim L(V, W) = m \times n$ .*

**Proof:** Fix  $\mathcal{B}, \mathcal{C}$  bases of  $V$  and  $W$ . We show

$$\begin{aligned}\theta : L(V, W) &\rightarrow M_{m,n}(F) \\ \alpha &\mapsto [\alpha]_{\mathcal{B}, \mathcal{C}}\end{aligned}$$

is an isomorphism.

- $\theta$  is linear:  $[\lambda_1 \alpha_1 + \lambda_2 \alpha_2]_{\mathcal{B}, \mathcal{C}} = \lambda_1 [\alpha_1]_{\mathcal{B}, \mathcal{C}} + \lambda_2 [\alpha_2]_{\mathcal{B}, \mathcal{C}}$ .
- $\theta$  is surjective: Consider  $A = (a_{ij})$ . Consider the map

$$\alpha : v_j \mapsto \sum_{i=1}^m a_{ij} w_i.$$

This can be extended by linearity, and  $[\alpha]_{\mathcal{B}, \mathcal{C}} = A$ .

- $\theta$  is injective: If  $[\alpha]_{\mathcal{B}, \mathcal{C}} = 0$ , then  $\alpha = 0$  for all  $v$ .

*Remark.* If  $\mathcal{B}, \mathcal{C}$  are bases of  $V, W$  and  $\varepsilon_{\mathcal{B}} : v \mapsto [v]_{\mathcal{B}}$ ,  $\varepsilon_{\mathcal{C}} : w \mapsto [w]_{\mathcal{C}}$ , then the following diagram commutes:

$$\begin{array}{ccc} V & \xrightarrow{\alpha} & W \\ \downarrow \varepsilon_{\mathcal{B}} & & \downarrow \varepsilon_{\mathcal{C}} \\ F^n & \xrightarrow{[\alpha]_{\mathcal{B}, \mathcal{C}}} & F^m \end{array}$$

## 6 Change of Basis and Equivalent Matrices

Let  $\alpha : V \rightarrow W$  with  $\mathcal{B}$  and  $\mathcal{C}$  bases of  $V, W$ . Then

$$[\alpha(v)]_{\mathcal{C}} = [\alpha]_{\mathcal{B}, \mathcal{C}} \cdot [v]_{\mathcal{B}}.$$

If  $Y \leq V$ , we can take  $\mathcal{B}$  a basis of  $V$ , such that  $(v_1, \dots, v_k, v_{k+1}, \dots, v_n)$  is a basis of  $V$ , and  $(v_1, \dots, v_k)$  is a basis  $\mathcal{B}'$  of  $Y$ , and  $(v_{k+1}, \dots, v_n)$  is a basis  $\mathcal{B}''$ .

Then if  $Z \leq W$ , we can take a basis  $\mathcal{C}$  of  $W$   $(w_1, \dots, w_l, w_{l+1}, \dots, w_m)$ , such that  $(w_1, \dots, w_l)$  is a basis  $\mathcal{C}'$  of  $Z$ , and  $(w_{l+1}, \dots, w_m)$  is a basis  $\mathcal{C}''$ . Then

$$[\alpha]_{\mathcal{B}, \mathcal{C}} = \begin{pmatrix} A & B \\ 0 & C \end{pmatrix}.$$

Then we can show that

$$A = [\alpha|_Y]_{\mathcal{B}', \mathcal{C}'},$$

if  $\alpha(Y) \leq Z$ . Moreover, we can show  $\alpha$  induces a homomorphism

$$\begin{aligned} \bar{\alpha} : V/Y &\rightarrow W/Z \\ v + Y &\mapsto \alpha(v) + Z \end{aligned}$$

This is well-defined as  $\alpha(v) \in Z$  for  $v \in Y$ , and  $[\bar{\alpha}]_{\mathcal{B}'', \mathcal{C}''} = C$ .

### 6.1 Change of Basis

Consider  $\alpha : V \rightarrow W$ , where  $V$  has two bases  $\mathcal{B} = \{v_1, \dots, v_n\}$  and  $\mathcal{B}' = \{v'_1, \dots, v'_n\}$  and  $W$  has two bases  $\mathcal{C} = \{w_1, \dots, w_n\}$  and  $\mathcal{C}' = \{w'_1, \dots, w'_m\}$ . We aim to find the relation between  $[\alpha]_{\mathcal{B}, \mathcal{C}}$  and  $[\alpha]_{\mathcal{B}', \mathcal{C}'}$ .

**Definition 6.1.** The change of basis matrix from  $\mathcal{B}'$  to  $\mathcal{B}$  is  $P = (p_{ij})$  given by

$$P = ([v'_1]_{\mathcal{B}}, \dots, [v'_n]_{\mathcal{B}}) = [\text{id}]_{\mathcal{B}', \mathcal{B}}.$$

**Lemma 6.1.**  $[v]_{\mathcal{B}} = P[v]_{\mathcal{B}'}$ .

**Proof:** In general  $[\alpha(v)]_{\mathcal{C}} = [\alpha]_{\mathcal{B}, \mathcal{C}}[v]_{\mathcal{B}}$ . If  $P = [\text{id}]_{\mathcal{B}', \mathcal{B}}$ , then

$$[v]_{\mathcal{B}} = [\text{id}(v)]_{\mathcal{B}} = [\text{id}]_{\mathcal{B}', \mathcal{B}}[v]_{\mathcal{B}'} = P[v]_{\mathcal{B}'}.$$

*Remark.*  $P$  is an  $n \times n$  invertible matrix, and  $P^{-1}$  is the change of basis matrix from  $\mathcal{B}$  to  $\mathcal{B}'$ . Indeed,

$$[\text{id}]_{\mathcal{B}, \mathcal{B}'}[\text{id}]_{\mathcal{B}', \mathcal{B}} = [\text{id}]_{\mathcal{B}', \mathcal{B}'} = \text{id},$$

and similarly.

Note while we know  $[v]_{\mathcal{B}} = P[v]_{\mathcal{B}'}$ , to compute a vector in  $\mathcal{B}'$ , we have  $[v]_{\mathcal{B}'} = P^{-1}[v]_{\mathcal{B}}$ . This is hard to do.

Similarly, we can also change basis  $\mathcal{C}$  to  $\mathcal{C}'$  in  $W$ . In this case, the change of basis matrix  $Q = [\text{id}]_{\mathcal{C}', \mathcal{C}}$  is  $m \times m$  and invertible.

Now given  $\alpha : V \rightarrow W$ , we wish to find how  $[\alpha]_{\mathcal{B}, \mathcal{C}}$  and  $[\alpha]_{\mathcal{B}', \mathcal{C}'}$ .

**Proposition 6.1.** *If  $A = [\alpha]_{\mathcal{B}, \mathcal{C}}$ ,  $A' = [\alpha]_{\mathcal{B}', \mathcal{C}'}$ ,  $P = [\text{id}]_{\mathcal{B}', \mathcal{B}}$ ,  $Q = [\text{id}]_{\mathcal{C}', \mathcal{C}}$ , then*

$$A' = Q^{-1}AP.$$

**Proof:** Combining the facts we know, we get

$$[\alpha(v)]_{\mathcal{C}} = Q[\alpha(v)]_{\mathcal{C}'} = Q[a]_{\mathcal{B}', \mathcal{C}'}[v]_{\mathcal{B}'} = QA'[v]_{\mathcal{B}'}.$$

But we also know

$$[\alpha(v)]_{\mathcal{C}} = [\alpha]_{\mathcal{B}, \mathcal{C}}[v]_{\mathcal{B}} = AP[v]_{\mathcal{B}'}.$$

But since this is true for any  $v \in V$ , we get  $QA' = AP$ , so  $A' = Q^{-1}AP$ .

**Definition 6.2** (Equivalent matrices). Two matrices  $A, B \in M_{m,n}(F)$  are equivalent if  $A' = Q^{-1}AP$ , where  $Q \in M_{m,m}$  and  $P \in M_{n,n}$  are invertible.

*Remark.* This defines an equivalence relation on  $M_{m,n}(F)$ , as

- $A = I_m^{-1}AI_n$ ,
- If  $A' = Q^{-1}AP$ , then  $A = (Q^{-1})^{-1}A'P^{-1}$ ,
- If  $A' = Q^{-1}AP$ ,  $A'' = (Q')^{-1}A'P'$ , then  $A'' = (QQ')^{-1}A(PP')$ .

**Proposition 6.2.** *Let  $V, W$  be vector spaces over  $F$ , with  $\dim_F V = n$ ,  $\dim_F W = m$ . Let  $\alpha : V \rightarrow W$  be a linear map. Then there exists  $\mathcal{B}, \mathcal{C}$  bases of  $V, W$  such that*

$$[\alpha]_{\mathcal{B}, \mathcal{C}} = \begin{pmatrix} I_r & 0 \\ 0 & 0 \end{pmatrix}.$$

**Proof:** Choose  $\mathcal{B}$  and  $\mathcal{C}$  wisely. Fix  $r \in \mathbb{N}$  such that  $\dim \text{Ker } \alpha = n - r$ . Let  $N(\alpha) = \text{Ker}(\alpha) = \{x \in V \mid \alpha(x) = 0\}$ . Fix any basis of  $N(\alpha)$ ,  $(v_{r+1}, \dots, v_n)$ , and extend it to a basis  $\mathcal{B} = (v_1, \dots, v_r, v_{r+1}, \dots, v_n)$ .

We claim that  $(\alpha(v_1), \dots, \alpha(v_r))$  is a basis of  $\text{Im } \alpha$ .

- First, if  $v = \sum \lambda_i v_i$ , then

$$\alpha(v) = \sum_{i=1}^n \lambda_i \alpha(v_i) = \sum_{i=1}^r \lambda_i \alpha(v_i).$$

Let  $y \in \text{Im } \alpha$ , so then

$$y = \sum_{i=1}^r \lambda_i \alpha(v_i).$$

So  $y \in \langle \alpha(v_1), \dots, \alpha(v_r) \rangle$ .

- Now, suppose that it is not free, so

$$\sum_{i=1}^r \lambda_i \alpha(v_i) = 0.$$

Then we get

$$\alpha \left( \sum_{i=1}^r \lambda_i v_i \right) = 0,$$

so

$$\sum_{i=1}^r \lambda_i v_i \in \text{Ker } \alpha.$$

Hence, we get that

$$\sum_{i=1}^r \lambda_i v_i = \sum_{i=1}^n \mu_i v_i.$$

But since  $(v_1, \dots, v_n)$  is a basis,  $\lambda_i = \mu_i = 0$ .

So we have  $(\alpha(v_1), \dots, \alpha(v_r))$  is a basis of  $\text{Im } \alpha$ , and  $(v_{r+1}, \dots, v_n)$  is a basis of  $\text{Ker } \alpha$ . Let  $\mathcal{C} = (\alpha(v_1), \dots, \alpha(v_r), w_{r+1}, \dots, w_m)$ . We get that

$$[\alpha]_{\mathcal{B}, \mathcal{C}} = (\alpha(v_1), \dots, \alpha(v_r), \alpha(v_{r+1}), \dots, \alpha(v_n)) = \begin{pmatrix} I_r & 0 \\ 0 & 0 \end{pmatrix}.$$

*Remark.* This proves another proof of the rank-nullity theorem:  $r(\alpha) + n(\alpha) = n$ .

**Corollary 6.1.** *Any  $m \times n$  matrix is equivalent to*

$$\begin{pmatrix} I_r & 0 \\ 0 & 0 \end{pmatrix},$$

where  $r = \text{rank}(\alpha)$ .

**Definition 6.3.** For  $a \in M_{m,n}(F)$ , the column rank  $r_c(A)$  of  $A$  is the dimension of the span of the column vectors of  $A$  in  $F^m$ . Similarly, the row rank is the column rank of  $A^T$ .

*Remark.* If  $\alpha$  is a linear map represented by  $A$  with respect to one basis, the column rank  $A$  equals the rank of  $\alpha$ .

**Proposition 6.3.** *Two matrices are equivalent if and only if  $r_c(A) = r_c(A')$ .*

**Proof:** If  $A$  and  $A'$  are equivalent then they coorespond to the same linear map  $\alpha$  except in two different bases.

Conversely, if  $r_c(A) = r_c(A') = r$ , then both  $A$  and  $A'$  are equivalent to

$$\begin{pmatrix} I_r & 0 \\ 0 & 0 \end{pmatrix},$$

hence are equivalent.

**Theorem 6.1.**  $r_c(A) = r_c(A^T)$ , so column rank equals row rank.

**Proof:** If  $r = r_c(A)$ , then

$$Q^{-1}AP = \begin{pmatrix} I_r & 0 \\ 0 & 0 \end{pmatrix}.$$

Take the transpose, to get

$$(Q^{-1}AP)^T = P^T A^T (Q^{-1})^T = P^T A^T (Q^T)^{-1} = \begin{pmatrix} I_r & 0 \\ 0 & 0 \end{pmatrix}.$$

Hence  $r_c(A^T) = r = r_c(A)$ .



## 7 Elementary operations and Elementary Matrices

This is a special case of the change of basis formula, when  $\alpha : V \rightarrow V$  is a map from a vector space to itself, called an endomorphism. Suppose  $\mathcal{B} = \mathcal{C}$  and  $\mathcal{B}' = \mathcal{C}'$ , and  $P$  is the change of basis matrix from  $\mathcal{B}'$  to  $\mathcal{B}$ . Then

$$[\alpha]_{\mathcal{B}', \mathcal{B}'} = P^{-1}[\alpha]_{\mathcal{B}, \mathcal{B}}P.$$

**Definition 7.1.** Let  $A, A'$  be  $n \times n$  matrices. We say that  $A$  and  $A'$  are similar if and only if  $A' = P^{-1}AP$  for a square invertible matrix  $P$ .

**Definition 7.2.** The elementary column operations on an  $m \times n$  matrix  $A$  are:

- (i) Swap columns  $i$  and  $j$ ;
- (ii) Replace column  $i$  by  $\lambda$  times column  $i$ ;
- (iii) Add  $\lambda$  times column  $i$  to column  $j$ , for  $i \neq j$ .

The elementary row operations are analogously defined.

Note elementary operations are invertible, and all operations can be realized through the action of elementary matrices:

- (i) For swapping columns  $i$  and  $j$ , we can take an identity matrix, but with  $a_{ij} = a_{ji} = 1$ , and  $a_{ii} = a_{jj} = 0$ .
- (ii) For multiplying column  $i$  by  $\lambda$ , we can take an identity matrix but with  $a_{ii} = \lambda$ .
- (iii) For adding  $\lambda$  times columns  $i$  to column  $j$ , we can take an identity matrix but with  $a_{ij} = \lambda$ .

An elementary columns (resp. row) operation can be done by multiplying  $A$  by the corresponding elementary matrix from the right (resp. left).

We will now show that any  $m \times n$  matrix is equivalent to

$$\begin{pmatrix} I_r & 0 \\ 0 & 0 \end{pmatrix}.$$

Start with a matrix  $A$ . If all entries are zero, we are done. Otherwise, pick  $a_{ij} = \lambda \neq 0$ . By swapping columns and rows, we can ensure  $a_{11} = \lambda$ . Multiplying column 1 by  $1/\lambda$ , we get  $a_{11} = 1$ . We can then clean out row 1 by subtracting a

suitable multiply of column 1 from every row, and similarly from column 1. This gives us a matrix

$$\begin{pmatrix} 1 & 0 & \cdots & 0 \\ 0 & & & \\ \vdots & & \tilde{A} & \\ 0 & & & \end{pmatrix}.$$

Iterating with  $\tilde{A}$ , a strictly smaller matrix, eventually gives

$$\begin{pmatrix} I_r & 0 \\ 0 & 0 \end{pmatrix} = Q^{-1}AP.$$

A variation of this is known as **Gauss' pivot algorithm**. If we only use row operations, we can reach the row-echelon form of the matrix:

- Assume that  $a_{i1} \neq 0$  for some  $i$ .
- Swap rows  $i$  and 1.
- Divide first row by  $\lambda = a_{i1}$ .
- Use 1 in  $a_{i1}$  to clean the first column.
- Iterate over all columns.

This procedure is what is usually done when solving a system of linear equations.

## 7.1 Representation of Square Invertible Matrix

**Lemma 7.1.** *If  $A$  is an  $n \times n$  square invertible matrix, then we can obtain  $I_n$  using either only row or column elementary operations.*

**Proof:** We prove for column operations; row operations are analogous. We proceed by induction on the number of rows.

- Suppose that we could write  $A$  in the form

$$\begin{pmatrix} I_h & 0 \\ * & * \end{pmatrix}.$$

Then we want to obtain the same structure as we go from  $h$  to  $h + 1$ .

- We show there exists  $j > h$  such that  $\lambda = a_{h+1,j} \neq 0$ . Otherwise, the row rank is less than  $n$ , as the first  $h + 1$  rows are linearly dependent. Hence  $\text{rank } A < n$ .
- We swap columns  $h + 1$  and  $j$ , so  $\lambda = a_{h+1,h+1} \neq 0$ , and then divide by  $\lambda$ .
- Finally, we can use the 1 in  $a_{h+1,h+1}$  to clear out the rest of the  $(h + 1)$ 'st row.

This gives  $AE_1 \dots E_c = I_n$ , or  $A^{-1} = E_1 \dots E_c$ . This is an algorithm for computing  $A^{-1}$ .

**Proposition 7.1.** *Any invertible square matrix is a product of elementary matrices.*

## 8 Dual Spaces and Dual Maps

**Definition 8.1.**  $V$  is a  $F$ -vector space. We say  $V^*$  is the dual of  $V$  if

$$V^* = L(V, F) = \{\alpha : V \rightarrow F \text{ linear}\}.$$

If  $\alpha : V \rightarrow F$  is linear, then we say  $\alpha$  is a linear form.

**Example 8.1.**

(i)  $\text{tr} : M_{n,n}(F) \rightarrow F$  is a linear map, so  $\text{tr} \in M_{n,n}^*(F)$ .

(ii) Let  $f : [0, 1] \rightarrow \mathbb{R}$  by  $x \mapsto f(x)$ , and  $Tf : \mathcal{C}^\infty([0, 1], \mathbb{R}) \rightarrow \mathbb{R}$  by

$$\phi \mapsto \int_0^1 f(x)\phi(x) \, dx.$$

Then  $Tf$  is a linear form.

**Lemma 8.1.** Let  $V$  be a vector space over  $F$  with a finite basis  $\mathcal{B} = \{e_1, \dots, e_n\}$ . Then there exists a basis for  $V^*$  given by  $\mathcal{B}^* = \{\varepsilon_1, \dots, \varepsilon_n\}$ , with

$$\varepsilon_j \left( \sum_{i=1}^n a_i e_i \right) = a_j.$$

Then  $\mathcal{B}^*$  is the dual basis of  $\mathcal{B}$ .

*Remark.* If we define the Kronecker symbols

$$\delta_{ij} = \begin{cases} 1 & i = j, \\ 0 & \text{otherwise,} \end{cases}$$

then we can equivalently define

$$\varepsilon_j \left( \sum_{i=1}^n a_i e_i \right) = a_j \iff \varepsilon_j(e_i) = \delta_{ij}.$$

**Proof:** Let  $(\varepsilon_1, \dots, \varepsilon_n)$  be defined as above.

We prove  $(\varepsilon_i)$  are free. Indeed, suppose

$$\sum_{j=1}^n \lambda_j \varepsilon_j = 0 \implies \sum_{j=1}^n \lambda_j \varepsilon_j(e_i) = 0 \implies \lambda_i = 0.$$

Now we show  $(\varepsilon_i)$  generates  $V^*$ . Pick  $\alpha \in V^*$ , then for  $x \in V$ , we have

$$\alpha(x) = \alpha\left(\sum_{j=1}^n \lambda_j e_j\right) = \sum_{j=1}^n \lambda_j \alpha(e_j).$$

On the other hand, consider the linear form

$$\sum_{j=1}^n \alpha(e_j) \varepsilon_j \in V^*.$$

Then we have

$$\begin{aligned} \sum_{j=1}^n \alpha(e_j) \varepsilon_j(x) &= \sum_{j=1}^n \alpha(e_j) \varepsilon_j\left(\sum_{k=1}^n \lambda_k e_k\right) = \sum_{j=1}^n \alpha(e_j) \sum_{k=1}^n \lambda_k \varepsilon_j(e_k) \\ &= \sum_{j=1}^n \alpha(e_j) \lambda_j = \alpha(x). \end{aligned}$$

Hence  $(\varepsilon_i)$  generates  $V^*$ .

**Corollary 8.1.** *If  $V$  is finite dimensional, then  $\dim V^* = \dim V$ .*

This is very different in infinite dimensions.

*Remark.* It is sometimes convenient to think of  $V^*$  as the space of row vector of length  $n$  over  $F$ . If  $(e_1, \dots, e_n)$  is a basis of  $v$  such that  $x = \sum x_i e_i$  and  $(\varepsilon_1, \dots, \varepsilon_n)$  is a basis of  $V^*$  such that  $\alpha = \sum \alpha_i \varepsilon_i$ , then

$$\begin{aligned} \alpha(x) &= \sum_{i=1}^n \alpha_i \varepsilon_i\left(\sum_{j=1}^n x_j e_j\right) = \sum_{i=1}^n \alpha_i \sum_{j=1}^n x_j \varepsilon_i(e_j) = \sum_{i=1}^n \alpha_i x_i \\ &= (\alpha_1 \quad \cdots \quad \alpha_n) \begin{pmatrix} x_1 \\ \vdots \\ x_n \end{pmatrix}. \end{aligned}$$

This gives a scalar product structure on  $V^*$ .

**Definition 8.2.** If  $U \leq V$ , we define the annihilator of  $U$  by

$$U^\circ = \{\alpha \in V^* \mid \alpha(u) = 0 \ \forall u \in U\}.$$

**Lemma 8.2.** (i)  $U^\circ \leq V^*$ .

(ii) If  $U \leq V$  and  $\dim V < \infty$ , then  $\dim V = \dim U + \dim U^\circ$ .

**Proof:** Suppose  $\alpha, \alpha' \in U^\circ$ . Then for all  $u \in U$ ,

$$(\alpha + \alpha')(u) = \alpha(u) + \alpha'(u) = 0,$$

and for all  $\lambda \in F$ ,  $(\lambda\alpha)(u) = \lambda\alpha(u) = 0$ . Hence  $U^\circ \leq V^*$ .

Now let  $U \leq V$ , and  $\dim V = n$ . Let  $(e_1, \dots, e_k)$  be a basis of  $U$  and complete it to a basis  $\mathcal{B} = (e_1, \dots, e_k, e_{k+1}, \dots, e_n)$  of  $V$ . Let  $(\varepsilon_1, \dots, \varepsilon_n)$  be the dual basis of  $\mathcal{B}$ . Then I claim  $U^\circ = \langle \varepsilon_{k+1}, \dots, \varepsilon_n \rangle$ .

Indeed, pick  $i > k$ , then  $\varepsilon_i(e_k) = \delta_{ik} = 0$ , so  $\varepsilon_i \in U^\circ$ . Now let  $\alpha \in U^\circ$ . Then  $(\varepsilon_1, \dots, \varepsilon_n)$  is a basis of  $V^*$  implies  $\alpha = \sum \alpha_i \varepsilon_i$ . But  $\alpha \in U^\circ \implies \alpha(e_i) = 0$ , which gives  $\alpha_i = 0$  for  $i \leq k$ . Hence  $\alpha \in \langle \varepsilon_{k+1}, \dots, \varepsilon_n \rangle$ .

**Definition 8.3.** Let  $V, W$  be vector spaces over  $F$ , and let  $\alpha \in L(V, W)$ . Then the map

$$\begin{aligned} \alpha^* : W^* &\rightarrow V^* \\ \varepsilon &\mapsto \varepsilon \circ \alpha \end{aligned}$$

is an element of  $L(W^*, V^*)$ . This is known as the dual map of  $\alpha$ .

**Proof:**  $\varepsilon \circ \alpha : V \rightarrow F$  is linear due to the linearity of  $\varepsilon$  and  $\alpha$ . Hence  $\varepsilon \circ \alpha \in V^*$ .

We show  $\alpha^*$  is linear. Let  $\theta_1, \theta_2 \in W^*$ . Then,

$$\alpha^*(\theta_1 + \theta_2) = (\theta_1 + \theta_2)(\alpha) = \theta_1 \circ \alpha + \theta_2 \circ \alpha = \alpha^*(\theta_1) + \alpha^*(\theta_2).$$

Similarly, if  $\lambda \in F$ , then

$$\alpha^*(\lambda\theta) = \lambda\alpha^*(\theta).$$

Hence  $\alpha^* \in L(W^*, V^*)$ .

**Proposition 8.1.** Let  $V, W$  be finite dimensional spaces over  $F$  with bases  $\mathcal{B}, \mathcal{C}$ . Let  $\mathcal{B}^*, \mathcal{C}^*$  be the dual bases for  $V^*, W^*$ . Then

$$[\alpha^*]_{\mathcal{C}^*, \mathcal{B}^*} = [\alpha]_{\mathcal{B}, \mathcal{C}}^T.$$

**Proof:** Let  $\mathcal{B} = (b_1, \dots, b_n)$ ,  $\mathcal{C} = (c_1, \dots, c_m)$ ,  $\mathcal{B}^* = (\beta_1, \dots, \beta_n)$ ,  $\mathcal{C}^* = (\gamma_1, \dots, \gamma_m)$ . Say  $[\alpha]_{\mathcal{B}, \mathcal{C}} = A = (a_{ij})$ . Recall  $\alpha^* : W^* \rightarrow V^*$ , so let us compute

$$\alpha^*(\gamma_r)(b_s) = \gamma_r \circ \alpha(b_s) = \gamma_r \left( \sum_t a_{ts} c_t \right) = \sum_t a_{ts} \gamma_r(c_t) = a_{rs}.$$

Say that

$$[\alpha^*]_{\mathcal{C}^*, \mathcal{B}^*} = (\alpha^*(\gamma_1) \quad \cdots \quad \alpha^*(\gamma_m)) = (m_{ij}).$$

Then we can find that

$$\alpha^*(\gamma_r) = \sum_{i=1}^n m_{ir} \beta_i,$$

so

$$\alpha^*(\gamma_r)(b_s) = m_{sr}.$$

This gives  $a_{rs} = m_{sr}$ , as desired.

## 9 Properties of the Dual Map

Recall if  $V, W$  are vector spaces over  $F$ , and  $\alpha \in L(V, W)$ , then we can construct a dual map

$$\begin{aligned}\alpha^* : W^* &\rightarrow V^* \\ \varepsilon &\mapsto \varepsilon \circ \alpha\end{aligned}$$

Moreover, if  $\mathcal{B}, \mathcal{C}$  are bases of  $V$  and  $W$ , and  $\mathcal{B}^*, \mathcal{C}^*$  are the dual bases of  $\mathcal{B}$  and  $\mathcal{C}$  respectively, then

$$[\alpha^*]_{\mathcal{C}^*, \mathcal{B}^*} = [\alpha]_{\mathcal{B}, \mathcal{C}}^T.$$

Now if  $\mathcal{E} = (e_1, \dots, e_n)$  is a basis of  $V$  and  $\mathcal{F} = (f_1, \dots, f_n)$  is another basis of  $V$ , then consider the change of basis matrix

$$P = [\text{id}]_{\mathcal{F}, \mathcal{E}}.$$

Consider  $\mathcal{E}^* = (\varepsilon_1, \dots, \varepsilon_n)$  and  $\mathcal{F}^* = (\eta_1, \dots, \eta_n)$ .

**Lemma 9.1.** *The change of basis matrix from  $\mathcal{F}^*$  to  $\mathcal{E}^*$  is*

$$(P^{-1})^T.$$

**Proof:** We have

$$[\text{id}]_{\mathcal{F}^*, \mathcal{E}^*} = [\text{id}]_{\mathcal{E}, \mathcal{F}}^T = ([\text{id}]_{\mathcal{F}, \mathcal{E}}^{-1})^T.$$

### 9.1 Properties of the Dual Map

**Lemma 9.2.** *Let  $V, W$  be vector spaces over  $F$ . Let  $\alpha \in L(V, W)$  and  $\alpha^* \in L(W^*, V^*)$ . Then*

- (i)  $\text{Ker}(\alpha^*) = (\text{Im } \alpha)^\circ$ . Hence  $\alpha^*$  is injective if and only if  $\alpha$  is surjective.
- (ii)  $\text{Im } \alpha^* \leq (\text{Ker } \alpha)^\circ$  with equality if  $V, W$  are finite dimensional. Hence in this case,  $\alpha^*$  is injective if and only if  $\alpha$  is injective.

There are many problems where the understanding of  $\alpha^*$  is simpler than the understanding of  $\alpha$ .



**Proof:**

- (i) Let  $\varepsilon \in W^*$ . Then  $\varepsilon \in \text{Ker } \alpha^* \iff \alpha^*(\varepsilon) = 0$ . But  $\alpha^*(\varepsilon) = \varepsilon(\alpha)$ , so for all  $x$ ,

$$\varepsilon(\alpha)(x) = \varepsilon(\alpha(x)) = 0.$$

This holds if and only if  $\varepsilon \in (\text{Im } \alpha)^\circ$ .

- (ii) We will first show that

$$\text{Im } \alpha^* \leq (\text{Ker } \alpha)^\circ.$$

Indeed, if  $\varepsilon \in \text{Im } \alpha^*$ , then  $\varepsilon = \alpha^*(\phi)$ , so for all  $u \in \text{Ker } \alpha$ ,

$$\varepsilon(u) = \alpha^*(\phi)(u) = \phi \circ \alpha(u) = \phi(0) = 0.$$

Hence  $\varepsilon \in (\text{Ker } \alpha)^\circ$ . In finite dimension, we can compare the dimension of  $\text{Im } \alpha^*$  and  $(\text{Ker } \alpha)^\circ$ . Indeed,

$$\dim(\text{Im } \alpha^*) = r(\alpha^*) = r([\alpha^*]_{\mathcal{C}^*, \mathcal{B}^*}) = r([\alpha]_{\mathcal{B}, \mathcal{C}}^T) = r([\alpha]_{\mathcal{B}, \mathcal{C}}) = r(\alpha).$$

Hence, we get

$$\dim(\text{Im } \alpha^*) = r(\alpha^*) = r(\alpha) = \dim V - \dim \text{Ker } \alpha = \dim[(\text{Ker } \alpha)^\circ].$$

Since the dimensions are the same, we get  $\text{Im } \alpha^* = (\text{Ker } \alpha)^\circ$ .

## 9.2 Double Dual

If  $V$  is a vector space over  $F$ , then  $V^* = L(V, F)$ .

We define the **bidual** as

$$V^{**} = (V^*)^* = L(V^*, F).$$

This is a very important space in infinite dimension. In general, there is no obvious connection between  $V$  and  $V^*$ . However, there is a large class of function spaces such that

$$V \cong V^{**}.$$

This is known as a reflexive space.

**Example 9.1.** For  $p > 2$ , define

$$L^p(\mathbb{R}) = \left\{ f : \mathbb{R} \rightarrow \mathbb{R} \mid \int_{\mathbb{R}} |f(x)|^p dx < \infty \right\}.$$

This is an example of a reflexive space.

In general, there is a canonical embedding of  $V$  into  $V^{**}$ . Indeed, pick  $v \in V$ . We define

$$\begin{aligned} \hat{v} : V^* &\rightarrow F \\ \varepsilon &\mapsto \varepsilon(v) \end{aligned}$$

Then this is linear, as

$$\hat{v}(\lambda_1 \varepsilon_1 + \lambda_2 \varepsilon_2) = (\lambda_1 \varepsilon_1 + \lambda_2 \varepsilon_2)(v) = \lambda_1 \varepsilon_1(v) + \lambda_2 \varepsilon_2(v) = \lambda_1 \hat{v}(\varepsilon_1) + \lambda_2 \hat{v}(\varepsilon_2).$$

**Theorem 9.1.** *If  $V$  is a finite dimensional vector space over  $F$ , then the hat map  $v \mapsto \hat{v}$  is an isomorphism.*

In infinite dimension, under certain assumption (e.g. Banach space) we can show that the hat map is injective.

**Proof:** If  $V$  is finite dimensional, then first note that for  $v \in V$ ,  $\hat{v} \in V^{**}$ . We show the hat map is linear: for  $v_1, v_2 \in V$ ,  $\lambda_1, \lambda_2 \in F$  and  $\varepsilon \in V^*$ ,

$$\widehat{\lambda_1 v_1 + \lambda_2 v_2}(\varepsilon) = \varepsilon(\lambda_1 v_1 + \lambda_2 v_2) = \lambda_1 \varepsilon(v_1) + \lambda_2 \varepsilon(v_2) = \lambda_1 \hat{v}_1(\varepsilon) + \lambda_2 \hat{v}_2(\varepsilon).$$

Now we show the hat map is injective. Let  $e \in V \setminus \{0\}$ . Then extend to a basis  $(e, e_2, \dots, e_n)$ . Let  $(\varepsilon, \varepsilon_2, \dots, \varepsilon_n)$  be the dual basis. Then

$$\hat{e}(\varepsilon) = \varepsilon(e) = 1.$$

Hence  $\hat{e} \neq \{0\}$ , so the hat map is injective.

Finally, we show the hat map is an isomorphism. We already know  $\dim V = \dim V^*$ , and as a result  $\dim V^* = \dim V^{**}$ . Thus, since the hat map is injective, it is an isomorphism.

**Lemma 9.3.** *Let  $V$  be a finite dimensional vector space over  $K$ , and let  $U \leq V$ . Then*

$$\hat{U} = U^{\circ\circ}.$$

Hence after identification of  $V$  and  $V^{**}$ , we get

$$U = U^{\circ\circ}.$$

**Proof:** We will show  $U \leq U^{\circ\circ}$ . Indeed, let  $u \in U$ . Then for all  $\varepsilon \in U^\circ$ ,  $\varepsilon(u) = 0$ . So for all  $\varepsilon \in U^\circ$ ,  $\hat{u}(\varepsilon) = \varepsilon(u) = 0$ . Hence  $\hat{u} \in U^{\circ\circ}$ , so  $\hat{U} \subset U^{\circ\circ}$ .

But then we can compute dimension to find

$$\dim U^{\circ\circ} = \dim V - \dim U^\circ = \dim U,$$

proving this lemma.

*Remark.* If  $T \leq V^*$ , then

$$T^\circ = \{v \in V \mid \theta(v) = 0, \forall \theta \in T\}.$$

**Lemma 9.4.** *Let  $V$  be a finite dimensional vector space over  $K$ . Let  $U_1, U_2 \leq V$ . Then,*

- (i)  $(U_1 + U_2)^\circ = U_1^\circ \cap U_2^\circ$ ,
- (ii)  $(U_1 \cap U_2)^\circ = U_1^\circ + U_2^\circ$ .

**Proof:**

- (i) Let  $\theta \in V^*$ , then

$$\begin{aligned} \theta \in (U_1 + U_2)^\circ &\iff \theta(u_1 + u_2) = 0 \iff \theta(u) = 0 \forall u \in U_1 \cup U_2 \\ &\iff \theta \in U_1^\circ \cap U_2^\circ. \end{aligned}$$

Hence  $(U_1 + U_2)^\circ = U_1^\circ \cap U_2^\circ$ .

- (ii) Looking at (i), we can take the annihilator of everything to get

$$(U_1 \cap U_2)^\circ = (U_1^\circ + U_2^\circ)^{\circ\circ} = U_1^\circ + U_2^\circ.$$

## 10 Bilinear Forms

**Definition 10.1.** Let  $U, V$  be vector spaces over  $K$ . Then

$$\phi : U \times V \rightarrow K$$

is a **bilinear form** if it is linear in both components.

**Example 10.1.**

(i) Take  $V \times V^* \rightarrow K$  by  $(v, \theta) \mapsto \theta(v)$ .

(ii) The scalar product on  $U = V = \mathbb{R}^n$  is  $\psi : \mathbb{R}^n \times \mathbb{R}^n \rightarrow \mathbb{R}$  by

$$\left( \begin{pmatrix} x_1 \\ \vdots \\ x_n \end{pmatrix}, \begin{pmatrix} y_1 \\ \vdots \\ y_n \end{pmatrix} \right) \mapsto \sum_{i=1}^n x_i y_i.$$

(iii) If  $U = V = \mathcal{C}([0, 1], \mathbb{R})$ , then we can define

$$\phi(f, g) = \int_0^1 f(t)g(t) \, dt.$$

This can be thought of as an infinite dimensional scalar product.

**Definition 10.2.** Let  $\mathcal{B} = (e_1, \dots, e_m)$  be a basis of  $U$ , and  $\mathcal{C} = (f_1, \dots, f_n)$  be a basis of  $V$ . If  $\phi : U \times V \rightarrow F$  is a bilinear form, then the matrix of  $\phi$  with respect to  $\mathcal{B}$  and  $\mathcal{C}$  is

$$[\phi]_{\mathcal{B}, \mathcal{C}} = (\phi(e_i, f_j)).$$

**Lemma 10.1.**

$$\phi(u, v) = [u]_{\mathcal{B}}^T [\phi]_{\mathcal{B}, \mathcal{C}} [v]_{\mathcal{C}}.$$

**Proof:** Let

$$u = \sum_{i=1}^m \lambda_i e_i, \quad v = \sum_{j=1}^n \mu_j f_j.$$

Since  $\phi$  is a bilinear form,

$$\begin{aligned} \phi(u, v) &= \phi\left(\sum_{i=1}^m \lambda_i e_i, \sum_{j=1}^n \mu_j f_j\right) = \sum_{i=1}^m \sum_{j=1}^n \lambda_i \mu_j \phi(e_i, f_j) \\ &= [u]_{\mathcal{B}}^T [\phi]_{\mathcal{B}, \mathcal{C}} [v]_{\mathcal{C}}. \end{aligned}$$

*Remark.*  $[\phi]_{\mathcal{B}, \mathcal{C}}$  is the only matrix satisfying this property.

**Definition 10.3.**  $\phi : U \times V \rightarrow K$  a bilinear form determines two linear maps:

$$\begin{aligned} \phi_L : U &\rightarrow V^* \\ \phi_L(u) : V &\rightarrow K \\ v &\mapsto \phi(u, v) \\ \phi_R : V &\rightarrow U^* \\ \phi_R(v) : U &\rightarrow K \\ u &\mapsto \phi(u, v) \end{aligned}$$

**Lemma 10.2.** Let  $\mathcal{B} = (e_1, \dots, e_m)$  a basis of  $U$ , and  $\mathcal{B}^* = (\varepsilon_1, \dots, \varepsilon_m)$  a dual basis of  $U^*$ , Similarly, let  $\mathcal{C} = (f_1, \dots, f_n)$  be a basis of  $V$ , and  $\mathcal{C}^* = (\eta_1, \dots, \eta_n)$  a dual basis of  $V^*$ .

Let  $A = [\phi]_{\mathcal{B}, \mathcal{C}}$ . Then,

$$\begin{aligned} [\phi_R]_{\mathcal{C}, \mathcal{B}^*} &= A, \\ [\phi_L]_{\mathcal{B}, \mathcal{C}^*} &= A^T. \end{aligned}$$

**Proof:** We have  $\phi_L(e_i, f_j) = \phi(e_i, f_j) = A_{ij}$ , and so

$$\phi_L(e_i) = \sum A_{ij} \eta_j.$$

Similarly,  $\phi_R(f_j)(e_i) = \phi(e_i, f_j) = A_{ij}$ , so

$$\phi_R(f_j) = \sum A_{ij} \varepsilon_i.$$

This naturally gives our result.

**Definition 10.4.** Let  $\text{Ker } \phi_L$  be the left kernel of  $\phi$ , and  $\text{Ker } \phi_R$  be the right kernel of  $\phi$ .

We say that  $\phi$  is non-degenerate if  $\text{Ker } \phi_L = \{0\}$  and  $\text{Ker } \phi_R = \{0\}$ . Otherwise, we say that  $\phi$  is degenerate.

**Lemma 10.3.** Let  $U, V$  be finite dimensional,  $\mathcal{B}, \mathcal{C}$  bases of  $U$  and  $V$ , and  $\phi : U \times V \rightarrow K$  a bilinear form. Let  $A = [\phi]_{\mathcal{B}, \mathcal{C}}$ .

Then  $\phi$  is non-degenerate if and only if  $A$  is invertible.

**Corollary 10.1.** If  $\phi$  is non-degenerate, then  $\dim U = \dim V$ .

**Proof:**  $\phi$  is non-degenerate if and only if  $\text{Ker } \phi_L = \{0\}$  and  $\text{Ker } \phi_R = \{0\}$ . But this implies  $\text{null}(A^T) = 0$  and  $\text{null}(A) = 0$ , hence by rank-nullity theorem, we must have  $\text{rank}(A^T) = \dim U$ , and  $\text{rank}(A) = \dim V$ . But this gives  $A$  invertible and  $\dim U = \dim V$ .

*Remark.* Taking  $\phi : \mathbb{R}^n \times \mathbb{R}^n \rightarrow \mathbb{R}$  by the scalar product, then  $\phi$  is non-degenerate, as in the standard basis  $\mathcal{B}$ ,

$$[\phi]_{\mathcal{B}, \mathcal{B}} = I_n.$$

**Corollary 10.2.** When  $U$  and  $V$  are finite dimensional, then choosing a non-degenerate bilinear form  $\phi : U \times V \rightarrow K$  is equivalent to choosing an isomorphism  $\phi_L : U \rightarrow V^*$ .

**Definition 10.5.** If  $T \subset U$ , we define

$$T^\perp = \{v \in V \mid \phi(t, v) = 0 \forall t \in T\}.$$

Similarly, if  $S \subset V$ , then

$${}^\perp S = \{u \in U \mid \phi(u, s) = 0 \forall s \in S\}.$$

**Proposition 10.1.** Let  $\mathcal{B}, \mathcal{B}'$  be two bases of  $U$ , and  $P = [\text{id}]_{\mathcal{B}', \mathcal{B}}$ , and  $\mathcal{C}, \mathcal{C}'$  two bases of  $V$ , and  $Q = [\text{id}]_{\mathcal{C}', \mathcal{C}}$ , then if  $\phi : U \times V \rightarrow K$  is a bilinear form, then

$$[\phi]_{\mathcal{B}', \mathcal{C}'} = P^T [\phi]_{\mathcal{B}, \mathcal{C}} Q.$$

**Proof:** We have

$$\phi(u, v) = [u]_{\mathcal{B}}^T [\phi]_{\mathcal{B}, \mathcal{C}} [v]_{\mathcal{C}} = (P[u]_{\mathcal{B}'}^T)^T [\phi]_{\mathcal{B}, \mathcal{C}} (Q[v]_{\mathcal{C}'}) = [u]_{\mathcal{B}'}^T (P^T [\phi]_{\mathcal{B}, \mathcal{C}} Q) [v]_{\mathcal{C}'},$$

which implies  $P^T [\phi]_{\mathcal{B}, \mathcal{C}} Q = [\phi]_{\mathcal{B}', \mathcal{C}'}$ .

**Definition 10.6.** The rank of  $\phi$  ( $\text{rank } \phi$ ) is the rank of any matrix representing  $\phi$ .

This is true as  $\text{rank}(P^T A Q) = \text{rank } A$ , if  $P$  and  $Q$  are invertible.

Note we could have equivalently defined  $\text{rank } \phi = \text{rank } \phi_L = \text{rank } \phi_R$ .

## 11 Determinant and Traces

**Definition 11.1.** If  $A \in M_n(K)$ , we define the trace of  $A$  as

$$\operatorname{tr} A = \sum_{i=1}^n A_{ii}.$$

*Remark.* The map  $M_n(K) \rightarrow K$  by  $A \mapsto \operatorname{tr} A$  is linear.

**Lemma 11.1.**  $\operatorname{tr}(AB) = \operatorname{tr}(BA)$ .

**Proof:**

$$\operatorname{tr}(AB) = \sum_{i=1}^n \left( \sum_{j=1}^n a_{ij} b_{ji} \right) = \sum_{j=1}^n \sum_{i=1}^n b_{ji} a_{ij} = \operatorname{tr}(BA).$$



# Index

- annihilator, 28
- basis, 6
- bidual, 32
- bilinear form, 35
- change of basis matrix, 20
- column rank, 23
- complement, 11
- dimension, 9
- direct sum, 10
- double dual, 32
- dual basis, 27
- dual map, 29
- dual space, 27
- elementary column operation, 24
- elementary matrices, 24
- elementary row operation, 24
- endomorphism, 24
- equivalent matrices, 21
- finite dimension, 5
- Gauss' pivot algorithm, 25
- image, 15
- isomorphism, 14
- kernel, 15
- linear form, 27
- linear independence, 6
- linear map, 13
- matrix, 17
- matrix of bilinear form, 35
- non-degenerate bilinear form, 37
- nullity, 16
- proper subspace, 10
- quotient, 4
- rank, 16
- rank of bilinear form, 38
- rank-nullity theorem, 16
- reflexive space, 32
- row echelon form, 25
- row rank, 23
- scalar product, 35
- similar matrices, 24
- span, 5
- Steinitz exchange lemma, 7
- subspace, 3
- subspace sum, 3
- trace, 39
- vector space, 2