

4. A. S. Smogorzhevskii and E. S. Stolova, Handbook on the Theory of Plane Curves of Third Order [in Russian], Fizmatgiz, Moscow (1961).
5. S. A. Kruglyak and Yu. S. Samoilenko, Funkts. Anal. Prilozhen., 14, No. 1, 60-62 (1980).
6. P. R. Halmos, Matematika [Russian translation], 15, No. 5, 74-83 (1971).
7. V. S. Sunder, Can. J. Math., XL, No. 1, 38-51 (1988).
8. S. A. Kruglyak, in: Representations and Quadratic Forms [in Russian], Kiev (1979), pp. 149-151.
9. P. R. Halmos, Tr. Amer. Math. Soc., 144, 381-389 (1969).

## LOCAL PLÜCKER FORMULAS FOR A SEMISIMPLE LIE GROUP

L. E. Positsel'skii

UDC 517.9

The local Plücker formulas connect the metrics on a holomorphic curve in  $\mathbb{CP}^n$  induced by the Plücker embeddings of the adjoined curves with the corresponding curvatures. The classical (global) Plücker formulas are obtained from the local ones via integration over the curve [3]. Givental' [2] remarked that the vector of curvatures is expressible through the vector of metrics by means of a Cartan matrix of type  $A_n$ . In the present paper we prove a generalization of this fact to the case of an arbitrary Cartan matrix, which was formulated by Givental' as a conjecture.

Let  $G$  be a complex simply connected semisimple Lie group of rank  $n$ ,  $K$  be a compact form of  $G$ , and  $\mathfrak{g}$  be the Lie algebra of  $G$ . Let  $F$  be the set of all Borel subgroups  $B$  of the group  $G$  (the flag variety of  $G$ ).

We shall consider linear bundles  $\mathcal{L}$  on  $F$ , to which there is a lift of the action of  $G$  on  $F$ , i.e., for  $g \in G$ ,  $B \in F$  there is given an isomorphism  $g_*: \mathcal{L}_B \rightarrow \mathcal{L}_{gBg^{-1}}$ . Fix a point  $B_0 \in F$ . The subgroup  $B_0$  acts on the fiber  $\mathcal{L}_{B_0}$  by means of some character  $\omega: B_0 \rightarrow \mathbb{C}^*$ . It is readily seen that the bundle  $\mathcal{L}$  with the action of  $G$  is uniquely recoverable from the character  $\omega$ ; we denote by  $\mathcal{L}(\omega)$  the result of this recovering process. In fact, any bundle  $\mathcal{L}$  admits a unique, up to an automorphism, action of  $G$ , and so the linear bundles over  $F$  are in a bijective correspondence with the characters of the group  $B_0$ . Let  $\omega_i: B_0 \rightarrow \mathbb{C}^*$ ,  $i = 1, \dots, n$ , be fundamental characters, and let  $\mathcal{L}_i = \mathcal{L}(\omega_i)$  be the corresponding bundles; then  $V_i = \Gamma(F, \mathcal{L}_i)^*$  are fundamental representations of  $G$  (the index  $i$  runs through the vertices of the Dynkin diagram).  $V_i$  is irreducible, and therefore on  $\mathbb{P}(V_i)$  there is a uniquely-defined Fubini-Study metric  $FS$ , corresponding to a unique, up to a constant,  $K$ -invariant Hermitian form on  $V_i$ . Let  $f: S \rightarrow F$  be a holomorphic curve, let  $\pi_i: F \rightarrow \mathbb{P}(V_i)$  be the projective morphisms associated with the sheaves  $\mathcal{L}_i$ , and let  $\varphi_i = f^*\pi_i^*(FS)$  be the induced metrics on  $S$ . We shall assume that the metrics  $\varphi_i$  are not identically equal to zero, and since the problem under examination is local, we shall deal with a neighborhood of a point where all of them are different from zero. Let  $\theta_i$  be the curvatures of the metrics  $\varphi_i$ .

Let us define a natural  $n$ -dimensional distribution  $\mathcal{N}$  on the flag variety  $F$ . The tangent space  $T_B F$  is isomorphic to  $\mathfrak{g}/\mathfrak{b}$ , where  $\mathfrak{b}$  is the Lie algebra of the group  $B$ . Set

$$\mathcal{N}(B) = \{x \in \mathfrak{g} : [x, [\mathfrak{b}, \mathfrak{b}]] \subset \mathfrak{b}\} / \mathfrak{b}.$$

**THEOREM 1.** Suppose the curve  $S$  is tangent to the distribution  $\mathcal{N}$ . Let  $\varphi$  be the vector of  $(1, 1)$ -forms composed of the metrics  $\varphi_i$  on  $S$ , and let  $\theta$  be the vector of curvatures. Then  $\theta = A\varphi$ , where  $A$  is the Cartan matrix of the group  $G$ .

The connection with the classical case is as follows. Let  $K = \mathrm{SU}(n+1)$ . Then  $F$  is the flag variety in  $\mathbb{CP}^{n+1}$ . The morphism  $\pi_i$  is the composition of the projection onto the Grassmannian which sends a flag into its  $i$ -plane, and the Plücker embedding;  $\pi_i(S)$  is a curve in  $\mathbb{CP}^n$ . The condition that  $S$  belongs to the distribution  $\mathcal{N}$  means that the preimage

---

V. M. Lomonosov State University, Moscow. Translated from Funktsional'nyi Analiz i Ego Prilozheniya, Vol. 25, No. 4, pp. 74-76, October-December, 1991. Original article submitted May 10, 1990.

in  $S$  of a point  $x \in \pi_1(S)$  is the moving flag of the curve  $\pi_1(S)$  at the point  $x$ , i.e.,  $\pi_1(S)$  are Plücker-embedded adjoined curves for  $\pi_1(S)$ . The assertion of the theorem is identical with the local Plücker formulas for the curve  $\pi_1(S)$ , given in [3].

Proof. A subgroup  $P \subset G$  is called a parabolic subgroup [1, 4] if it contains a Borel subgroup. The subgroups  $P_I(B)$  containing  $B$  correspond to subsets  $I$  of the vertex set of the Dynkin diagram, and the type  $I$  of a parabolic subgroup does not depend on the choice of the Borel subgroup contained in it. Let  $F_I \simeq G/P_I(B)$  be the variety of parabolic subgroups of type  $I$ . Instead of the subsets  $\{i\}$  and  $\{1, \dots, \hat{i}, \dots, n\}$  we shall write the indices  $i$  and  $i'$ , respectively.

I. The map  $\pi_i$  factors as the composition of the canonical projection  $\rho_i: F \rightarrow F_{i'}$ ,  $\rho_i(B) = P_{i'}(B)$  and an embedding  $\tau_i: F_{i'} \rightarrow \mathbb{P}(V_i)$ .

Let us show that the map  $\pi_i$  is constant on the fibers of  $\rho_i$ , i.e., the bundle  $\mathcal{L}_i$  is trivial along these fibers. The fibers of  $\rho_i$  are the sets of all Borel subgroups that are contained in a fixed subgroup of the form  $P_{i'}(B)$ , or, equivalently, all Borel subgroups in the semisimple subgroup  $G_{i'} \subset G$  generated by all simple roots except the  $i$ -th. Now  $\mathcal{L}(\omega_i)|_{F(G_{i'})} = \mathcal{L}(\omega_i|_{B \cap G_{i'}})$ , but  $\omega_i|_{B \cap G_{i'}} = 1$  by definition. The morphisms  $\tau_i$  are embeddings, since they are morphisms of  $G$ -spaces, and  $P_{i'}(B)$  is a maximal proper subgroup of  $G$ .

II. Let  $\mathcal{N}_i$  be the one-dimensional distribution on  $F$  defined as  $\mathcal{N}_i(B) = \mathfrak{p}_i(B)/\mathfrak{h}$ , where  $\mathfrak{p}_i(B)$  is the Lie algebra of the group  $P_i(B)$ . Then, as is readily seen,  $\mathcal{N}(B) = \bigoplus_{i=1}^n \mathcal{N}_i(B)$ .

III.  $d\pi_i(\mathcal{N}_j) = 0$  for  $i \neq j$ . Indeed,  $P_j(B) \subset P_{i'}(B)$ .

IV. Let  $\theta(\mathcal{L})$  be the curvature of a  $K$ -invariant metric  $\|\cdot\|$  in an arbitrary bundle  $\mathcal{L}$  over  $F$  equipped with an action of  $G$ . Let  $s$  be a local section of  $\mathcal{L}$ . Then one has the following formula [3]:

$$\theta(\mathcal{L}) = -\partial\bar{\partial} \ln \|s\|,$$

from which it follows, in particular, that the mapping  $\mathcal{L} \mapsto \theta(\mathcal{L})$  takes tensor products into sums.

V.  $\varphi_i = f^*\theta(\mathcal{L}_i)$ . Let us show that  $\pi_i^*(FS) = \theta(\mathcal{L}_i)$ . Let  $s^*$  be a local section of  $\mathcal{L}_i^*$ . Then  $\theta(\mathcal{L}_i) = \partial\bar{\partial} \ln \|s^*\|$ . The section  $s^*$  yields a local lift  $\tilde{\pi}_i: U \rightarrow V_i$ ,  $U \subset F$ , by the rule  $\tilde{\pi}_i(u)(t) = \langle s^*(u), t(u) \rangle$ ,  $t \in \Gamma(F, \mathcal{L}_i)$ ,  $u \in U$ . Therefore, by definition,  $\pi_i^*(FS) = \partial\bar{\partial} \ln \|\tilde{\pi}_i\| = \partial\bar{\partial} \ln \|s^*\|$ .

VI.  $\theta_i = f^*\theta(\mathcal{N}_i)$ , where the  $\mathcal{N}_i$  are regarded as abstract bundles. In fact, by III, the metric  $\varphi_i$  on  $S$  is lifted by means of the projection on  $\mathcal{N}_i$ , i.e., is induced by the isomorphism  $\tau_S \simeq f^*\mathcal{N}_i$ .

VII.  $(\mathcal{N}) = (\mathcal{L})^A$ , i.e.,  $\mathcal{N}_i \simeq \bigotimes_j \mathcal{L}_j^{\otimes a_{ij}}$ . The bundles  $\mathcal{N}_i$  correspond, as is readily seen, to the characters effecting the  $\text{Ad}^{-1}$ -action of  $B$  on the lines  $N_i \subset \mathfrak{g}/\mathfrak{b}$ , i.e., to the characters  $\alpha_i$  corresponding to the simple roots, and  $(\alpha) = (\omega)^A$ .

The needed assertion follows upon combining the results IV-VII.

Note. In the case  $K = \text{Sp}(2n)$ , i.e., the case of a Cartan matrix of type  $C_n$ , the theorem is proved in [2] by means of reduction to the classical case.

The author thanks A. Yu. Vaintrob for posing the problem and for unceasing help and support, and to A. B. Givental', who suggested an essential simplification of the proof.

#### LITERATURE CITED

1. A. Borel, Linear Algebraic Groups, W. A. Benjamin, New York (1969).
2. A. B. Givental', "Plücker formulas and Cartan matrices," Usp. Mat. Nauk, 44, No. 3, 155-156 (1989).
3. Ph. A. Griffiths and J. Harris, Principles of Algebraic Geometry, Wiley-Interscience, New York (1978).
4. J. E. Humphreys, Linear Algebraic Groups, Graduate Texts in Math., No. 21, Springer, New York (1975).