



# Communications in Partial Differential Equations

Publication details, including instructions for authors and subscription information:  
<http://www.tandfonline.com/loi/lpde20>

## Uniform estimates and blow-up behavior for solutions of $-\delta(u)=v(x)e^u$ in two dimensions

Haïm Brezis <sup>a</sup> & Frank Merle <sup>b</sup>

<sup>a</sup> Université Paris VI and Rutgers University,

<sup>b</sup> CNRS, Université Paris VI,

Version of record first published: 28 Jul 2009.

To cite this article: Haïm Brezis & Frank Merle (1991): Uniform estimates and blow-up behavior for solutions of  $-\delta(u)=v(x)e^u$  in two dimensions, Communications in Partial Differential Equations, 16:8-9, 1223-1253

To link to this article: <http://dx.doi.org/10.1080/03605309108820797>

PLEASE SCROLL DOWN FOR ARTICLE

Full terms and conditions of use: <http://www.tandfonline.com/page/terms-and-conditions>

This article may be used for research, teaching, and private study purposes. Any substantial or systematic reproduction, redistribution, reselling, loan, sub-licensing, systematic supply, or distribution in any form to anyone is expressly forbidden.

The publisher does not give any warranty express or implied or make any representation that the contents will be complete or accurate or up to date. The accuracy of any instructions, formulae, and drug doses should be independently verified with primary sources. The publisher shall not be liable for any loss, actions, claims, proceedings, demand, or costs or damages whatsoever or howsoever caused arising directly or indirectly in connection with or arising out of the use of this material.

UNIFORM ESTIMATES AND BLOW-UP BEHAVIOR FOR SOLUTIONS  
OF  $-\Delta u = V(x)e^u$  IN TWO DIMENSIONS

Haïm BREZIS  
Université Paris VI  
and  
Rutgers University

Frank MERLE  
CNRS, Université Paris VI

Introduction

In this paper we deal with the equation

$$(*) \quad \begin{cases} -\Delta u = V(x)e^u & \text{in } \Omega \subset \mathbb{R}^2, \\ u = 0 & \text{on } \partial\Omega, \end{cases}$$

where  $\Omega$  is a bounded domain (except in Section II.3) and  $V(x)$  is a given function in  $L^p(\Omega)$  for some  $1 < p \leq \infty$ . We assume that  $u \in L^1(\Omega)$  and  $e^u \in L^{p'}(\Omega)$  (where  $p'$  is the conjugate exponent of  $p$ ) so that  $(*)$  has a meaning in the sense of distributions.

A first question is whether one can conclude that  $u \in L^\infty(\Omega)$ . As we will see in Section II the answer is positive. Next we turn, in Section III, to a more delicate issue, namely the question of uniform estimates. Suppose we have a sequence  $(u_n)$  of solutions of

$$(**) \quad \begin{cases} -\Delta u_n = V_n(x)e^{u_n} & \text{in } \Omega, \\ u_n = 0 & \text{on } \partial\Omega, \end{cases}$$

with

$$\|V_n\|_{L^p} \leq C_1$$

and

$$\|e^{u_n}\|_{L^{p'}} \leq C_2.$$

Can one conclude that

$$\|u_n\|_{L^\infty} \leq C_3$$

where  $C_3$  depends only on  $C_1, C_2$  and  $\Omega$ ? We prove that the answer is positive under a smallness condition, namely  $C_1 C_2 < 4\pi/p'$  (see Corollary 3). The answer is also positive under a domination condition, namely  $|V_n| \leq W$  for a fixed  $W \in L^p(\Omega), 1 < p < \infty$  (and then  $C_3$  depends also on  $W$ , see Corollary 5).

A deeper result (see Corollary 6) asserts that if  $V_n \geq 0$  then  $(u_n)$  is bounded in  $L^\infty_{loc}(\Omega)$ , i.e. for every compact subset  $K$  of  $\Omega$  we have

$$\|u_n\|_{L^\infty(K)} \leq C_3$$

where  $C_3$  depends only on  $C_1, C_2$  and  $K$ . Surprisingly such an estimate does not hold up to the boundary. Given any  $1 < p \leq \infty$  we construct in Example 6 (Section III.3) sequences  $(u_n)$  and  $(V_n)$  satisfying (\*\*) with  $V_n \geq 0$

$$\|V_n\|_{L^p} \leq C_1$$

$$\|e^{u_n}\|_{L^{p'}} \leq C_2$$

and  $\|u_n\|_{L^\infty} \rightarrow +\infty$ .

A corollary of our methods also yields the following. Suppose  $u_n$  satisfies

$$-\Delta u_n = V_n e^{u_n} \text{ in } \Omega$$

with

$$0 < a \leq V_n \leq b < \infty$$

and

$$\inf_\Omega u_n \geq -M > -\infty$$

(here no boundary condition is imposed). Then for every compact subset  $K$  of  $\Omega$ ,  $\sup_K u_n$  can be estimated just in terms of  $a, b, M, K$  and  $\Omega$  (see Corollary 8).

Finally we turn to the general case where no boundary condition is imposed and  $(u_n)$  is not bounded below. More precisely let  $(u_n)$  be a sequence of solutions of

$$-\Delta u_n = V_n e^{u_n} \text{ in } \Omega$$

with

$$V_n \geq 0 \text{ in } \Omega, \|V_n\|_{L^p} \leq C_1 \text{ and } \|e^{u_n}\|_{L^{p'}} \leq C_2,$$

for some  $1 < p \leq \infty$ .

Then we have the following alternative (see Theorem 3):

either

$$(i) \quad (u_n) \text{ is bounded in } L_{loc}^\infty(\Omega)$$

or

$$(ii) \quad u_n \rightarrow -\infty \text{ uniformly on compact subsets of } \Omega$$

or

(iii) there is a finite nonempty set  $S$  such that  $u_n \rightarrow -\infty$  uniformly on compact subsets of  $\Omega \setminus S$  and  $u_n \rightarrow +\infty$  on  $S$  (in a sense to be precised later).

In this case  $V_n e^{u_n}$  converges to a finite sum of Dirac masses  $\sum \alpha_i \delta_{a_i}$  with coefficients  $\alpha_i \geq 4\pi/p'$ .

Such behavior is well illustrated by the sequence

$$u_n(x) = \log \frac{8n^2}{(1+n^2|x|^2)^2}$$

which satisfies  $-\Delta u_n = e^{u_n}$ ,  $\|e^{u_n}\|_{L^1} \leq C$ ,  $u_n(x) \rightarrow -\infty$  for all  $x \neq 0$  and

$u_n(0) \rightarrow +\infty$ . Here  $e^{u_n}$  converges to  $8\pi\delta_0$ .

We thank Congming Li for raising questions which led us to Theorem 2 and Corollary 3 (Theorem 2 is used in [3]). Some of our results (in particular Corollary 4 and Theorem 4) are connected to earlier works of Nagasaki and Suzuki (see [6] and [7]) who consider mostly the case where the  $V_n$ 's are constants. A. Chang

and P. Yang [2] have also studied blow-up sequences for related equations on  $S^2$  (see e.g. their Concentration Lemma). However their approach involves  $H^1$  norms and is quite different from ours.

In a forthcoming work we shall consider similar issues for the equation  $-\Delta u = V(x)u^p$  in  $\Omega \subset \mathbb{R}^N$ ,  $N \geq 3$ . The plan of the paper is the following:

#### Introduction

##### I. A basic inequality

##### II. $L^\infty$ -boundedness for a single solution of $-\Delta u = Ve^u$

###### II.1. The case of a bounded domain

###### II.2. Some variants and counterexamples

###### II.3. The case $\Omega = \mathbb{R}^2$

##### III. Uniform $L^\infty$ bounds and blow-up behavior for solutions of $-\Delta u = Ve^u$

###### III.1. Some easy cases

###### III.2. The main results

###### III.3. Variants and counterexamples.

##### I. A basic inequality

Assume  $\Omega \subset \mathbb{R}^2$  is a bounded domain and let  $u$  be a solution of

$$(1) \quad \begin{cases} -\Delta u = f(x) & \text{in } \Omega, \\ u = 0 & \text{on } \partial\Omega, \end{cases}$$

with  $f \in L^1(\Omega)$ . Set  $\|f\|_1 = \int_{\Omega} |f(x)| dx$ .

**Theorem 1.** For every  $\delta \in (0, 4\pi)$  we have

$$(2) \quad \int_{\Omega} \exp \left[ \frac{(4\pi - \delta)|u(x)|}{\|f\|_1} \right] dx \leq \frac{4\pi^2}{\delta} (\text{diam } \Omega)^2.$$

**Proof.** Let  $R = \frac{1}{2} \text{diam } \Omega$  so that  $\Omega \subset B_R$  for some ball of radius  $R$ . Extend  $f$  to be zero outside  $\Omega$  and set, for  $x \in \mathbb{R}^2$ ,

$$\bar{u}(x) = \frac{1}{2\pi} \int_{B_R} \log\left(\frac{2R}{|x-y|}\right) |f(y)| dy$$

so that

$$-\Delta \bar{u} = |f| \quad \text{on } \mathbb{R}^2.$$

Note that  $\bar{u}(x) \geq 0$  for  $x \in B_R$  since  $\frac{2R}{|x-y|} \geq 1 \quad \forall x, y \in B_R$ . It follows from the maximum principle that  $|u| \leq \bar{u}$  on  $\Omega$  and thus

$$(3) \quad \int_{\Omega} \exp\left[\frac{(4\pi-\delta)|u(x)|}{\|f\|_1}\right] dx \leq \int_{B_R} \exp\left[\frac{(4\pi-\delta)\bar{u}(x)}{\|f\|_1}\right] dx.$$

We now estimate the right-hand side of (3) using Jensen's inequality

$$F\left(\int w(y)\varphi(y)dy\right) \leq \int w(y)F(\varphi(y))dy$$

with  $F(t) = \exp t$ ,  $w(y) = \frac{|f(y)|}{\|f\|_1}$  and  $\varphi(y) = \frac{(4\pi-\delta)}{2\pi} \log\left(\frac{2R}{|x-y|}\right)$ . We obtain

$$\begin{aligned} \int_{B_R} \exp\left[\frac{(4\pi-\delta)\bar{u}(x)}{\|f\|_1}\right] dx &\leq \int_{B_R} dx \int_{B_R} \left(\frac{2R}{|x-y|}\right)^{2-\frac{\delta}{2\pi}} \frac{|f(y)|}{\|f\|_1} dy \\ &= \int_{B_R} \frac{|f(y)|}{\|f\|_1} \left[ \int_{B_R} \left(\frac{2R}{|x-y|}\right)^{2-\frac{\delta}{2\pi}} dx \right] dy. \end{aligned}$$

But, for  $y \in B_R$ , we have

$$\int_{B_R} \left(\frac{2R}{|x-y|}\right)^{2-\frac{\delta}{2\pi}} dx \leq \int_{B_R} \left(\frac{2R}{|x|}\right)^{2-\frac{\delta}{2\pi}} dx = \frac{4\pi^2}{\delta} (\text{diam } \Omega)^2$$

and the estimate (2) follows.

A simple consequence of Theorem 1 is

**Corollary 1.** Let  $u$  be a solution of (1) with  $f \in L^1(\Omega)$ . Then for every constant  $k > 0$

$$e^{k|u|} \in L^1(\Omega).$$

Proof. Let  $0 < \epsilon < 1/k$ . We may split  $f$  as  $f = f_1 + f_2$  with  $\|f_1\|_1 < \epsilon$  and  $f_2 \in L^\infty(\Omega)$ . Write  $u = u_1 + u_2$  where  $u_i$  are the solutions of

$$\begin{cases} -\Delta u_i = f_i & \text{in } \Omega, \\ u_i = 0 & \text{on } \partial\Omega. \end{cases}$$

Choosing, for example,  $\delta = (4\pi-1)$  in Theorem 1 we find  $\int_{\Omega} \exp\left[\frac{|u_1(x)|}{\|f_1\|_1}\right] < \infty$  and thus  $\int_{\Omega} \exp[k|u_1|] < \infty$ . The conclusion follows since  $|u| \leq |u_1| + |u_2|$  and  $u_2 \in L^\infty(\Omega)$ .

Remark 1. The conclusion of Theorem 1 could also be deduced from BMO estimates and the John-Nirenberg inequality [4].

Remark 2. There is a local form of Corollary 1, namely if  $u \in L^1_{loc}(\Omega)$  and  $\Delta u \in L^1_{loc}(\Omega)$ , then for every  $k > 0$ ,  $e^{k|u|} \in L^1_{loc}(\Omega)$ . [Here we use the well-known fact that  $u \in L^1_{loc}(\Omega)$  and  $\Delta u \in L^1_{loc}(\Omega)$  imply  $\nabla u \in L^1_{loc}(\Omega)$ .]

Remark 3. In Corollary 1,  $e^{k|u|} \in L^1$  but  $\|e^{k|u|}\|_1$  can not be estimated in terms of  $k$  and  $\|f\|_1$ . For example, we may have a sequence  $(f_n)$  such that  $\|f_n\|_1 \leq 1$ ,  $f_n \rightarrow \delta_{x_0}$  and then  $u_n \rightarrow u$  with  $u(x) \simeq \frac{1}{2\pi} \log \frac{1}{|x-x_0|}$  as  $x \rightarrow x_0$  so that  $\int e^{k|u|} = \infty$  for  $k \geq 4\pi$ .

## II. $L^\infty$ -boundedness for a single solution of $-\Delta u = Ve^u$ .

### II.1. The case of a bounded domain.

Let  $u$  satisfy the nonlinear equation

$$(4) \quad \begin{cases} -\Delta u = V(x)e^u & \text{in } \Omega, \\ u = 0 & \text{on } \partial\Omega, \end{cases}$$

where  $\Omega$  is a bounded domain in  $\mathbb{R}^2$  and  $V(x)$  is a given function on  $\Omega$ .

**Corollary 2.** Suppose  $u$  is a solution of (4) with  $V \in L^p(\Omega)$  and  $e^u \in L^{p'}(\Omega)$  for some  $1 < p \leq \infty$ . Then  $u \in L^\infty(\Omega)$ .

**Proof.** By Corollary 1 we know that  $e^{ku} \in L^1(\Omega) \forall k > 0$ , i.e.,  $e^u \in L^r(\Omega) \forall r < \infty$ . It follows that  $Ve^u \in L^{p-\delta} \forall \delta > 0$  if  $p < \infty$ , and  $Ve^u \in L^r(\Omega) \forall r < \infty$  if  $p = \infty$ . Standard elliptic estimates imply that  $u \in L^\infty(\Omega)$ .

**Remark 4.** The conclusion of Corollary 2 still holds for a solution  $u$  of

$$\begin{cases} -\Delta u = V(x)e^u + f(x) & \text{in } \Omega, \\ u = g & \text{on } \partial\Omega, \end{cases}$$

with  $g \in L^\infty(\partial\Omega)$  and  $f \in L^q(\Omega)$  for some  $q > 1$ . Indeed let  $w$  be the solution of

$$\begin{cases} -\Delta w = f & \text{in } \Omega, \\ w = g & \text{on } \partial\Omega, \end{cases}$$

so that  $w \in L^\infty(\Omega)$ . The function  $\tilde{u} = u - w$  satisfies

$$\begin{cases} -\Delta \tilde{u} = (Ve^w)e^{\tilde{u}} & \text{in } \Omega, \\ \tilde{u} = 0 & \text{on } \partial\Omega, \end{cases}$$

and we are reduced to the assumptions of Corollary 2.

**Remark 5.** There is a local version of Corollary 2, namely if  $u \in L^1_{\text{loc}}(\Omega)$  satisfies

$$-\Delta u = Ve^u$$

with  $V \in L^p_{\text{loc}}(\Omega)$  and  $e^u \in L^{p'}_{\text{loc}}(\Omega)$  for some  $1 < p \leq \infty$ , then  $u \in L^\infty_{\text{loc}}(\Omega)$ .

This follows easily from Remark 2.

## II.2. Some variants and counterexamples.

1. The conclusion of Corollary 2 fails when  $p = 1$  (we may only say that  $u^+ \in L^\infty(\Omega)$ ). Here is an example:



**Example 1.** Let  $0 < a < 1$ . The function

$$u = -a \log(\log \frac{e}{r}) \quad \text{with } r = |x| \quad \text{satisfies}$$

$$(5) \quad \begin{cases} -\Delta u = Ve^u & \text{in } \Omega = B_1 \\ u = 0 & \text{on } \partial\Omega \end{cases}$$

with  $V = -\frac{a}{r^2(\log \frac{e}{r})^{2-a}}$ . Note that  $V \in L^1(\Omega)$ ,  $e^u \in L^\infty(\Omega)$  and nevertheless  $u \notin L^\infty(\Omega)$  since  $u(x) \rightarrow -\infty$  as  $x \rightarrow 0$ . The same function  $u$  with  $a < 0$  provides an example where  $u$  satisfies (5) with  $V \in L^1(\Omega)$ ,  $Ve^u \in L^1(\Omega)$  and nevertheless  $u^+ \notin L^\infty(\Omega)$  since  $u(x) \rightarrow +\infty$  as  $x \rightarrow 0$ .

2. The function  $e^u$  is in some sense the "critical nonlinearity" for which a statement such as Corollary 2 holds. Suppose, for example, that  $u$  satisfies

$$\begin{cases} -\Delta u = V(x)e^{u^\alpha} & \text{in } \Omega \\ u = 0 & \text{on } \partial\Omega \end{cases}$$

with  $u \geq 0$ ,  $\alpha > 1$ ,  $V \in L^p(\Omega)$  and  $e^{u^\alpha} \in L^{p'}(\Omega)$ ,  $1 < p \leq \infty$ . In general, we may not infer that  $u \in L^\infty(\Omega)$ .

**Example 2.** Consider first the case  $p = \infty$ . Fix  $1 < \gamma < 2 - (1/\alpha)$ . In  $\Omega = B_1$  set

$$u(x) = |\log(r^2(\log \frac{e}{r})^\gamma)|^{1/\alpha}.$$

For  $r$  small we have

$$e^{u^\alpha} = \frac{1}{r^{2(\log \frac{e}{r})^\gamma}}$$

and therefore  $e^{u^\alpha} \in L^1(\Omega)$ . On the other hand  $u$  satisfies  $-\Delta u = Ve^{u^\alpha}$  where  $V$  is defined by  $V = (-\Delta u)e^{-u^\alpha}$ . An easy computation shows that

$$V \sim |\log r|^{\gamma-2+(1/\alpha)} \quad \text{as } r \rightarrow 0$$

and hence  $V \in L^\infty(\Omega)$ . Nevertheless  $u \notin L^\infty(\Omega)$ .

When  $1 < p < \infty$  we may use the function  $u$  above and write

$$-\Delta u = (Ve^{\frac{1}{p}u^\alpha})e^{\frac{1}{p'}u^\alpha}$$

The function  $\tilde{u} = (p')^{-1/\alpha}u$  satisfies  $-\Delta \tilde{u} = \tilde{V}e^{\tilde{u}^\alpha}$  with  $\tilde{V} = (p')^{-1/\alpha}Ve^{\frac{1}{p}u^\alpha}$  so that  $\tilde{V} \in L^p(\Omega)$  and  $e^{\tilde{u}^\alpha} \in L^{p'}(\Omega)$ .

3. There is a version of Corollary 2 for subsolutions. Assume  $u$  satisfies

$$\begin{aligned} -\Delta u &\leq V(x)e^u \quad \text{in } \Omega, \\ u &\leq 0 \quad \text{on } \partial\Omega, \end{aligned}$$

with  $V \in L^p(\Omega)$  and  $e^u \in L^{p'}(\Omega)$  for some  $1 < p \leq \infty$ . Then  $u^+ \in L^\infty(\Omega)$ .

### II.3. The case $\Omega = \mathbb{R}^2$ .

The main result is the following.

Theorem 2. Suppose  $u \in L^1_{\text{loc}}(\mathbb{R}^2)$  satisfies

$$-\Delta u = V(x)e^u \quad \text{in } \mathbb{R}^2$$

with  $V \in L^p(\mathbb{R}^2)$  and  $e^u \in L^{p'}(\mathbb{R}^2)$  for some  $1 < p \leq \infty$ . Then  $u \in L^\infty(\mathbb{R}^2)$ .

Proof. Fix  $0 < \epsilon < 1/p'$  and split  $Ve^u$  as  $Ve^u = f_1 + f_2$  with  $\|f_1\|_{L^1(\mathbb{R}^2)} < \epsilon$  and  $f_2 \in L^\infty(\mathbb{R}^2)$ . Let  $B_r$  be the ball of radius  $r$  centered at  $x_0$ . We denote by  $C$  various constants independent of  $x_0$  (but possibly depending on  $\epsilon$ ). Let  $u_1$  be the solution of

$$\begin{cases} -\Delta u_1 = f_1 & \text{in } B_1, \\ u_1 = 0 & \text{on } \partial B_1. \end{cases}$$

By Theorem 1 (applied with  $\delta = 4\pi - 1$ ) we have

$$\int_{B_1} \exp\left[\frac{1}{\epsilon} |u_1|\right] \leq C$$

and in particular  $\|u_1\|_{L^1(B_1)} \leq C$ . We also have  $\|u_2\|_{L^\infty(B_1)} \leq C$ . Let

$u_3 = u - u_1 - u_2$  so that  $\Delta u_3 = 0$  on  $B_1$ . The mean value theorem for harmonic functions implies that

$$(6) \quad \|u_3^+\|_{L^\infty(B_{1/2})} \leq C \|u_3^+\|_{L^1(B_1)}.$$

On the other hand we have

$$u_3^+ \leq u^+ + |u_1| + |u_2|$$

and since

$$p' \int_{\mathbb{R}^2} u^+ \leq \int_{\mathbb{R}^2} e^{p'u} \leq C$$

we see that  $\|u_3^+\|_{L^1(B_1)} \leq C$ . Combining this with (6) we find that

$$\|u_3^+\|_{L^\infty(B_{1/2})} \leq C. \text{ Finally we write}$$

$$(7) \quad -\Delta u = Ve^u = (Ve^{u_1})e^{u_2+u_3} = g$$

with  $\|g\|_{L^{1+\delta}(B_{1/2})} \leq C$  for some  $\delta > 0$  (since  $e^{u_2+u_3} \in L^\infty(B_{1/2})$ ),

$V \in L^p(B_1)$  and  $e^{u_1} \in L^{1/\epsilon}(B_1)$  with  $1/\epsilon > p'$ ). Using once more the mean value theorem and standard elliptic estimates we deduce from (7) that

$$\|u^+\|_{L^\infty(B_{1/4})} \leq C \|u^+\|_{L^1(B_{1/2})} + C \|g\|_{L^{1+\delta}(B_{1/2})} \leq C.$$

Since  $C$  is independent of  $x_0$  we conclude that  $u^+ \in L^\infty(\mathbb{R}^2)$ .

### III. Uniform $L^\infty$ bounds and blow-up behavior for solutions of $-\Delta u = V(x)e^u$ .

In this section we consider a sequence  $(u_n)$  of solutions of

$$(8) \quad -\Delta u_n = V_n(x)e^{u_n} \quad \text{in } \Omega$$

where  $\Omega$  is a bounded domain in  $\mathbb{R}^2$ . We seek a uniform bound for  $\|u_n\|_{L^\infty}$

(resp.  $\|u_n\|_{L^\infty_{loc}}$ ) under various assumptions. We start with:

### III.1. Some easy cases

There are two different kinds of assumptions which lead easily to uniform bounds:

- a) Smallness assumption.
- b) Uniform domination.

#### a) Smallness assumption

Corollary 3. Assume  $(u_n)$  is a sequence of solutions of (8) with  $u_n = 0$  on  $\partial\Omega$ , such that

$$(9) \quad \|V_n\|_{L^p} \leq C \text{ for some } 1 < p \leq \infty$$

and

$$(10) \quad \int_{\Omega} |V_n| e^{u_n} \leq \epsilon_0 < 4\pi/p' \quad \forall n.$$

Then  $\|u_n\|_{L^\infty} \leq C$ .

Proof. Fix  $\delta > 0$  such that  $4\pi - \delta > \epsilon_0(p' + \delta)$ . By Theorem 1 we have

$$\int_{\Omega} e^{(p' + \delta)u_n} \leq C.$$

Therefore  $e^{u_n}$  is bounded in  $L^{p' + \delta}(\Omega)$  and so  $V_n e^{u_n}$  is bounded in  $L^q(\Omega)$  for some  $q > 1$ . Hence  $u_n$  is bounded in  $L^\infty(\Omega)$ .

Remark 6. The smallness condition (10) is sharp. Given any  $1 < p \leq \infty$  one can construct a sequence  $(u_n)$  of solutions of (8) satisfying (9) and

$$(11) \quad \int_{\Omega} |V_n| e^{u_n} = 4\pi/p'$$

such that  $\|u_n\|_{L^\infty} \rightarrow \infty$ .

**Example 3.** Set

$$f_n(x) = \begin{cases} \frac{1}{p} n^2 & \text{if } |x| < 1/n, \\ 0 & \text{otherwise} \end{cases}.$$

Let  $u_n$  be the solution of

$$\begin{cases} -\Delta u_n = f_n & \text{in } B_1, \\ u_n = 0 & \text{on } \partial B_1. \end{cases}$$

Note that  $u_n$  satisfies (8) with  $V_n$  being defined by  $V_n = f_n e^{-u_n}$ . An easy computation shows that (9) and (11) hold. Moreover  $\|u_n\|_{L^\infty} = u_n(0) = \frac{1}{p}(2 \log n + 1)$ .

Here is a variant of Corollary 3 where no boundary condition is imposed.

**Corollary 4.** Assume  $(u_n)$  is a sequence of solutions of (8) such that, for some  $1 < p \leq \infty$ ,

$$(12) \quad \|V_n\|_{L^p} \leq C_1,$$

$$(13) \quad \|u_n^+\|_{L^1} \leq C_2$$

and

$$(14) \quad \int_{\Omega} |V_n| e^{u_n} \leq \epsilon_0 < 4\pi/p'.$$

Then  $(u_n^+)$  is bounded in  $L_{loc}^\infty(\Omega)$ .

**Proof.** Without loss of generality we may assume that  $\Omega = B_R$ . Split  $u_n$  as  $u_n = u_{1n} + u_{2n}$  where  $u_{1n}$  is the solution of

$$(15) \quad \begin{cases} -\Delta u_{1n} = V_n e^{u_n} & \text{in } \Omega, \\ u_{1n} = 0 & \text{on } \partial\Omega; \end{cases}$$

so that  $\Delta u_{2n} = 0$  in  $\Omega$ . By the mean value theorem for harmonic functions we have

$$\|u_{2n}^+\|_{L^\infty(B_{R/2})} \leq C \|u_{2n}^+\|_{L^1(B_R)} \leq C \left[ \|u_n^+\|_{L^1(B_R)} + \|u_{1n}\|_{L^1(B_R)} \right] \leq C.$$

Using (15), the smallness condition (14) and Theorem 1 we see that  $(e^{u_{1n}})$  is bounded in  $L^{p'+\delta}(B_R)$  for some  $\delta > 0$ . Therefore  $(V_n e^{u_n})$  is bounded in  $L^q(B_{R/2})$  for some  $q > 1$ . Using (15) once more we see that  $(u_{1n})$  is bounded in  $L^\infty(B_{R/4})$ . Therefore  $(u_n)$  is bounded in  $L^\infty(B_{R/4})$ .

b) Uniform domination

Corollary 5. Assume  $(u_n)$  is a sequence of solutions of (8) with  $u_n = 0$  on  $\partial\Omega$ , satisfying, for some  $1 < p < \infty$ ,

$$(16) \quad \|e^{u_n}\|_{L^{p'}} \leq C$$

and one of the following conditions:

either

$$(17) \quad |V_n(x)| \leq W(x) \quad \forall n, \text{ with } W \in L^p(\Omega)$$

or

$$(18) \quad V_n \rightarrow V \text{ in } L^p(\Omega).$$

Then  $\|u_n\|_{L^\infty} \leq C$ .

Proof. Assume first that (17) holds. For every  $\epsilon > 0$  we have

$$|V_n| e^{u_n} \leq W e^{u_n} \leq \epsilon e^{p' u_n} + \frac{1}{\epsilon^{1/(p-1)}} W^p.$$

By (16) we may fix  $\epsilon > 0$  small enough so that

$$(19) \quad \epsilon \int_{\Omega} e^{p' u_n} \leq \alpha < 4\pi/p' \quad \forall n.$$

We have  $|u_n| \leq u_{1n} + u_2$  where  $u_{1n}$  is the solution of

$$\begin{cases} -\Delta u_{1n} = \epsilon e^{p'u_n} & \text{in } \Omega, \\ u_{1n} = 0 & \text{on } \partial\Omega \end{cases}$$

and  $u_2$  is the solution of

$$\begin{cases} -\Delta u_2 = \frac{1}{\epsilon^{1/(p-1)}} W^p & \text{in } \Omega, \\ u_2 = 0 & \text{on } \partial\Omega. \end{cases}$$

By Theorem 1 and (19) we see that  $e^{u_{1n}}$  is bounded in  $L^{p'+\delta}(\Omega)$  for some  $\delta > 0$  and, by Corollary 1,  $e^{u_2} \in L^k(\Omega)$  for every  $k \geq 1$ . Thus  $|V_n| e^{u_n} \leq e^{u_{1n}} (e^{u_2 n} W)$  remains bounded in some  $L^q$ ,  $q > 1$ , and the conclusion follows.

Assume now that (18) holds. Suppose, by contradiction, that  $\|u_n\|_{L^\infty}$  is not bounded. We may then extract a subsequence such that  $\|u_{n_k}\|_{L^\infty} \rightarrow \infty$ . By passing to a further subsequence (still denoted  $n_k$ ) we may assume that  $|V_{n_k}| \leq W$  for some  $W \in L^p$  (see e.g. [1]), Théorème IV.9). We are therefore reduced to the previous case.

III.2. The main results

We now turn to the study of a sequence  $(u_n)$  of solutions of (8) under the assumptions

$$(20) \qquad V_n \geq 0 \text{ in } \Omega, \ \|V_n\|_{L^p} \leq C_1 \text{ and } \|e^{u_n}\|_{L^{p'}} \leq C_2$$

for some  $1 < p \leq \infty$ . A typical example is the sequence

$$u_n(x) = \log \frac{8n^2}{(1+n^2|x|^2)^2}$$

which satisfies  $-\Delta u_n = e^{u_n}$  in  $\mathbb{R}^2$  and  $\|e^{u_n}\|_{L^1(\mathbb{R}^2)} = 8\pi$ . Note that  $u_n(x) \rightarrow -\infty$  for all  $x \neq 0$  and  $u_n(0) \rightarrow +\infty$ . This example provides a very good description of the blow-up mechanism in the general case under the assumption (20). In fact, if a sequence  $(u_n)$  becomes unbounded then there is a

finite set  $S$  (possibly empty) where  $u_n$  tends to  $+\infty$  and elsewhere  $u_n$  tends to  $-\infty$ .

More precisely, define the "blow-up" set as follows:

$$S = \left\{ x \in \Omega; \text{there exists a sequence } x_n \text{ in } \Omega \text{ such that } x_n \rightarrow x \right. \\ \left. \text{and } u_n(x_n) \rightarrow +\infty \right\}.$$

Then we have

**Theorem 3.** Assume  $(u_n)$  is a sequence of solutions of (8) satisfying, for some  $1 < p \leq \infty$ ,

$$(21) \quad V_n \geq 0 \text{ in } \Omega,$$

$$(22) \quad \|V_n\|_{L^p} \leq C_1$$

and

$$(23) \quad \|e^{u_n}\|_{L^{p'}} \leq C_2.$$

Then, there exists a subsequence  $(u_{n_k})$  satisfying the following alternative:

either

$$(i) \quad (u_{n_k}) \text{ is bounded in } L_{loc}^\infty(\Omega)$$

or

$$(ii) \quad u_{n_k}(x) \rightarrow -\infty \text{ uniformly on compact subsets of } \Omega$$

or

$$(iii) \quad \text{the blow-up set } S \text{ (relative to } (u_{n_k})) \text{ is finite, nonempty and}$$

$u_{n_k}(x) \rightarrow -\infty$  uniformly on compact subsets of  $\Omega \setminus S$ . In addition  $V_{n_k} e^{u_{n_k}}$  converges in the sense of measures on  $\Omega$  to  $\sum_i \alpha_i \delta_{a_i}$  with  $\alpha_i \geq 4\pi/p'$   $\forall i$  and

$$S = \bigcup_i \{a_i\}.$$

Before giving the proof of Theorem 3 we mention some Corollaries.

**Corollary 6.** Assume  $(u_n)$  is a sequence of solutions of (8) with  $u_n = 0$  on  $\partial\Omega$ ,



satisfying (21), (22), and (23).

Then  $(u_n)$  is bounded in  $L^{\infty}_{loc}(\Omega)$ .

**Proof.** By the maximum principle  $u_n \geq 0$  on  $\Omega$  and therefore cases (ii) and (iii) in Theorem 3 are excluded for all subsequences. Therefore the (full) sequence  $(u_n)$  is bounded in  $L^{\infty}_{loc}(\Omega)$ .

**Remark 7.** One may wonder whether the conclusion of Corollary 6 holds uniformly up to the boundary (since we impose here the boundary condition  $u_n = 0$  on  $\partial\Omega$ ). This is not true as is shown in Section III.3 (Example 6). However it is plausible that a stronger assumption about the  $V_n$ 's yields an estimate up to the boundary. For example, here is an

**Open problem 1:** Suppose  $(u_n)$  is a sequence of solutions of (8) with  $u_n = 0$  on  $\partial\Omega$  satisfying (21),

$$(24) \quad V_n \rightarrow V \text{ in } C^0(\overline{\Omega})$$

and

$$(25) \quad \|e^{u_n}\|_{L^1} \leq C.$$

Can one conclude that  $\|u_n\|_{L^{\infty}} \leq C$ ?

**Remark 8.** The conclusion of Corollary 6 also fails if we remove assumption (21) (i.e.  $V_n \geq 0$  on  $\Omega$ ); see [8].

Another obvious consequence of Theorem 3 is:

**Corollary 7.** Assume  $(u_n)$  is a sequence of solutions of (8) satisfying (21), (22) and (23). Assume in addition

$$(26) \quad u_n \geq -M \text{ in } \Omega, \forall n$$

for some positive constant  $M$ , or more generally

$$(27) \quad \|u_n^-\|_{L^1} \leq M \quad \forall n.$$

Then  $(u_n)$  is bounded in  $L^{\infty}_{\text{loc}}(\Omega)$ .

**Corollary 8.** Assume  $(u_n)$  is a sequence of solutions of (8) satisfying (26) and

$$(28) \quad 0 < a \leq V_n \leq b < \infty \quad \text{in } \Omega$$

for some constants  $a, b$ .

Then  $(u_n)$  is bounded in  $L^{\infty}_{\text{loc}}(\Omega)$ .

**Proof.** In view of Corollary 7 we have only to show that  $(e^{u_n})$  is bounded in  $L^1_{\text{loc}}(\Omega)$ . We may always assume that  $M = 0$ , i.e.  $u_n \geq 0$  (this amounts to replace  $u_n$  by  $u_n + M$ ). Let  $\varphi_1$  be the first eigenfunction of  $-\Delta$  on  $\Omega$  with zero Dirichlet conditions and let  $\lambda_1$  be the corresponding eigenvalue.

Multiplying (8) by  $\varphi_1$  and integrating we obtain

$$\int_{\Omega} V_n e^{u_n} \varphi_1 = \int_{\Omega} u_n \frac{\partial \varphi_1}{\partial \nu} + \lambda_1 \int_{\Omega} u_n \varphi_1$$

where  $\nu$  is the outward normal. Using (28),  $u_n \geq 0$  and  $\frac{\partial \varphi_1}{\partial \nu} \leq 0$  we obtain

$$a \int_{\Omega} e^{u_n} \varphi_1 \leq \lambda_1 \int_{\Omega} u_n \varphi_1$$

This provides an upper bound for  $\int_{\Omega} e^{u_n} \varphi_1$ . Therefore  $(e^{u_n})$  is bounded in  $L^1_{\text{loc}}(\Omega)$  and the conclusion follows.

**Remark 9.** There are two natural questions suggested by Corollary 8:

**Open problem 2:** Suppose  $(u_n)$  is a sequence of solutions of (8) with  $u_n = 0$  on  $\partial\Omega$  satisfying (28). Can one conclude that  $(u_n)$  is bounded in  $L^{\infty}(\Omega)$ ? Is this true if we assume in addition that  $\|e^{u_n}\|_{L^1} \leq C$ ?

**Open problem 3:** Assume  $(u_n)$  is a sequence of solutions of (8) satisfying (26) and (28). Let  $K$  be a compact subset of  $\Omega$ . What is the optimal bound for  $\sup_K u_n$  as a function of  $M$ ? Does one have

$$(29) \quad \sup_K u_n \leq C_1 M + C_2$$

for some positive constants  $C_1, C_2$  depending only on  $a, b, K$  and  $\Omega$ ? Can one take  $C_1 = 1$  if  $V_n(x) \equiv 1$ ?

[Note that (29) holds with  $C_1 = 1$  for the special sequence

$$u_n(x) = \log \frac{8n^2}{(1+n^2|x|^2)^2}].$$

Proof of Theorem 3. Since  $(V_n e^{u_n})$  is bounded in  $L^1(\Omega)$  we may extract a subsequence (still denoted  $V_n e^{u_n}$ ) such that  $V_n e^{u_n}$  converges in the sense of measures on  $\Omega$  to some nonnegative bounded measure  $\mu$ , i.e.

$$(30) \quad \int V_n e^{u_n} \psi \rightarrow \int \psi d\mu$$

for every  $\psi \in C_c(\Omega)$ .

Definition: We say that a point  $x_0 \in \Omega$  is a regular point if there is a function  $\psi \in C_c(\Omega)$ ,  $0 \leq \psi \leq 1$ , with  $\psi = 1$  in some neighborhood of  $x_0$ , such that

$$(31) \quad \int \psi d\mu < 4\pi/p'.$$

It follows from Corollary 4 (applied in a small ball around  $x_0$ ) that if  $x_0$  is a regular point then there is some  $R_0 > 0$  such that

$$(32) \quad (u_n^+) \text{ is bounded in } L^p(B_{R_0}(x_0)).$$

[Note that (13) holds since  $(e^{u_n})$  is bounded in  $L^{p'}(\Omega)$ ].

We denote by  $\Sigma$  the set of nonregular points in  $\Omega$ . Clearly  $x_0 \in \Sigma$  iff  $\mu(\{x_0\}) \geq 4\pi/p'$ . Since  $\mu$  is a bounded measure (with  $\int d\mu \leq C_1 C_2$ ) it follows that  $\Sigma$  is finite and

$$\text{card}(\Sigma) \leq C_1 C_2 p' / 4\pi.$$

We now split the proof of Theorem 3 into 3 steps.

Step 1:  $S = \Sigma$ .

Clearly  $S \subset \Sigma$  by (32). Conversely, suppose  $x_0 \in \Sigma$ . Then we have

(33)  $\forall R > 0, \lim \|u_n^+\|_{L^\infty(B_R(x_0))} = +\infty.$

Otherwise there would be some  $R_0 > 0$  and a subsequence such that

$\|u_{n_k}^+\|_{L^\infty(B_{R_0}(x_0))} \leq C.$  In particular  $\|e^{u_{n_k}}\|_{L^\infty(B_{R_0}(x_0))} \leq C$  and therefore

$$\int_{B_R(x_0)} v_{n_k} e^{u_{n_k}} \leq CC_1 R^{2/p'} \text{ for all } R < R_0.$$

This implies (31) for some suitable  $\psi$ . Therefore  $x_0$  is regular – a contradiction.

Hence we have established (33). Choose  $R > 0$  small enough so that  $B_R(x_0)$  does not contain any other point of  $\Sigma$ . Let  $x_n \in B_R(x_0)$  be such that

$$u_n^+(x_n) = \max_{B_R(x_0)} u_n^+ \rightarrow +\infty.$$

We claim that  $x_n \rightarrow x_0$ . Otherwise there would be a subsequence

$x_{n_k} \rightarrow \bar{x} \neq x_0$  and  $\bar{x} \notin \Sigma$ , i.e.  $\bar{x}$  is a regular point. This is impossible in view of (32). Hence we have established that  $x_0 \in S$ . This completes the proof of Step 1.

Step 2:  $S = \emptyset$  implies (i) or (ii) holds.

By (32)  $(u_n^+)$  is bounded in  $L^\infty_{loc}(\Omega)$  and therefore  $f_n = v_n e^{u_n}$  is bounded in  $L^p_{loc}(\Omega)$ . This implies that  $\mu \in L^1(\Omega) \cap L^p_{loc}(\Omega)$ . Let  $v_n$  be the solution of

$$\begin{cases} -\Delta v_n = f_n & \text{in } \Omega, \\ v_n = 0 & \text{on } \partial\Omega. \end{cases}$$

Clearly,  $v_n \rightarrow v$  uniformly on every compact subset of  $\Omega$ , where  $v$  is the solution of

$$\begin{cases} -\Delta v = \mu & \text{in } \Omega, \\ v = 0 & \text{on } \partial\Omega. \end{cases}$$

Let  $w_n = u_n - v_n$  so that  $\Delta w_n = 0$  on  $\Omega$  and  $w_n^+$  is bounded in  $L^\infty_{loc}(\Omega)$ . By Harnack's principle we find that:

either

(34) a subsequence  $(w_{n_k})$  is bounded in  $L^\infty_{loc}(\Omega)$ ,

or

(35)  $(w_n)$  converges uniformly to  $-\infty$  on compact subsets of  $\Omega$ .

Case (i) corresponds to (34) and case (ii) to (35).

Step 3.  $S \neq \emptyset$  implies (iii) holds.

By (32)  $(u_n^+)$  is bounded in  $L_{loc}^\infty(\Omega \setminus S)$  and therefore  $f_n = V_n e^{u_n}$  is bounded in  $L_{loc}^\infty(\Omega \setminus S)$ . This implies that  $\mu$  is a bounded measure on  $\Omega$  with  $\mu \in L_{loc}^p(\Omega \setminus S)$ . (A basic difference with Step 2 is that here  $\mu$  is a measure, not an  $L^1$  function, and, as will be shown later,  $\mu$  is a sum of Dirac masses). Let  $v_n$ ,  $v$  and  $w_n$  be defined as in Step 2. Then  $v_n \rightarrow v$  uniformly on compact subsets of  $\Omega \setminus S$ . As above, by Harnack's principle, we find that

either

(36) a subsequence  $(w_{n_k})$  is bounded in  $L_{loc}^\infty(\Omega \setminus S)$

or

(37)  $(w_n)$  converges to  $-\infty$  uniformly on compact subsets of  $\Omega \setminus S$ .

We claim that (36) does not happen. Fix some point  $x_0 \in S$  and  $R > 0$  small so that  $x_0$  is the only point of  $S$  in  $\bar{B}_R(x_0)$ . Assume (36) holds, so that  $(w_{n_k})$  is bounded in  $L^\infty(\partial B_R(x_0))$  and similarly for  $(v_n)$ . Therefore  $(u_{n_k})$  is bounded in  $L^\infty(\partial B_R(x_0))$ , say by  $C$ . Let  $z_{n_k}$  be the solution of

$$\begin{cases} -\Delta z_{n_k} = f_{n_k} & \text{in } B_R(x_0), \\ z_{n_k} = -C & \text{on } \partial B_R(x_0). \end{cases}$$

By the maximum principle  $u_{n_k} \geq z_{n_k}$  in  $B_R(x_0)$ .

In particular

$$(38) \quad \int_{\partial B_R(x_0)} e^{p' z_{n_k}} \leq \int_{\partial B_R(x_0)} e^{p' u_{n_k}} \leq C_2^{p'}.$$

On the other hand  $z_{n_k} \rightarrow z$  a.e. (even uniformly on compact subsets of  $B_R(x_0) \setminus \{x_0\}$ ) where  $z$  is the solution of

$$\begin{cases} -\Delta z = \mu & \text{in } B_R(x_0), \\ z = -C & \text{on } \partial B_R(x_0). \end{cases}$$

Finally note that since  $x_0 \in S$  is not a regular point we have  $\mu(\{x_0\}) \geq 4\pi/p'$ .

This implies that  $\mu \geq \frac{4\pi}{p'} \delta_{x_0}$  and therefore

$$z(x) \geq \frac{2}{p'} \log \frac{1}{|x-x_0|} + o(1) \text{ as } x \rightarrow x_0.$$

Thus  $e^{p'z} \geq C/|x-x_0|^2$  with  $C > 0$ . Hence  $\int_{B_R(x_0)} e^{p'z} = \infty$ . On the other

hand, by (38) and Fatou's Lemma we find that

$$\int e^{p'z} \leq C_2^{p'}.$$

A contradiction. Hence we have shown that (37) holds. Consequently  $(u_n)$  converges to  $-\infty$  uniformly on compact subsets of  $\Omega \setminus S$ . Therefore  $V_n e^{u_n} \rightarrow 0$  in  $L^p_{loc}(\Omega \setminus S)$  and hence  $\mu$  is supported on  $S$ . This means that  $\mu = \sum_i \alpha_i \delta_{a_i}$  with  $S = \bigcup_i \{a_i\}$ . The argument above gives that  $\alpha_i \geq 4\pi/p'$  for each  $i$ .

**Remark 10.** The conclusion (iii) in Theorem 3 involves a finite sum of Dirac masses  $\sum_i \alpha_i \delta_{a_i}$  with coefficients  $\alpha_i \geq 4\pi/p'$ . The  $\alpha_i$ 's as well as the  $a_i$ 's can

be chosen arbitrarily. More precisely given any finite set  $S = \bigcup_{i=1}^k \{a_i\}$  and any

$\alpha_i > 4\pi/p'$  there exist sequences  $(u_n)$  and  $(V_n)$  as in Theorem 3 such that

$V_n e^{u_n}$  converges to  $\sum_{i=1}^k \alpha_i \delta_{a_i}$ .

To construct such sequences we proceed as follows. Set, for  $1 \leq i \leq k$ ,

$$v_{i,n} = \begin{cases} -\frac{A_i}{4} n^{2\beta_i} |x-a_i|^2 + \frac{A_i}{4} & \text{if } |x-a_i| < 1/n^{\beta_i}, \\ \frac{A_i}{2} \log\left(\frac{1}{n^{\beta_i} |x-a_i|}\right) & \text{if } |x-a_i| \geq 1/n^{\beta_i} \end{cases}$$

where  $A_i = \alpha_i/\pi > 4/p'$  and  $\beta_i$  is defined by the relation  $\beta_i(-\frac{A_i}{2} - \frac{2}{p'}) = 1$ .

Let  $u_n = \sum_{i=1}^k v_{i,n} + \sigma_n$  where  $\sigma_n = ((k-1) + \frac{2}{p'} \sum_{i=1}^k \beta_i) \log n$ . A direct

computation shows that  $V_n = (-\Delta u_n)e^{-u_n}$  satisfies (21) and (22); moreover  $(e^{u_n})$  is bounded in  $L^{p'}$  and  $V_n e^{u_n}$  converges to  $\sum_{i=1}^k \alpha_i \delta_{a_i}$ .

We believe that under additional conditions on the  $V_n$ 's the  $\alpha_i$ 's in Theorem 3 cannot take arbitrary values ( $> 4\pi/p'$ ):

Open problem 4: Assume  $(u_n)$  is a sequence of solutions of (8) satisfying

$V_n \geq 0$  on  $\Omega$ ,  $V_n \rightarrow V$  uniformly in  $\bar{\Omega}$  with  $V_n, V \in C^0(\bar{\Omega})$  and  $\|e^{u_n}\|_{L^1} \leq C$ . Assume  $S \neq \emptyset$  so that case (iii) holds. Can one conclude that  $V_{n_k} e^{u_{n_k}}$  converges to  $8\pi \sum m_i \delta_{a_i}$  with  $m_i \in \mathbb{N}$ ?

Evidence in favor of a positive answer comes from the fact that after a blow-up near  $a_i$  we are led to a solution of  $-\Delta v = ce^v$  on  $\mathbb{R}^2$  with  $c = V(a_i)$  and  $\int_{\mathbb{R}^2} e^v < \infty$ . It follows from the result of [3] that  $\int_{\mathbb{R}^2} ce^v = 8\pi$ . On the other hand, the blow-up analysis gives (formally)  $\alpha_i = \int_{\mathbb{R}^2} ce^v$ .

In Theorem 3 the assumption  $\|e^{u_n}\|_{L^{p'}} \leq C$  provides some kind of bound from above for  $(u_n)$  and plays an important role in proving that the blow-up set  $S$  is finite. If we drop that assumption little can be said in the general case. For instance, we may have a sequence  $(u_n)$  of solutions of

$$-\Delta u_n = e^{u_n} \quad \text{on } \Omega$$

(with  $\|e^{u_n}\|_{L^1} \rightarrow \infty$ ) such that

$$\begin{cases} u_n \rightarrow +\infty & \text{on a line } S, \\ u_n \rightarrow -\infty & \text{in } \Omega \setminus S. \end{cases}$$

Example 4. The sequence

$$u_n(x, y) = 2nx - 2 \log(1 + e^{2nx}) + \log 8n^2$$

satisfies  $-\Delta u_n = e^{u_n}$ ,  $u_n(0, y) \rightarrow +\infty$  and  $u_n(x, y) \rightarrow -\infty$  for  $x \neq 0$ .

However, if we assume some bound from below for the  $u_n$ 's then there are

only two possibilities: either  $S = \Omega$  (total blow-up) or  $S$  is (locally) finite.

**Theorem 4.** Assume  $(u_n)$  is a sequence of solutions of (8) satisfying, for some  $1 < p \leq \infty$ , (21), (22) and

$$(39) \quad \|u_n^-\|_{L^1} \leq C$$

Then, there exists a subsequence  $(u_{n_k})$  satisfying the following alternative:

either

(i)  $u_{n_k} \rightarrow +\infty$  uniformly on compact subsets of  $\Omega$

or

(ii) the blow-up set  $S$  (relative to  $(u_{n_k})$ ) is locally finite (i.e. for each  $x \in \Omega$  there is some neighborhood  $N(x)$  of  $x$  such that  $N(x) \cap S$  is finite). Moreover  $(u_{n_k})$  is bounded in  $L_{loc}^p(\Omega \setminus S)$ .

**Remark 11.** Both cases in the alternative may occur:

**Example of (i).** Let  $v$  be any solution of  $-\Delta v = e^v$  in  $\mathbb{R}^2$ . Then  $u_n = v + n$  satisfies  $-\Delta u_n = V_n e^{u_n}$  with  $V_n = e^{-n}$  and  $u_n \rightarrow +\infty$  everywhere.

**Example of (ii).** Recall that  $v_n(x) = \log \frac{8n^2}{(1+n^2|x|^2)^2}$  satisfies  $-\Delta v_n = e^{v_n}$ .

Thus  $u_n = v_n + \log n^2$  satisfies  $-\Delta u_n = V_n e^{u_n}$  with  $V_n = 1/n^2$ . Note that  $u_n(0) \rightarrow +\infty$  while  $u_n(x)$  remains bounded for  $x \neq 0$ .

**Proof of Theorem 4.** Without loss of generality we may assume that

$$(40) \quad u_n \geq 0 \quad \text{in } \Omega.$$

Indeed, by Kato's inequality [5] we have

$$(41) \quad \Delta u_n^- \geq -(\Delta u_n) \chi_{\{u_n \leq 0\}} = V_n e^{u_n} \chi_{\{u_n \leq 0\}} \geq -|V_n|.$$

It follows from (39), (41) and standard elliptic estimates that  $(u_n^-)$  is bounded in



$L^{\infty}_{\text{loc}}(\Omega)$ . Passing to a smaller domain and adding a constant to  $(u_n)$  we may always assume that (40) holds.

We now split the proof into 3 cases.

Case 1: There exists a compact subset  $K \subset \Omega$  and a subsequence  $(u_{n_k})$  such that

$$(42) \quad \int_K V_{n_k} e^{u_{n_k}} \rightarrow +\infty.$$

Then (i) holds.

Indeed, let  $K'$  be any compact subset of  $\Omega$ . Using (40) we obtain

$$u_{n_k}(x) \geq \int_{\Omega} G(x,y) V_{n_k}(y) e^{u_{n_k}(y)} dy,$$

where  $G$  is the Green's function of  $-\Delta$  with Dirichlet condition on  $\partial\Omega$ . Since  $G(x,y) \geq \alpha > 0 \forall x \in K', \forall y \in K$  we see that, for  $x \in K'$ ,

$$u_{n_k}(x) \geq \alpha \int_K V_{n_k} e^{u_{n_k}} \rightarrow +\infty.$$

Case 2.  $(V_n e^{u_n})$  is bounded in  $L^1_{\text{loc}}(\Omega)$  and there exists a compact subset  $K \subset \Omega$  such that, for a subsequence,

$$\int_K u_{n_k} \rightarrow +\infty.$$

Then (i) holds.

Indeed, let  $K'$  be any compact subset of  $\Omega$ . Let  $\omega$  be an open set such that  $K \cup K' \subset \omega \subset\subset \Omega$ . In  $\omega$ , split  $u_n$  as  $u_n = u_{1n} + u_{2n}$  where  $u_{1n}$  is the solution of

$$\begin{cases} -\Delta u_{1n} = V_n e^{u_n} & \text{in } \omega, \\ u_{1n} = 0 & \text{on } \partial\omega. \end{cases}$$

Note that  $(u_{1n})$  is bounded in  $L^1(\omega)$  and  $u_{2n}$  satisfies

$$\begin{cases} -\Delta u_{2n} = 0 & \text{in } \omega, \\ u_{2n} \geq 0 & \text{on } \partial\omega \end{cases}$$

Thus  $u_{2n} \geq 0$  in  $\omega$  and by Harnack's principle

$$(43) \quad \sup_{K \cup K'} u_{2n} \leq C \inf_{K \cup K'} u_{2n} \leq C \inf_{K'} u_n.$$

On the other hand

$$\int_K u_{2n} \leq C \sup_K u_{2n} \leq C \sup_{K \cup K'} u_{2n}$$

and

$$\int_K u_{2n} = \int_K u_n - \int_K u_{1n} \geq \int_K u_n - C.$$

It follows that  $\inf_{K'} u_{n_k} \rightarrow +\infty$  and thus (i) holds.

We are left with:

Case 3:  $(V_n e^{u_n})$  and  $(u_n)$  are bounded in  $L^1_{loc}(\Omega)$ . Then (ii) holds.

We proceed here as in the proof of Theorem 3. We extract a subsequence (still denoted  $V_n e^{u_n}$ ) such that  $V_n e^{u_n}$  converges in the sense of measures to some nonnegative (possibly unbounded) measure  $\mu$ , i.e.

$$\int V_n e^{u_n} \psi \rightarrow \int \psi d\mu$$

for every  $\psi \in C_c(\Omega)$ . We say that a point  $x_0 \in \Omega$  is a regular point if there is a function  $\psi \in C_c(\Omega)$ ,  $0 \leq \psi \leq 1$ , with  $\psi = 1$  in some neighborhood of  $x_0$ , such that

$$\int \psi d\mu < 4\pi/p'.$$

It follows from Corollary 4 (applied in a small ball around  $x_0$ ) that if  $x_0$  is a regular point then there is some  $R_0 > 0$  such that

$$(44) \quad (u_n) \text{ is bounded in } L^\infty(B_{R_0}(x_0)).$$

We denote by  $\Sigma$  the set of nonregular points in  $\Omega$ . Clearly  $x_0 \in \Sigma$  if  $\mu(\{x_0\}) \geq 4\pi/p'$ . It follows that  $\Sigma$  is locally finite and for every compact subset  $K$  of  $\Omega$

$$\text{card}(\Sigma \cap K) \leq (p'/4\pi) \int_K d\mu.$$

We have  $S = \Sigma$  as in the proof of Theorem 3 (Step 1). Thus  $S$  is locally finite and by (44)  $(u_n)$  is bounded in  $L^{\infty}_{\text{loc}}(\Omega \setminus S)$ , i.e. (ii) holds.

### III.3. Variants and counterexamples

1. Suppose that instead of a sequence of solutions of (8) we have a sequence of subsolutions, i.e.

$$-\Delta u_n \leq V_n(x)e^{u_n} \quad \text{in } \Omega.$$

It is easy to adapt the arguments of Section III.1 to obtain estimates for  $\|u_n^+\|_{L^{\infty}}$  under smallness or uniform domination assumption. However the analogue of Corollary 6 for subsolutions does not hold as may be seen from the following:

Example 5. There is a sequence  $(u_n)$  satisfying

$$\begin{cases} -\Delta u_n \leq e^{u_n} & \text{in } \Omega = B_1, \\ u_n = 0 & \text{on } \partial\Omega \end{cases}$$

with

$$\int_{\Omega} e^{u_n} \leq C$$

and such that  $u_n(0) \rightarrow +\infty$ . First, note that the function

$$\varphi_{\epsilon}(x) = \log \frac{8\epsilon^2}{(\epsilon^2 + |x|^2)^2}$$

satisfies

$$-\Delta \varphi_{\epsilon} = e^{\varphi_{\epsilon}} \quad \forall \epsilon > 0$$

and

$$\int_{\mathbb{R}^2} e^{\varphi_{\epsilon}} = 8\pi \quad \forall \epsilon > 0.$$

Hence the function  $u_n = \varphi_{1/n}^+$  has all the required properties. The same example can be used to produce sequences  $(v_n)$  and  $(V_n)$  such that

$$\begin{cases} -\Delta v_n \leq V_n e^{v_n} & \text{in } \Omega = B_1 \\ v_n = 0 & \text{on } \partial\Omega \end{cases}$$

such that  $V_n \geq 0$ ,  $\|V_n\|_{L^p} \leq C$ ,  $\|e^{v_n}\|_{L^{p'}} \leq C$ ,  $1 < p < \infty$ , and  $v_n(0) \rightarrow +\infty$ .

It suffices to take  $v_n = \frac{1}{p}$ ,  $u_n$  and  $V_n = \frac{1}{p} e^{\frac{1}{p} u_n}$ .

2. The same kind of example shows that the conclusion of Theorem 2 does not hold uniformly. More precisely there are sequences  $(u_n)$  and  $(V_n)$  such that

$$-\Delta u_n = V_n e^{u_n} \quad \text{on } \mathbb{R}^2$$

with  $\|V_n\|_{L^p(\mathbb{R}^2)} \leq C$ ,  $\|e^{u_n}\|_{L^{p'}(\mathbb{R}^2)} \leq C$ ,  $1 < p \leq \infty$ , such that  $u_n(0) \rightarrow +\infty$ .

One may take for instance  $u_n = \frac{1}{p} \varphi_{1/n}$  and  $V_n = \frac{1}{p} \exp(\frac{1}{p} \varphi_{1/n})$ .

3. The conclusion of Corollary 6 cannot be strengthened to  $\|u_n\|_{L^\infty} \leq C$ . There are sequences  $(u_n)$  and  $(V_n)$  satisfying

$$-\Delta u_n = V_n e^{u_n} \quad \text{in } \Omega = B_1$$

$$u_n = 0 \quad \text{on } \partial\Omega$$

$$V_n \geq 0 \quad \text{in } \Omega$$

$$\|V_n\|_{L^p} \leq C,$$

$$\|e^{u_n}\|_{L^{p'}} \leq C,$$

with  $1 < p \leq \infty$  and such that  $\|u_n\|_{L^\infty} \rightarrow \infty$ . It suffices to construct such an

example when  $p = \infty$ . For a general  $1 < p < \infty$  we may use the  $p = \infty$

example and note that  $\tilde{u}_n = \frac{1}{p} u_n$  satisfies  $-\Delta \tilde{u}_n = \tilde{V}_n e^{\tilde{u}_n}$  with

$\tilde{V}_n = \frac{1}{p} V_n \exp(\frac{1}{p} u_n)$  so that  $\|\tilde{V}_n\|_{L^p} \leq C$  and  $\|e^{\tilde{u}_n}\|_{L^{p'}} \leq C$ .

**Example 6.** Let  $\Omega$  be the unit disc centered at  $(1,0)$ . Set  $a_\epsilon = (d_\epsilon, 0)$  with

$\epsilon < d_\epsilon < 1$ . Let  $A > 1$  be a constant and let

$$f_\epsilon = \begin{cases} \frac{4A}{\epsilon^2} & \text{in } B_\epsilon(a_\epsilon) \\ 0 & \text{otherwise.} \end{cases}$$

Let  $u_\epsilon$  be the solution of

$$\begin{cases} -\Delta u_\epsilon = f_\epsilon & \text{in } \Omega, \\ u_\epsilon = 0 & \text{on } \partial\Omega. \end{cases}$$

Let  $V_\epsilon$  be defined by

$$V_\epsilon = f_\epsilon e^{-u_\epsilon}$$

so that  $-\Delta u_\epsilon = V_\epsilon e^{u_\epsilon}$ . We claim that, for an appropriate choice of  $d_\epsilon$ , we have

$$(45) \quad \|V_\epsilon\|_{L^\infty} \leq C$$

$$(46) \quad \int_\Omega e^{u_\epsilon} \leq C.$$

while  $u_\epsilon(a_\epsilon) \rightarrow +\infty$ .

Verification of (45). Let  $v_\epsilon$  be the solution of

$$\begin{cases} -\Delta v_\epsilon = f_\epsilon & \text{in } B_{d_\epsilon}(a_\epsilon), \\ v_\epsilon = 0 & \text{on } \partial B_{d_\epsilon}(a_\epsilon). \end{cases}$$

By the maximum principle we have  $v_\epsilon \leq u_\epsilon$  in  $B_{d_\epsilon}(a_\epsilon)$  so that

$$\|V_\epsilon\|_{L^\infty} = \|f_\epsilon e^{-u_\epsilon}\|_{L^\infty} \leq \frac{4A}{\epsilon^2} \|e^{-v_\epsilon}\|_{L^\infty(B_{d_\epsilon}(a_\epsilon))}.$$

But  $v_\epsilon$  is given explicitly by

$$v_\epsilon = \begin{cases} -\frac{4A}{\epsilon^2} r^2 + \alpha_\epsilon & 0 \leq r < \epsilon \\ 2A \log\left(\frac{d_\epsilon}{r}\right) & \epsilon < r < d_\epsilon \end{cases}$$

where  $r = |x - a_\epsilon|$  and  $\alpha_\epsilon = A + 2A \log(\frac{d_\epsilon}{\epsilon})$ . Thus

$$\|e^{-v_\epsilon}\|_{L^\infty(B_\epsilon(a_\epsilon))} = e^{A - \alpha_\epsilon} = (\frac{\epsilon}{d_\epsilon})^{2A}.$$

Hence (45) holds provided

$$(47) \quad \frac{1}{\epsilon^{2A}} (\frac{\epsilon}{d_\epsilon})^{2A} \leq C.$$

Verification of (46). Let  $G$  be the half-plane

$$G = \{(x_1, x_2) \in \mathbb{R}^2; x_1 > 0\}.$$

Let  $w_\epsilon$  be the solution of

$$\begin{cases} -\Delta w_\epsilon = f_\epsilon & \text{in } G, \\ w_\epsilon = 0 & \text{on } \partial G. \end{cases}$$

By the maximum principle we have  $u_\epsilon \leq w_\epsilon$  in  $\Omega$  and thus

$$\int_\Omega e^{u_\epsilon} \leq \int_\Omega e^{w_\epsilon}.$$

But  $w_\epsilon$  is given explicitly by

$$w_\epsilon = \begin{cases} -\frac{4A}{\epsilon^2} |x - a_\epsilon|^2 + \beta_\epsilon + 2A \log|x - a'_\epsilon| & \text{if } |x - a_\epsilon| < \epsilon \\ 2A \log\left(\frac{|x - a'_\epsilon|}{|x - a_\epsilon|}\right) & \text{otherwise} \end{cases}$$

where  $a'_\epsilon = -a_\epsilon$  and  $\beta_\epsilon = A - 2A \log \epsilon$ . We have

$$w_\epsilon(x) \leq C + 2A \log\left(\frac{d_\epsilon}{\epsilon}\right) \quad \text{if } |x - a_\epsilon| < \epsilon$$

(since  $|x - a'_\epsilon| < |x - a_\epsilon| + 2d_\epsilon \leq \epsilon + 2d_\epsilon \leq 3d_\epsilon$ ),

$$w_\epsilon(x) \leq C + 2A \log\left(\frac{d_\epsilon}{|x - a_\epsilon|}\right) \quad \text{if } \epsilon \leq |x - a_\epsilon| < d_\epsilon$$

(since  $|x - a'_\epsilon| < 3d_\epsilon$ ) and

$$w_\epsilon \leq C \quad \text{if } |x - a_\epsilon| \geq d_\epsilon$$

(since  $|x-a'_\epsilon| < |x-a_\epsilon| + 2d_\epsilon \leq 3|x-a_\epsilon|$ ).

It follows that

$$\begin{aligned} \int_{\Omega} e^{w_\epsilon} &\leq C \epsilon^{2(\frac{d_\epsilon}{\epsilon})^{2A}} + C \int_{\epsilon}^{\frac{d_\epsilon}{\epsilon}} \left(\frac{d_\epsilon}{r}\right)^{2A} r \, dr + C \\ &\leq C \epsilon^{2(\frac{d_\epsilon}{\epsilon})^{2A}} + C. \end{aligned}$$

Hence (47) and (46) can be achieved by choosing  $d_\epsilon = \epsilon^{1-(1/A)}$ . Finally we have

$$u_\epsilon(a_\epsilon) \geq v_\epsilon(a_\epsilon) = \alpha_\epsilon \geq 2A \log\left(\frac{d_\epsilon}{\epsilon}\right) \rightarrow +\infty \text{ as } \epsilon \rightarrow 0. \text{ Note that in this}$$

Example  $\int_{\Omega} V_\epsilon e^{u_\epsilon} = 4A\pi$  can be made arbitrarily close to  $4\pi$ , showing once

more that assumption (10) in Corollary 3 is sharp.

4. One may combine the techniques of Sections III.1 and Section III.2. Assume for example that all the assumptions of Corollary 6 hold with  $1 < p < \infty$  and in addition

$$|V_n(x)| \leq W(x) \text{ in some fixed neighbourhood of } \partial\Omega$$

with  $W \in L^p$ . Then  $\|u_n\|_{L^\infty} \leq C$ .

Acknowledgement : The second author (F.M.) thanks New York University and Rutgers University for their hospitality during the preparation of this paper ; while at NYU, F.M. was partly supported by NSF Grant DMS-8806731.

#### References

- [1] H. Brezis, *Analyse Fonctionnelle*, Masson (1983).
- [2] S.Y.A. Chang - P.C. Yang, Prescribing Gaussian curvature on  $S^2$ , *Acta Math.* **159** (1987), 215-259.
- [3] W. Chen - C. Li, Classification of solutions of some nonlinear elliptic equations, preprint (1990).
- [4] F. John - L. Nirenberg, On functions of bounded mean oscillation, *Comm. Pure Appl. Math.* **14** (1960), 93-140.

- [5] T. Kato, Schrödinger operators with singular potentials, Israel J. Math. 13 (1972), 135–148.
- [6] K. Nagasaki – T. Suzuki, Asymptotic analysis for two dimensional elliptic eigenvalue problems with exponentially-dominated nonlinearities, Asymptotic Analysis (to appear).
- [7] T. Suzuki, Introduction to geometric potential theory, preprint (1990).
- [8] S. Wang, An example of a blow-up sequence for  $-\Delta u = V(x)e^u$  (to appear).

Received December 1990

Revised April 1991