Homework set 6 — due March 05

Problem 1. Compute

$$\oint_{\alpha} \frac{z - i + 1}{z^2 - z(1 + i)} dz$$

along the following three circles, all positively oriented:


- (i) The circle of radius 2 centred at $z_0 = 1$
- (ii) The circle of radius 1 centred at $z_0 = 1 + i$
- (iii) The circle of radius 1/2 centred at $z_0 = 2$

Problem 2. Use the function $f(z) = \frac{e^{iz}}{z}$ and the contour sketched in the figure to show that

$$\int_0^\infty \frac{\sin(x)}{x} \mathrm{d}x = \frac{\pi}{2}.$$

Hints: (a) Since the integrand is even and continuous at x = 0,

$$\int_0^\infty \frac{\sin(x)}{x} \mathrm{d}x = \frac{1}{2} \lim_{R \to \infty} \lim_{\epsilon \to 0} \left[\int_{-R}^{-\epsilon} \frac{\sin(x)}{x} \mathrm{d}x + \int_{\epsilon}^R \frac{\sin(x)}{x} \mathrm{d}x \right].$$



(b) On the two curves c_{ϵ} and c_R , you are allowed to exchange the order to integrating and taking limits.


Problem 3. Let $0 < a \le 1$. Use the contour sketched in the figure to show that

$$\int_0^\infty e^{-(1+ia)^2 t^2} dt = \frac{\sqrt{\pi}}{2(1+ia)}.$$

Conclude that

$$\int_0^\infty \cos(t^2)dt = \int_0^\infty \sin(t^2)dt = \sqrt{\frac{\pi}{8}}.$$

Hint: You can use that $\int_0^\infty \mathrm{e}^{-t^2} dt = \sqrt{\pi}/2$. While evaluating the integral along the vertical segment, you may find the bound $t^2 \le t$ useful, which is valid for $0 \le t \le 1$.



Problem 4. Let f, g be two holomorphic functions in a domain Ω . Let α be a simple closed curve in Ω such that its complete interior lies in Ω . Show that if f(z) = g(z) for every $z \in \alpha$, then f(z) = g(z) for all z in the interior of α .