

Worksheet 3: Phototropic growth in plants

Name: _____ Class: _____ No.: _____


Topic notes



- **Phototropism**（向光性） is growth in response to light direction. Shoots usually grow towards light (**positive phototropism**（向光性）). The left diagram shows a bending coleoptile（胚芽鞘）: part of the seedling's shoot that protects young leaves.
- The tip senses light from one side and contains **auxin**（生長素）, a chemical substance that is a **plant hormone**. More **auxin**（生長素） **passes down** the **shaded** side, stimulating cells in the elongation（延長） region to **elongate**（延長） and promote growth.
- Agar（瓊脂） allows chemical substances to pass through it.



Coleoptile（胚芽鞘）: more **auxin**（生長素） on the **shaded** right



Cell detail: **on the right,** cells **elongate（延長） more**

Light from the left

↓

The right side is **shaded**

↓

More **auxin（生長素） passes down the shaded side**

↓

On the right, cells **elongate（延長） more**


↓

The shoot **bends left**


Part 1 | Context and fair comparisons

Aim: investigate how shading position affects coleoptile（胚芽鞘） growth direction.

Design A · Light blocked at the tip



Design B · Light blocked below the tip



1. Identify the variables: tick the appropriate factors in each row. You may choose more than one.

Variable type	Factors to choose from
Independent variable (the factor changed on purpose)	☐ Time exposed to light ☐ The coleoptile（胚芽鞘）'s bending response ☐ Light direction and intensity ☐ Coleoptile（胚芽鞘） type, starting height and growth stage ☑ **Light-blocking treatment** ☐ Temperature and water supply
Dependent variable (the result measured or observed)	☐ Time exposed to light ☑ **The coleoptile（胚芽鞘）'s bending response** ☐ Light direction and intensity ☐ Coleoptile（胚芽鞘） type, starting height and growth stage ☐ Light-blocking treatment ☐ Temperature and water supply
Controlled variables (the factors kept the same)	☑ **Time exposed to light** ☐ The coleoptile（胚芽鞘）'s bending response ☑ **Light direction and intensity** ☑ **Coleoptile（胚芽鞘） type, starting height and growth stage** ☐ Light-blocking treatment ☑ **Temperature and water supply**

2. What assumptions does this inquiry make?

An assumption is a condition we have not proved but take as true, to help make the experiment results valid.

Choose all suitable assumptions. You may choose more than one.

☑ **The coleoptiles（胚芽鞘） start in similar growth conditions.**

☑ **The light-blocking cap and sleeve do not limit elongation（延長） or bending.**

☑ **The treatments do not cause clear temperature differences between setups.**

☐ If the plants are the same type, their starting size does not need to be controlled.

3. Control group

To find out whether blocking light affects the coleoptile（胚芽鞘）'s growth direction, how would you design the control group?

(Hint: The control group provides a baseline for comparison. It helps you confirm that a change is caused by the independent variable rather than other external factors, so the control group does not receive the independent variable treatment.)


Add an intact, uncovered coleoptile（胚芽鞘） as the comparison baseline for the two shading treatments; keep all other conditions the same.

Part 2 | Results and analysis


2.1 Main investigation: effect of shading on growth

Light is on the left. The sleeve covers below the exposed tip without restricting growth.


Setup A · Light blocked at the tip



Setup B · Light blocked below the tip



Setup C · No light blocked




4. Record each setup's bending direction in the table.

Setup	Bending direction
A	(Bends left ／ Bends right ／ **No clear bending**)
B	(**Bends left** ／ Bends right ／ No clear bending)
C	(**Bends left** ／ Bends right ／ No clear bending)


5. Compare C and A: after **blocking light at the tip**, what changes?
   (**No clear bending towards the light** ／ It shows the same bending towards light)
6. Compare C and B: when the lower part **does not** receive light directly, does the coleoptile (胚芽鞘) still bend?
   (**It can still bend towards light** ／ No clear bending towards the light)
7. Which part may **sense light from one side**?
   (**Tip** ／ Only the lower part)

Extension investigation: effect of tip removal on growth

Setup C · No light blocked



Setup D · Tip removed



8. Record whether each setup **elongate**s (延長) in the table.


Setup	Elongation (延長)
C	(**Yes** ／ No)
D	(Yes ／ **No**)

9. What does comparing elongation (延長) in C and D support?
   ☑ **The presence of the tip also affects the coleoptile (胚芽鞘)'s growth / elongation (延長)**
   □ The presence of the tip does not affect the coleoptile (胚芽鞘)'s growth / elongation (延長)


2.3 Agar（瓊脂）

All cut-tip coleoptiles（胚芽鞘） start at similar sizes and stay in darkness. Blank agar（瓊脂） has not touched a tip; treated agar（瓊脂） has.


E
Blank agar（瓊脂）
Position: Centre




F
Treated agar（瓊脂）
Position: Centre



G
Treated agar（瓊脂）
Position: Left



H
Treated agar（瓊脂）
Position: Right



10. Record each setup's angle and bending direction.

Setup	Angle (°)	Bending direction
E	0°	（Bends left ／ Bends right ／ No clear bending）
F	0°	（Bends left ／ Bends right ／ No clear bending）
G	35°	（Bends left ／ Bends right ／ No clear bending）
H	35°	（Bends left ／ Bends right ／ No clear bending）

11. What does the comparison of E and F support?

☑ The tip may produce a growth-promoting effect that can be transferred

□ All agar（瓊脂） produces the same growth-promoting effect

12. What does the comparison of F, G and H support?


☑ The position of treated agar（瓊脂） affects the direction of coleoptile（胚芽鞘） bending

□ The position of treated agar（瓊脂） does not affect the direction of coleoptile（胚芽鞘） bending

13. When treated agar（瓊脂） is placed on the **left** side of a coleoptile（胚芽鞘） with its tip removed, compared with the right side, the left side receives （more ／ less） substance X, causing cells on the left to **elongate（延長）** （more ／ less） , finally causing （left ／ right） **bending** of the coleoptile（胚芽鞘）.

Part 3 | Integrated explanation

14. Use the experimental results and topic notes to explain why the coleoptile（胚芽鞘） in the diagram bends to the right.



Light is on the right and sensed at the tip. More auxin（生長素） is on the shaded left side and passes down. Left-side cells elongate（延長） more, causing bending towards the light on the right (positive phototropism（向光性）).