

VE203  
Discrete Math  
RC5

University of Michigan  
Shanghai Jiao Tong University  
Joint Institute

A set of navigation icons typically found in Beamer presentations, including symbols for back, forward, search, and other slide controls.

1 Ring (Optional)

## 2 Modular Arithmetic

- Modulo
- Fermat's (Little) Theorem
- Congruences
- Fast Modular Exponentiation
- Chinese Remainder Theorem
- RSA Cryptography

### 3 Q&A

## Definition

$\mathcal{R}_1.\langle R, + \rangle$  is an abelian group.

$\mathcal{R}_7$ . Multiplication is associative.

$\mathcal{R}_3$ . For all  $a, b, c \in R$ , the left distributive law,  $a \cdot (b + c) = (a \cdot b) + (a \cdot c)$  and the right distributive law  $(a + b) \cdot c = (a \cdot c) + (b \cdot c)$  hold.

# Ring

We are well aware that axioms  $\mathcal{R}_1, \mathcal{R}_2$ , and  $\mathcal{R}_3$  for a ring hold in any subset of the complex numbers that is a group under addition and that is closed under multiplication. For example,  $\langle \mathbb{Z}, +, \cdot \rangle, \langle \mathbb{Q}, +, \cdot \rangle, \langle \mathbb{R}, +, \cdot \rangle$ , and  $\langle \mathbb{C}, +, \cdot \rangle$  are rings.

0

Consider the cyclic group  $\langle \mathbb{Z}_n, + \rangle$ . If we define for  $a, b \in \mathbb{Z}_n$  the product  $ab$  as the remainder of the usual product of integers when divided by  $n$ , it can be shown that  $\langle \mathbb{Z}_n, +, \cdot \rangle$  is a ring. We shall feel free to use this fact. For example, in  $\mathbb{Z}_{10}$  we have  $(3)(7) = 1$ . This operation on  $\mathbb{Z}_n$  is multiplication modulo  $n$ . We do not check the ring axioms here. From now on,  $\mathbb{Z}_n$  will always be the ring  $\langle \mathbb{Z}_n, +, \cdot \rangle$ .

# Property

If  $R$  is a ring with additive identity  $0$  , then for any  $a, b \in R$  we have

1.  $0a = a0 = 0$ ,
2.  $a(-b) = (-a)b = -(ab)$ ,
3.  $(-a)(-b) = ab$ .

179

1999, 2000, 2001, 2002, 2003, 2004, 2005, 2006, 2007, 2008, 2009, 2010, 2011, 2012, 2013, 2014, 2015, 2016, 2017, 2018, 2019, 2020, 2021, 2022, 2023, 2024, 2025, 2026, 2027, 2028, 2029, 2030, 2031, 2032, 2033, 2034, 2035, 2036, 2037, 2038, 2039, 2040, 2041, 2042, 2043, 2044, 2045, 2046, 2047, 2048, 2049, 2050, 2051, 2052, 2053, 2054, 2055, 2056, 2057, 2058, 2059, 2060, 2061, 2062, 2063, 2064, 2065, 2066, 2067, 2068, 2069, 2070, 2071, 2072, 2073, 2074, 2075, 2076, 2077, 2078, 2079, 2080, 2081, 2082, 2083, 2084, 2085, 2086, 2087, 2088, 2089, 2090, 2091, 2092, 2093, 2094, 2095, 2096, 2097, 2098, 2099, 2100, 2101, 2102, 2103, 2104, 2105, 2106, 2107, 2108, 2109, 2110, 2111, 2112, 2113, 2114, 2115, 2116, 2117, 2118, 2119, 2120, 2121, 2122, 2123, 2124, 2125, 2126, 2127, 2128, 2129, 2130, 2131, 2132, 2133, 2134, 2135, 2136, 2137, 2138, 2139, 2140, 2141, 2142, 2143, 2144, 2145, 2146, 2147, 2148, 2149, 2150, 2151, 2152, 2153, 2154, 2155, 2156, 2157, 2158, 2159, 2160, 2161, 2162, 2163, 2164, 2165, 2166, 2167, 2168, 2169, 2170, 2171, 2172, 2173, 2174, 2175, 2176, 2177, 2178, 2179, 2180, 2181, 2182, 2183, 2184, 2185, 2186, 2187, 2188, 2189, 2190, 2191, 2192, 2193, 2194, 2195, 2196, 2197, 2198, 2199, 2200, 2201, 2202, 2203, 2204, 2205, 2206, 2207, 2208, 2209, 2210, 2211, 2212, 2213, 2214, 2215, 2216, 2217, 2218, 2219, 2220, 2221, 2222, 2223, 2224, 2225, 2226, 2227, 2228, 2229, 2230, 2231, 2232, 2233, 2234, 2235, 2236, 2237, 2238, 2239, 2240, 2241, 2242, 2243, 2244, 2245, 2246, 2247, 2248, 2249, 2250, 2251, 2252, 2253, 2254, 2255, 2256, 2257, 2258, 2259, 2260, 2261, 2262, 2263, 2264, 2265, 2266, 2267, 2268, 2269, 2270, 2271, 2272, 2273, 2274, 2275, 2276, 2277, 2278, 2279, 2280, 2281, 2282, 2283, 2284, 2285, 2286, 2287, 2288, 2289, 2290, 2291, 2292, 2293, 2294, 2295, 2296, 2297, 2298, 2299, 2300, 2301, 2302, 2303, 2304, 2305, 2306, 2307, 2308, 2309, 2310, 2311, 2312, 2313, 2314, 2315, 2316, 2317, 2318, 2319, 2320, 2321, 2322, 2323, 2324, 2325, 2326, 2327, 2328, 2329, 2330, 2331, 2332, 2333, 2334, 2335, 2336, 2337, 2338, 2339, 2340, 2341, 2342, 2343, 2344, 2345, 2346, 2347, 2348, 2349, 2350, 2351, 2352, 2353, 2354, 2355, 2356, 2357, 2358, 2359, 2360, 2361, 2362, 2363, 2364, 2365, 2366, 2367, 2368, 2369, 2370, 2371, 2372, 2373, 2374, 2375, 2376, 2377, 2378, 2379, 2380, 2381, 2382, 2383, 2384, 2385, 2386, 2387, 2388, 2389, 2390, 2391, 2392, 2393, 2394, 2395, 2396, 2397, 2398, 2399, 2400, 2401, 2402, 2403, 2404, 2405, 2406, 2407, 2408, 2409, 2410, 2411, 2412, 2413, 2414, 2415, 2416, 2417, 2418, 2419, 2420, 2421, 2422, 2423, 2424, 2425, 2426, 2427, 2428, 2429, 2430, 2431, 2432, 2433, 2434, 2435, 2436, 2437, 2438, 2439, 2440, 2441, 2442, 2443, 2444, 2445, 2446, 2447, 2448, 2449, 2450, 2451, 2452, 2453, 2454, 2455, 2456, 2457, 2458, 2459, 2460, 2461, 2462, 2463, 2464, 2465, 2466, 2467, 2468, 2469, 2470, 2471, 2472, 2473, 2474, 2475, 2476, 2477, 2478, 2479, 2480, 2481, 2482, 2483, 2484, 2485, 2486, 2487, 2488, 2489, 2490, 2491, 2492, 2493, 2494, 2495, 2496, 2497, 2498, 2499, 2500, 2501, 2502, 2503, 2504, 2505, 2506, 2507, 2508, 2509, 2510, 2511, 2512, 2513, 2514, 2515, 2516, 2517, 2518, 2519, 2520, 2521, 2522, 2523, 2524, 2525, 2526, 2527, 2528, 2529, 2530, 2531, 2532, 2533, 2534, 2535, 2536, 2537, 2538, 2539, 2540, 2541, 2542, 2543, 2544, 2545, 2546, 2547, 2548, 2549, 2550, 2551, 2552, 2553, 2554, 2555, 2556, 2557, 2558, 2559, 2560, 2561, 2562, 2563, 2564, 2565, 2566, 2567, 2568, 2569, 2570, 2571, 2572, 2573, 2574, 2575, 2576, 2577, 2578, 2579, 2580, 2581, 2582, 2583, 2584, 2585, 2586, 2587, 2588, 2589, 2590, 2591, 2592, 2593, 2594, 2595, 2596, 2597, 2598, 2599, 2600, 2601, 2602, 2603, 2604, 2605, 2606, 2607, 2608, 2609, 2610, 2611, 2612, 2613, 2614, 2615, 2616, 2617, 2618, 2619, 2620, 2621, 2622, 2623, 2624, 2625, 2626, 2627, 2628, 2629, 2630, 2631, 2632, 2633, 2634, 2635, 2636, 2637, 2638, 2639, 2640, 2641, 2642, 2643, 2644, 2645, 2646, 2647, 2648, 2649, 2650, 2651, 2652, 2653, 2654, 2655, 2656, 2657, 2658, 2659, 2660, 2661, 2662, 2663, 2664, 2665, 2666, 2667, 2668, 2669, 2670, 2671, 2672, 2673, 2674, 2675, 2676, 2677, 2678, 2679, 2680, 26

( )

— — — — —

\_\_\_\_\_

and  $-(-ab)$  is the element that when added to  $-(ab)$  gives 0. This is  $ab$  by definition of  $-(ab)$  and by the uniqueness of an inverse in a group. Thus,  $(-a)(-b) = ab$ .



1.  $\phi(a + b) = \phi(a) + \phi(b)$ ,
2.  $\phi(ab) = \phi(a)\phi(b)$ .

1.  $\phi(a + b) = \phi(a) + \phi(b)$ ,
2.  $\phi(ab) = \phi(a)\phi(b)$ .

## Exercise 1

In Exercises 1 through 6, compute the product in the given ring.

1.  $(12)(16)$  in  $\mathbb{Z}_{24}$
2.  $(16)(3)$  in  $\mathbb{Z}_{32}$
3.  $(11)(-4)$  in  $\mathbb{Z}_{15}$
4.  $(20)(-8)$  in  $\mathbb{Z}_{26}$
5.  $(2, 3)(3, 5)$  in  $\mathbb{Z}_5 \times \mathbb{Z}_9$
6.  $(-3, 5)(2, -4)$  in  $\mathbb{Z}_4 \times \mathbb{Z}_{11}$

# Exercise 1 Solution

1. 0
2. 16
3. 1
4. 22
5. (1, 6)
6. (2, 2)

## Exercise 2

Decide whether the indicated operations of addition and multiplication are defined (closed) on the set, and give a ring structure. If a ring is not formed, tell why this is the case.

1.  $n\mathbb{Z}$  with the usual addition and multiplication
2.  $\mathbb{Z}^+$  with the usual addition and multiplication
3.  $\mathbb{Z} \times \mathbb{Z}$  with addition and multiplication by components
4. The set of all pure imaginary complex numbers  $ri$  for  $r \in \mathbb{R}$  with the usual addition and multiplication

# Exercise 2 Solution

1. Yes,  $n\mathbb{Z}$  for  $n \in \mathbb{Z}^+$  is a commutative ring.
2. No,  $\mathbb{Z}^+$  is not a ring; there is no identity for addition.
3. Yes,  $\mathbb{Z} \times \mathbb{Z}$  is a commutative ring.
4. No,  $\mathbb{R}i$  is not closed under multiplication.

# Exercise 3

Show that  $a^2 - b^2 = (a + b)(a - b)$  for all  $a$  and  $b$  in a ring  $R$  if and only if  $R$  is commutative.

# Exercise 3 Solution

Now  $(a + b)(a - b) = a^2 + ba - ab - b^2$  is equal to  $a^2 - b^2$  if and only if  $ba - ab = 0$ , that is, if and only if  $ba = ab$ . But  $ba = ab$  for all  $a, b \in R$  if and only if  $R$  is commutative.

## 1 Ring (Optional)

## 2 Modular Arithmetic

- Modulo
- Fermat's (Little) Theorem
- Congruences
- Fast Modular Exponentiation
- Chinese Remainder Theorem
- RSA Cryptography

## 3 Q&A



## 2 Modular Arithmetic

## 2 Modular Arithmetic

- **Modulo**
- Fermat's (Little) Theorem
- Congruences
- Fast Modular Exponentiation
- Chinese Remainder Theorem
- RSA Cryptography

### 3 Q&A

# Modular Arithmetic

## Definition

Given  $a, b \in \mathbb{Z}$ ,  $a$  and  $b$  are said to be congruent modulo  $n$ , i.e.,

$$a \equiv b \pmod{n}$$

if  $n \mid b - a$ , i.e.,  $b = a + nk$  for some  $k \in \mathbb{Z}$

# Modular Arithmetic

Determine whether 17 is congruent to 5 modulo 6 and whether 24 and 14 are congruent modulo 6 .

Solution: Because 6 divides  $17 - 5 = 12$ , we see that  $17 \equiv 5 \pmod{6}$ . However, because  $24 - 14 = 10$  is not divisible by 6 , we see that  $24 \not\equiv 14 \pmod{6}$ .

# Theorem 1

Let  $m$  be a positive integer. If  $a \equiv b \pmod{m}$  and  $c \equiv d \pmod{m}$ , then

$$a + c \equiv b + d \pmod{m}$$

and

$$ac \equiv bd \pmod{m}$$

Proof: We use a direct proof. Because  $a \equiv b \pmod{m}$  and  $c \equiv d \pmod{m}$ , there are integers  $s$  and  $t$  with  $b = a + sm$  and  $d = c + tm$ . Hence,

$$b + d = (a + sm) + (c + tm) = (a + c) + m(s + t)$$

and

$$bd = (a + sm)(c + tm) = ac + m(at + cs + stm)$$

Hence,

$$a + c \equiv b + d \pmod{m} \quad \text{and} \quad ac \equiv bd \pmod{m}$$

# Example

Because  $7 \equiv 2 \pmod{5}$  and  $11 \equiv 1 \pmod{5}$ , it follows from Theorem 1 that

$$18 = 7 + 11 \equiv 2 + 1 = 3 \pmod{5}$$

and that

$$77 = 7 \cdot 11 \equiv 2 \cdot 1 = 2 \pmod{5}$$

# Exercise 4

Find each of these values.

a)  $(-133 \bmod 23 + 261 \bmod 23) \bmod 23$

b)  $(457 \bmod 23 \cdot 182 \bmod 23) \bmod 23$

# Exercise 4 Solution

a) Working modulo 23 , we have  $-133 + 261 = 128 \equiv 13$ , so the answer is 13 .

b) Working modulo 23 , we have  $457 \cdot 182 \equiv 20 \cdot 21 = 420 \equiv 6$ .

## 1 Ring (Optional)

## 2 Modular Arithmetic

- Modulo
- Fermat's (Little) Theorem
- Congruences
- Fast Modular Exponentiation
- Chinese Remainder Theorem
- RSA Cryptography

## 3 Q&A



# Fermat's (Little) Theorem

## Theorem I

Given  $a \in \mathbb{Z}$  and  $p \in \mathbb{P}$ , such that  $(a, p) = 1$ , then

$$a^{p-1} \equiv 1 \pmod{p}$$

## Theorem II

Given  $a \in \mathbb{Z}$  and  $p \in \mathbb{P}$ , then

$$a^p \equiv a \pmod{p}$$

# Fermat's (Little) Theorem

Let us compute the remainder of  $8^{103}$  when divided by 13 . Using Fermat's theorem, we have

$$\begin{aligned} 8^{103} &\equiv (8^{12})^8 (8^7) \equiv (1^8) (8^7) \equiv 8^7 \equiv (-5)^7 \\ &\equiv (25)^3 (-5) \equiv (-1)^3 (-5) \equiv 5 \pmod{13} \end{aligned}$$

# Exercise 5

Show that  $2^{11,213} - 1$  is not divisible by 11 .

# Exercise 5 Solution

By Fermat's theorem,  $2^{10} \equiv 1 \pmod{11}$ , so

$$\begin{aligned} 2^{11,213} - 1 &\equiv \left[ (2^{10})^{1,121} \cdot 2^3 \right] - 1 \equiv [1^{1,121} \cdot 2^3] - 1 \\ &\equiv 2^3 - 1 \equiv 8 - 1 \equiv 7 \pmod{11} \end{aligned}$$

# Euler's Theorem

## Theorem

For  $m \in \mathbb{N} \setminus \{0\}$  and  $a \in \mathbb{Z}$  such that  $\gcd(a, m) = 1$ ,

$$a^{\varphi(m)} \equiv 1 \pmod{m}$$

where  $\varphi(m)$  is the number of invertible integers modulo  $m$ .

# Euler's Theorem

1. Compute  $\varphi(p^2)$  where  $p$  is a prime.
2. Compute  $\varphi(pq)$  where both  $p$  and  $q$  are primes.

# Euler's Theorem

1. All positive integers less than  $p^2$  that are not divisible by  $p$  are relatively prime to  $p$ . Thus we delete from the  $p^2 - 1$  integers less than  $p^2$  the integers  $p, 2p, 3p, \dots, (p-1)p$ . There are  $p-1$  integers deleted, so  $\phi(p^2) = (p^2 - 1) - (p - 1) = p^2 - p$
2. We delete from the  $pq - 1$  integers less than  $pq$  those that are multiples of  $p$  or of  $q$  to obtain those relatively prime to  $pq$ . The multiples of  $p$  are  $p, 2p, 3p, \dots, (q-1)p$  and the multiples of  $q$  are  $q, 2q, 3q, \dots, (p-1)q$ . Thus we delete a total of  $(q-1) + (p-1) = p + q - 2$  elements, so  

$$\phi(pq) = (pq - 1) - (p + q - 2) = pq - p - q + 1 = (p-1)(q-1).$$

## 1 Ring (Optional)

## 2 Modular Arithmetic

- Modulo
- Fermat's (Little) Theorem
- **Congruences**
- Fast Modular Exponentiation
- Chinese Remainder Theorem
- RSA Cryptography

## 3 Q&A



# Congruences

A congruence of the form

$$ax \equiv b \pmod{m},$$

where  $m$  is a positive integer,  $a$  and  $b$  are integers, and  $x$  is a variable, is called a linear congruence. Such congruences arise throughout number theory and its applications.

How can we solve the linear congruence  $ax \equiv b \pmod{m}$ , that is, how can we find all integers  $x$  that satisfy this congruence? One method that we will describe uses an integer  $\bar{a}$  such that  $\bar{a}a \equiv 1 \pmod{m}$ , if such an integer exists. Such an integer  $\bar{a}$  is said to be an inverse of  $a$  modulo  $m$ . Theorem 1 guarantees that an inverse of  $a$  modulo  $m$  exists whenever  $a$  and  $m$  are relatively prime.

# Congruences

Let  $d$  be the gcd of positive integers  $a$  and  $m$ . The congruence  $ax \equiv b \pmod{m}$  has a solution if and only if  $d$  divides  $b$ . When this is the case, the solutions are the integers in exactly  $d$  distinct residue classes modulo  $m$ .

# Congruences

Find all solutions of the congruence  $12x \equiv 27 \pmod{18}$ .

Solution: The gcd of 12 and 18 is 6 , and 6 is not a divisor of 27.

Thus by the preceding theorem, there are no solutions.

# Solving Congruences

What are the solutions of the linear congruence  
 $101x \equiv 583 \pmod{4620}$ ?

# Solving Congruences

1. The gcd of 101 and 4620 is 1 , and 1 is a divisor of 583. Thus by the preceding theorem, there is a solution.

The steps used by the Euclidean algorithm to find  $\gcd(101, 4620)$  are

$$101 = 1 \cdot 75 + 26$$

$$75 = 2 \cdot 26 + 23$$

$$26 = 1 \cdot 23 + 3$$

$$23 = 7 \cdot 3 + 2$$

$$3 = 1 \cdot 2 + 1$$

$$2 = 2 \cdot 1$$

In each step we eliminate the remainder by expressing it as a linear combination of the divisor and the dividend. We obtain

$$\begin{aligned} 1 &= 3 - 1 \cdot 2 \\ &= 3 - 1 \cdot (23 - 7 \cdot 3) = -1 \cdot 23 + 8 \cdot 3 \\ &= -1 \cdot 23 + 8 \cdot (26 - 1 \cdot 23) = 8 \cdot 26 - 9 \cdot 23 \\ &= 8 \cdot 26 - 9 \cdot (75 - 2 \cdot 26) = -9 \cdot 75 + 26 \cdot 26 \\ &= -9 \cdot 75 + 26 \cdot (101 - 1 \cdot 75) = 26 \cdot 101 - 35 \cdot 75 \\ &= 26 \cdot 101 - 35 \cdot (4620 - 45 \cdot 101) = -35 \cdot 4620 + 1601 \cdot 101 \end{aligned}$$

# Solving Congruences

4. 1601 is an inverse of 101 modulo 4620.



# Solving Congruences

5. Multiplying both sides of the congruence by 1601 shows that  $1601 \cdot 101x \equiv 1601 \cdot 583 \pmod{4620}$ . Because  $161701 \equiv 1 \pmod{4620}$  and  $933383 \equiv 143 \pmod{4620}$ , it follows that if  $x$  is a solution, then  $x \equiv 143 \pmod{4620}$ .

# Solving Congruences

5. Multiplying both sides of the congruence by 1601 shows that  $1601 \cdot 101x \equiv 1601 \cdot 583 \pmod{4620}$ . Because  $161701 \equiv 1 \pmod{4620}$  and  $933383 \equiv 143 \pmod{4620}$ , it follows that if  $x$  is a solution, then  $x \equiv 143 \pmod{4620}$ .

## 1 Ring (Optional)

## 2 Modular Arithmetic

- Modulo
- Fermat's (Little) Theorem
- Congruences
- **Fast Modular Exponentiation**
- Chinese Remainder Theorem
- RSA Cryptography

## 3 Q&A

# Fast Modular Exponentiation

In cryptography it is important to be able to find  $b^n \bmod m$  efficiently, where  $b$ ,  $n$ , and  $m$  are large integers. It is impractical to first compute  $b^n$  and then find its remainder when divided by  $m$  because  $b^n$  will be a huge number. Instead, we can use an algorithm that employs the binary expansion of the exponent  $n$ . Before we present this algorithm, we illustrate its basic idea. We will explain how to use the binary expansion of  $n$ , say  $n = (a_{k-1} \dots a_1 a_0)_2$ , to compute  $b^n$ . First, note that

$$b^n = b^{a_{k-1} \cdot 2^{k-1} + \dots + a_1 \cdot 2 + a_0} = b^{a_{k-1} \cdot 2^{k-1}} \dots b^{a_1 \cdot 2} \cdot b^{a_0}$$

# Fast Modular Exponentiation

This shows that to compute  $b^n$ , we need only compute the values of  $b, b^2, (b^2)^2 = b^4, (b^4)^2 = b^8, \dots, b^{2^k}$ . Once we have these values, we multiply the terms  $b^{2^j}$  in this list, where  $a_j = 1$ . (For efficiency, after multiplying by each term, we reduce the result modulo  $m$ .) This gives us  $b^n$ . For example, to compute  $3^{11}$  we first note that  $11 = (1011)_2$ , so that  $3^{11} = 3^8 3^2 3^1$ . By successively squaring, we find that  $3^2 = 9, 3^4 = 9^2 = 81$ , and  $3^8 = (81)^2 = 6561$ . Consequently,  $3^{11} = 3^8 3^2 3^1 = 6561 \cdot 9 \cdot 3 = 177,147$ .

The algorithm successively finds  $b \bmod m, b^2 \bmod m, b^4 \bmod m, \dots, b^{2^{k-1}} \bmod m$  and multiplies together those terms  $b^{2^j} \bmod m$  where  $a_j = 1$ , finding the remainder of the product when divided by  $m$  after each multiplication.

# Example

$$2^{2021} \bmod 2021$$

# Example

Consider  $2^{2021} \bmod 2021$ . We first note that the binary representation of 2021 is

$$2021 = 2^{10} + 2^9 + 2^8 + 2^7 + 2^6 + 2^5 + 2^2 + 2^0.$$

Then we know that  $2^{2021} = 2^{2^{10}+2^9+2^8+2^7+2^6+2^5+2^2+2^0} = 2^{2^{10}} \times 2^{2^9} \times 2^{2^8} \times 2^{2^7} \times 2^{2^6} \times 2^{2^5} \times 2^{2^2} \times 2^{2^0}$

Calculating that

$$2^1 \equiv 2 \pmod{2021}$$

$$2^{2^2} \equiv 16 \pmod{2021}$$

$$2^{2^5} \equiv 747 \pmod{2021}$$

$$2^{2^6} \equiv 213 \pmod{2021}$$

$$2^{2^7} \equiv 907 \pmod{2021}$$

$$2^{2^8} \equiv 102 \pmod{2021}$$

$$2^{2^9} \equiv 299 \pmod{2021}$$

$$2^{2^{10}} \equiv 477 \pmod{2021}$$

$$2^{2021} \equiv 477 \times 299 \times 102 \times 907 \times 213 \times 747 \times 16 \times 2 \equiv 1322 \pmod{2021}$$



## 1 Ring (Optional)

## 2 Modular Arithmetic

- Modulo
- Fermat's (Little) Theorem
- Congruences
- Fast Modular Exponentiation
- Chinese Remainder Theorem
- RSA Cryptography

## 3 Q&A

$$x \equiv 2 \pmod{7}$$
$$105 \equiv 0 \pmod{3}, 105 \equiv 0 \pmod{5}, 105 \equiv 0 \pmod{7}$$

$$x \equiv 3 \pmod{8}$$

$$x \equiv 1 \pmod{15}$$

$$x \equiv 11 \pmod{20}$$

$$x \equiv 1 \pmod{3}$$

$$x \equiv 11 \pmod{4}$$

$$x \equiv 11 \pmod{5}$$

$$x \equiv 1 \pmod{5}$$

$$x \equiv 1 \pmod{3}$$

$$x \equiv 5 \pmod{6}$$

$$x \equiv 1 \pmod{15}$$

$$x \equiv 1 \pmod{15}$$

— 20 — (mod 100)

$$x \equiv -29 \pmod{120}$$



We apply the technique of the Chinese Remainder Theorem with

$$a_1 = 6, \quad a_2 = 13, \quad a_3 = 9, \quad a_4 = 19,$$

## Exercise 6 Solution

We compute

$$z_1 = m/m_1 = m_2 m_3 m_4 = 16 \cdot 21 \cdot 25 = 8400$$

$$z_2 = m/m_2 = m_1 m_3 m_4 = 11 \cdot 21 \cdot 25 = 5775$$

$$z_3 = m/m_3 = m_1 m_2 m_4 = 11 \cdot 16 \cdot 25 = 4400$$

$$z_4 = m/m_4 = m_1 m_3 m_3 = 11 \cdot 16 \cdot 21 = 3696$$

$$y_1 \equiv z_1^{-1} \pmod{m_1} \equiv 8400^{-1} \pmod{11} \equiv 7^{-1} \pmod{11} \equiv 8 \pmod{11}$$

$$y_2 \equiv z_2^{-1} \pmod{m_2} \equiv 5775^{-1} \pmod{16} \equiv 15^{-1} \pmod{16} \equiv 15 \pmod{16}$$

$$y_3 \equiv z_3^{-1} \pmod{m_3} \equiv 4400^{-1} \pmod{21} \equiv 11^{-1} \pmod{21} \equiv 2 \pmod{21}$$

$$y_4 \equiv z_4^{-1} \pmod{m_4} \equiv 3696^{-1} \pmod{25} \equiv 21^{-1} \pmod{25} \equiv 6 \pmod{25}$$

$$w_1 \equiv y_1 z_1 \pmod{m} \equiv 8 \cdot 8400 \pmod{92400} \equiv 67200 \pmod{92400}$$

$$w_2 \equiv y_2 z_2 \pmod{m} \equiv 15 \cdot 5775 \pmod{92400} \equiv 86625 \pmod{92400}$$

$$w_3 \equiv y_3 z_3 \pmod{m} \equiv 2 \cdot 4400 \pmod{92400} \equiv 8800 \pmod{92400}$$

$$w_4 \equiv y_4 z_4 \pmod{m} \equiv 6 \cdot 3696 \pmod{92400} \equiv 22176 \pmod{92400}$$



$$x \equiv a_1 w_1 + a_2 w_2 + a_3 w_3 + a_4 w_4 \pmod{92400}$$

$$\equiv 6 \cdot 67200 + 13 \cdot 86625 + 9 \cdot 8800 + 19 \cdot 22176 \pmod{92400}$$

$$\equiv 2029869 \pmod{92400}$$

$$\equiv 51669 \pmod{92400}$$

## 2

-

Trapdoor Function: Want to find a (bijective) trapdoor function

- Easy to compute.
- HARD to invert.
- Unless one has the secret key.

- Easy to compute.
- HARD to invert.
- Unless one has the secret key.

- $$d(y) = y^D = x^{ED} \equiv x \pmod{n}, \quad \text{e.g., } d(y) = y^{191} \pmod{323}$$



$$1980 = 97 \times 20 + 40$$

$$6 = 5 \times 1 + 1$$

$$\begin{aligned} 1 &= 6 - 5 \\ &= 6 - (17 - 6 \times 2) = 6 \times 3 - 17 \\ &= (40 - 17 \times 2) \times 3 - 17 = 40 \times 3 - 17 \times 7 \\ &= 40 \times 3 - (97 - 40 \times 2) \times 7 = 40 \times 17 - 97 \times 7 \\ &= (1980 - 97 \times 20) \times 17 - 97 \times 7 \\ &= 1980 \times 17 - 97 \times 347 \end{aligned}$$

Then



$$279^{2^{10}} \equiv 1054^2 \equiv -279 \pmod{2077}$$

Hence

$$\begin{aligned} 279^{1871} &\equiv 279^{2^0} + 2^5 + 2^6 + 2^9 + 2^{10} \\ &\equiv 279^{2^0} \cdot 279^{2^5} \cdot 279^{2^6} \cdot 279^{2^9} \cdot 279^{2^{10}} \\ &\equiv (279)(341)(-31)(1054)(-279) \\ &\equiv (-403)(-31)(1054)(-279) \\ &\equiv (31)(1054)(-279) \\ &\equiv (-558)(-279) \\ &\equiv 1984 \pmod{2077} \end{aligned}$$

- Modulo
- Fermat's (Little) Theorem
- Congruences
- Fast Modular Exponentiation
- Chinese Remainder Theorem
- RSA Cryptography

## 3

# Q&A

# Q&A