

---

# Chaotic Mixing in a Torus Map

Jean-Luc Thiffeault

Department of Mathematics

Imperial College London

with

Steve Childress

Courant Institute of Mathematical Sciences

New York University

<http://www.ma.imperial.ac.uk/~jeanluc>

# Experiment of Rothstein et al. (1999)

---





Regular array of magnets



[Rothstein, Henry, and Gollub, Nature **401**, 770 (1999)]

# Persistent Pattern

Disordered array ( $i = 2, 20, 50, 50.5$ )



# Local vs Global Regimes of Mixing

---

## Local theory:

- Based on distribution of Lyapunov exponents.

# Local vs Global Regimes of Mixing

---

## Local theory:

- Based on distribution of Lyapunov exponents.
- Applies if initial scale of concentration small.

# Local vs Global Regimes of Mixing

---

## Local theory:

- Based on distribution of Lyapunov exponents.
- Applies if initial scale of concentration small.

- [Antonsen et al., Phys. Fluids (1996)]  
[Balkovsky and Fouxon, PRE (1999)]  
[Son, PRE (1999)]

Average over angles  
Statistical model  
Statistical model

# Local vs Global Regimes of Mixing

---

## Local theory:

- Based on distribution of Lyapunov exponents.
- Applies if initial scale of concentration small.

- [Antonsen et al., Phys. Fluids (1996)]  
[Balkovsky and Fouxon, PRE (1999)]  
[Son, PRE (1999)]

Average over angles  
Statistical model  
Statistical model

## Global theory:

- Eigenfunction of advection–diffusion operator.

# Local vs Global Regimes of Mixing

---

## Local theory:

- Based on distribution of Lyapunov exponents.
- Applies if initial scale of concentration small.

- [Antonsen et al., Phys. Fluids (1996)]  
[Balkovsky and Fouxon, PRE (1999)]  
[Son, PRE (1999)]

Average over angles  
Statistical model  
Statistical model

## Global theory:

- Eigenfunction of advection–diffusion operator.
- Applies if initial scale large, or for inverse cascade.



# Local vs Global Regimes of Mixing

---

## Local theory:

- Based on distribution of Lyapunov exponents.
- Applies if initial scale of concentration small.
- [Antonsen et al., Phys. Fluids (1996)]      Average over angles  
[Balkovsky and Fouxon, PRE (1999)]      Statistical model  
[Son, PRE (1999)]      Statistical model

## Global theory:

- Eigenfunction of advection–diffusion operator.
- Applies if initial scale large, or for inverse cascade.
- [Pierrehumbert, Chaos Sol. Frac. (1994)]      Strange eigenmode  
[Fereday et al., Wonhas and Vassilicos, PRE (2002)]      Baker's map  
[Sukhatme and Pierrehumbert, PRE (2002)]      Unified description

# Local vs Global Regimes of Mixing

---

## Local theory:

- Based on distribution of Lyapunov exponents.
- Applies if initial scale of concentration small.

- [Antonsen et al., Phys. Fluids (1996)]  
[Balkovsky and Fouxon, PRE (1999)]  
[Son, PRE (1999)]

Average over angles  
Statistical model  
Statistical model

## Global theory:

- Eigenfunction of advection–diffusion operator.
- Applies if initial scale large, or for inverse cascade.
- Today: Focus on Global theory.

# Local vs Global Regimes of Mixing

---

## Local theory:

- Based on distribution of Lyapunov exponents.
- Applies if initial scale of concentration small.

- [Antonsen et al., Phys. Fluids (1996)]  
[Balkovsky and Fouxon, PRE (1999)]  
[Son, PRE (1999)]

Average over angles  
Statistical model  
Statistical model

## Global theory:

- Eigenfunction of advection–diffusion operator.
- Applies if initial scale large, or for inverse cascade.
- Today: Focus on Global theory.
- Map allows analytical results (enough said!).

# The Map

---

We consider a diffeomorphism of the 2-torus  $\mathbb{T}^2 = [0, 1]^2$ ,

$$\mathcal{M}(\mathbf{x}) = \mathbb{M} \cdot \mathbf{x} + \phi(\mathbf{x}),$$

where

$$\mathbb{M} = \begin{pmatrix} 2 & 1 \\ 1 & 1 \end{pmatrix}; \quad \phi(\mathbf{x}) = \frac{K}{2\pi} \begin{pmatrix} \sin 2\pi x_1 \\ \sin 2\pi x_1 \end{pmatrix};$$

$\mathbb{M} \cdot \mathbf{x}$  is the **Arnold cat map**.

The map  $\mathcal{M}$  is **area-preserving** and **chaotic**.

For  $K = 0$  the stretching of phase-space elements is **uniform in space** (homogeneous). For small  $K$  it is **hyperbolic**.

# Advection and Diffusion

Iterate the map and apply the **heat operator** to a scalar field (**which we call temperature for concreteness**) distribution  $\theta^{(i-1)}(\boldsymbol{x})$ ,

$$\theta^{(i)}(\boldsymbol{x}) = \mathcal{H}_\epsilon \theta^{(i-1)}(\mathcal{M}^{-1}(\boldsymbol{x}))$$

where  $\epsilon$  is the **diffusivity**, with the **heat operator**  $\mathcal{H}_\epsilon$  and **kernel**  $h_\epsilon$

$$\begin{aligned}\mathcal{H}_\epsilon \theta(\boldsymbol{x}) &:= \int_{\mathbb{T}^2} h_\epsilon(\boldsymbol{x} - \boldsymbol{y}) \theta(\boldsymbol{y}) \, d\boldsymbol{y}; \\ h_\epsilon(\boldsymbol{x}) &= \sum_{\boldsymbol{k}} \exp(2\pi i \boldsymbol{k} \cdot \boldsymbol{x} - \boldsymbol{k}^2 \epsilon).\end{aligned}$$

In other words: **advect** instantaneously and then **diffuse** for one unit of time.

# Transfer Matrix

Fourier expand  $\theta^{(i)}(\mathbf{x})$ ,

$$\theta^{(i)}(\mathbf{x}) = \sum_{\mathbf{k}} \hat{\theta}_{\mathbf{k}}^{(i)} e^{2\pi i \mathbf{k} \cdot \mathbf{x}}.$$

The effect of advection and diffusion becomes

$$\hat{\theta}^{(i)}(\mathbf{x}) = \sum_{\mathbf{q}} \mathbb{T}_{\mathbf{k}\mathbf{q}} \hat{\theta}_{\mathbf{q}}^{(i-1)},$$

with the [transfer matrix](#),

$$\begin{aligned} \mathbb{T}_{\mathbf{k}\mathbf{q}} &:= \int_{\mathbb{T}^2} \exp \left( 2\pi i (\mathbf{q} \cdot \mathbf{x} - \mathbf{k} \cdot \mathcal{M}(\mathbf{x})) - \epsilon \mathbf{q}^2 \right) d\mathbf{x}, \\ &= e^{-\epsilon \mathbf{q}^2} \delta_{0, Q_2} i^{Q_1} J_{Q_1} \left( (k_1 + k_2) K \right), \quad \mathbf{Q} := \mathbf{k} \cdot \mathbb{M} - \mathbf{q}, \end{aligned}$$

where the  $J_Q$  are the Bessel functions of the first kind.

# Variance: A measure of mixing

---

In the absence of diffusion ( $\epsilon = 0$ ), the **variance**  $\sigma^{(i)}$

$$\sigma^{(i)} := \int_{\mathbb{T}^2} |\theta^{(i)}(\mathbf{x})|^2 d\mathbf{x} = \sum_{\mathbf{k}} \sigma_{\mathbf{k}}^{(i)}, \quad \sigma_{\mathbf{k}}^{(i)} := |\hat{\theta}_{\mathbf{k}}^{(i)}|^2$$

is **preserved**. (We assume the spatial mean of  $\theta$  is zero.)

For  $\epsilon > 0$  the variance **decays**.

We consider the case  $\epsilon \ll 1$ , of greatest practical interest.

# Variance: A measure of mixing

---

In the absence of diffusion ( $\epsilon = 0$ ), the **variance**  $\sigma^{(i)}$

$$\sigma^{(i)} := \int_{\mathbb{T}^2} |\theta^{(i)}(\mathbf{x})|^2 d\mathbf{x} = \sum_{\mathbf{k}} \sigma_{\mathbf{k}}^{(i)}, \quad \sigma_{\mathbf{k}}^{(i)} := |\hat{\theta}_{\mathbf{k}}^{(i)}|^2$$

is **preserved**. (We assume the spatial mean of  $\theta$  is zero.)

For  $\epsilon > 0$  the variance **decays**.

We consider the case  $\epsilon \ll 1$ , of greatest practical interest.

**Three phases:**

- The variance is initially **constant**;



# Variance: A measure of mixing

---

In the absence of diffusion ( $\epsilon = 0$ ), the **variance**  $\sigma^{(i)}$

$$\sigma^{(i)} := \int_{\mathbb{T}^2} |\theta^{(i)}(\mathbf{x})|^2 d\mathbf{x} = \sum_{\mathbf{k}} \sigma_{\mathbf{k}}^{(i)}, \quad \sigma_{\mathbf{k}}^{(i)} := |\hat{\theta}_{\mathbf{k}}^{(i)}|^2$$

is **preserved**. (We assume the spatial mean of  $\theta$  is zero.)

For  $\epsilon > 0$  the variance **decays**.

We consider the case  $\epsilon \ll 1$ , of greatest practical interest.

**Three phases:**

- The variance is initially **constant**;
- It then undergoes a rapid **superexponential** decay;

# Variance: A measure of mixing

In the absence of diffusion ( $\epsilon = 0$ ), the **variance**  $\sigma^{(i)}$

$$\sigma^{(i)} := \int_{\mathbb{T}^2} |\theta^{(i)}(\mathbf{x})|^2 d\mathbf{x} = \sum_{\mathbf{k}} \sigma_{\mathbf{k}}^{(i)}, \quad \sigma_{\mathbf{k}}^{(i)} := |\hat{\theta}_{\mathbf{k}}^{(i)}|^2$$

is **preserved**. (We assume the spatial mean of  $\theta$  is zero.)


For  $\epsilon > 0$  the variance **decays**.

We consider the case  $\epsilon \ll 1$ , of greatest practical interest.

**Three phases:**

- The variance is initially **constant**;
- It then undergoes a rapid **superexponential** decay;
- $\theta^{(i)}$  settles into an eigenfunction of the A–D operator that sets the **exponential** decay rate.

# Decay of Variance



## Constant (**Stirring**) Phase

---

- Initially, the variance is essentially constant because of the minuscule diffusivity.

## Constant (**Stirring**) Phase

---

- Initially, the variance is essentially constant because of the minuscule diffusivity.
- However, there is a cascade of the variance to larger wavenumbers under the action of  $\mathbb{M}^{-1}$  in the map. (Neglect  $K$  term.)

## Constant (**Stirring**) Phase

---

- Initially, the variance is essentially constant because of the minuscule diffusivity.
- However, there is a cascade of the variance to larger wavenumbers under the action of  $\mathbb{M}^{-1}$  in the map. (Neglect  $K$  term.)
- This is the well-known “filamentation” effect in chaotic flows: the stretching and folding action of the flow causes rapid variation of the temperature across the folds.

## Constant (Stirring) Phase


---

- Initially, the variance is essentially constant because of the minuscule diffusivity.
- However, there is a cascade of the variance to larger wavenumbers under the action of  $\mathbb{M}^{-1}$  in the map. (Neglect  $K$  term.)
- This is the well-known “filamentation” effect in chaotic flows: the stretching and folding action of the flow causes rapid variation of the temperature across the folds.
- Can no longer neglect diffusion after a number of iterations

$$i_1 \simeq 1 + (\log \epsilon^{-1} / \log \Lambda^2) \simeq 6 \quad \text{for } \epsilon = 10^{-5},$$

where  $\Lambda = (3 + \sqrt{5})/2$  is the largest eigenvalue of  $\mathbb{M}^{-1}$ .

# Variance: 5 iterations for $K = 0.3$ and $\epsilon = 10^{-3}$





# Superexponential Phase

---

For small  $K$  and  $\mathbf{k}$ , we have  $J_0((k_1 + k_2)K) \gg J_1((k_1 + k_2)K)$ , so we set  $K = 0$  and retain **only** the  $Q_1 = 0$  term in the transfer matrix,

$$\mathbb{T}_{\mathbf{k}\mathbf{q}} = e^{-\epsilon \mathbf{q}^2} \delta_{\mathbf{0},\mathbf{Q}} + \mathcal{O}((k_1 + k_2)^2 K^2) ;$$

The **nonvanishing** matrix elements of  $\mathbb{T}$  have  $\mathbf{k} = \mathbf{q} \cdot \mathbb{M}^{-1}$ .

# Superexponential Phase

For small  $K$  and  $\mathbf{k}$ , we have  $J_0((k_1 + k_2)K) \gg J_1((k_1 + k_2)K)$ , so we set  $K = 0$  and retain **only** the  $Q_1 = 0$  term in the transfer matrix,

$$\mathbb{T}_{\mathbf{k}q} = e^{-\epsilon q^2} \delta_{\mathbf{0},Q} + \mathcal{O}((k_1 + k_2)^2 K^2) ;$$

The **nonvanishing** matrix elements of  $\mathbb{T}$  have  $\mathbf{k} = \mathbf{q} \cdot \mathbb{M}^{-1}$ .

If initially the variance is concentrated in a single wavenumber  $\mathbf{q}_0$ , then after one iteration it will all be in  $\mathbf{q}_0 \cdot \mathbb{M}^{-1}$ , after two in  $\mathbf{q}_0 \cdot \mathbb{M}^{-2}$ , etc.

The **length of  $\mathbf{q}$**  is multiplied by  **$\Lambda$**  at each iteration.

# Superexponential Phase

For small  $K$  and  $\mathbf{k}$ , we have  $J_0((k_1 + k_2)K) \gg J_1((k_1 + k_2)K)$ , so we set  $K = 0$  and retain **only** the  $Q_1 = 0$  term in the transfer matrix,

$$\mathbb{T}_{\mathbf{k}\mathbf{q}} = e^{-\epsilon \mathbf{q}^2} \delta_{\mathbf{0},\mathbf{Q}} + \mathcal{O}((k_1 + k_2)^2 K^2) ;$$

The **nonvanishing** matrix elements of  $\mathbb{T}$  have  $\mathbf{k} = \mathbf{q} \cdot \mathbb{M}^{-1}$ .

If initially the variance is concentrated in a single wavenumber  $\mathbf{q}_0$ , then after one iteration it will all be in  $\mathbf{q}_0 \cdot \mathbb{M}^{-1}$ , after two in  $\mathbf{q}_0 \cdot \mathbb{M}^{-2}$ , etc.

The **length of  $\mathbf{q}$**  is multiplied by  $\Lambda$  at each iteration.

But each at each step the **variance** is multiplied by the diffusive decay factor  $\exp(-\epsilon \mathbf{q}^2)$ , with  $\mathbf{q}$  getting exponentially larger; the net decay is thus **superexponential**.

# Exponential Phase

---

- In the superexponential phase we completely neglected the effect of the wave term in the map.

# Exponential Phase

---

- In the superexponential phase we completely neglected the effect of the wave term in the map.
- We described the action as a perfect cascade to large wavenumbers, so that the variance was irrevocably moved to small scales and dissipated extremely rapidly.

# Exponential Phase

---

- In the superexponential phase we completely neglected the effect of the wave term in the map.
- We described the action as a perfect cascade to large wavenumbers, so that the variance was irrevocably moved to small scales and dissipated extremely rapidly.
- There can be no eigenfunction in such a situation, since the mode structure changes completely at each iteration.

# Exponential Phase

---

- In the superexponential phase we completely neglected the effect of the wave term in the map.
- We described the action as a perfect cascade to large wavenumbers, so that the variance was irrevocably moved to small scales and dissipated extremely rapidly.
- There can be no eigenfunction in such a situation, since the mode structure changes completely at each iteration.
- This direct cascade process dominates at first, but it is so efficient that eventually we must examine the effect of the wave term ( $\sin$ ), which is felt through the higher-order Bessel functions in the transfer matrix.

# Exponential Phase

---

- In the superexponential phase we completely neglected the effect of the wave term in the map.
- We described the action as a perfect cascade to large wavenumbers, so that the variance was irrevocably moved to small scales and dissipated extremely rapidly.
- There can be no eigenfunction in such a situation, since the mode structure changes completely at each iteration.
- This direct cascade process dominates at first, but it is so efficient that eventually we must examine the effect of the wave term ( $\sin$ ), which is felt through the higher-order Bessel functions in the transfer matrix.
- Can the wave term lead to the formation of an eigenfunction of the advection–diffusion operator, which would imply exponential decay?



# An Eigenfunction?

---

Recall:

$$\mathbb{T}_{\mathbf{k}\mathbf{q}} = e^{-\epsilon \mathbf{q}^2} \delta_{0, Q_2} i^{Q_1} J_{Q_1} ((k_1 + k_2) K), \quad \mathbf{Q} := \mathbf{k} \cdot \mathbb{M} - \mathbf{q},$$

Consider a matrix element for which  $Q_1 \neq 0$ . This means that the **initial** ( $\mathbf{q}$ ) and **final** ( $\mathbf{k}$ ) wavenumbers connected by that matrix element can **differ** from  $\mathbf{k} \cdot \mathbb{M} = \mathbf{q}$  by  $Q_1$  in their **first component**.

# An Eigenfunction?

Recall:

$$\mathbb{T}_{\mathbf{k}\mathbf{q}} = e^{-\epsilon \mathbf{q}^2} \delta_{0, Q_2} i^{Q_1} J_{Q_1}((k_1 + k_2) K), \quad \mathbf{Q} := \mathbf{k} \cdot \mathbb{M} - \mathbf{q},$$

Consider a matrix element for which  $Q_1 \neq 0$ . This means that the **initial** ( $\mathbf{q}$ ) and **final** ( $\mathbf{k}$ ) wavenumbers connected by that matrix element can **differ** from  $\mathbf{k} \cdot \mathbb{M} = \mathbf{q}$  by  $Q_1$  in their **first component**.

Is it possible for a wavenumber to be **mapped back onto itself** by such a coupling? Seek solutions to


$$(q_1 \ q_2) \cdot \mathbb{M} = (q_1 + Q_1 \ q_2) \implies (q_1 \ q_2) = (0 \ Q_1).$$

The matrix element connecting the  $(0 \ Q_1)$  mode to itself is


$$\mathbb{T}_{(0 \ Q_1), (0 \ Q_1)} = e^{-\epsilon Q_1^2} i^{Q_1} J_{Q_1}(Q_1 K).$$

# Eigenfunction for $K = 0.3$ and $\epsilon = 10^{-3}$

(Renormalised by decay rate)



$i = 25$



$i = 30$

# Decay Rate

---

For small  $K$ , the dominant Bessel function is  $J_1$ , so the decay factor  $\mu^2$  for the variance is given by

$$\mu = \left| \mathbb{T}_{(0\ 1), (0\ 1)} \right| = e^{-\epsilon} J_1(K) = \frac{1}{2}K + \mathcal{O}(\epsilon K, K^2) .$$

Hence, for small  $K$  the **decay rate is limited by the  $(0\ 1)$  mode**. The decay rate is independent of  $\epsilon$  for  $\epsilon \rightarrow 0$ .

# Decay Rate

---

For small  $K$ , the dominant Bessel function is  $J_1$ , so the decay factor  $\mu^2$  for the variance is given by

$$\mu = \left| \mathbb{T}_{(0\ 1), (0\ 1)} \right| = e^{-\epsilon} J_1(K) = \frac{1}{2}K + \mathcal{O}(\epsilon K, K^2) .$$

Hence, for small  $K$  the **decay rate is limited by the  $(0\ 1)$  mode**. The decay rate is independent of  $\epsilon$  for  $\epsilon \rightarrow 0$ .

This is an analogous result to the baker's map [**Fereday et al., Wonhas and Vassilicos, PRE (2002)**], except that here the agreement with numerical results is good for  $K$  quite close to unity.

# Decay Rate

---

For small  $K$ , the dominant Bessel function is  $J_1$ , so the decay factor  $\mu^2$  for the variance is given by


$$\mu = \left| \mathbb{T}_{(0\ 1), (0\ 1)} \right| = e^{-\epsilon} J_1(K) = \frac{1}{2}K + \mathcal{O}(\epsilon K, K^2).$$

Hence, for small  $K$  the **decay rate is limited by the  $(0\ 1)$  mode**. The decay rate is independent of  $\epsilon$  for  $\epsilon \rightarrow 0$ .

This is an analogous result to the baker's map [**Fereday et al., Wonhas and Vassilicos, PRE (2002)**], except that here the agreement with numerical results is good for  $K$  quite close to unity.

This is because in the baker's map the **discontinuity** generates many slowly-decaying harmonics at each step.

# Decay Rate as $\epsilon \rightarrow 0$





# Transition from Superexponential to Exponential

---

- Now that the mechanism of exponential decay is understood, we can go back and describe the condition for **breakdown** of the superexponential solution.



# Transition from Superexponential to Exponential

---

- Now that the mechanism of exponential decay is understood, we can go back and describe the condition for **breakdown** of the superexponential solution.
- The superexponential decay **depletes** the variance very rapidly until all that is left is variance in the exponentially decaying mode  $(0 \ 1)$ .

# Transition from Superexponential to Exponential

---

- Now that the mechanism of exponential decay is understood, we can go back and describe the condition for **breakdown** of the superexponential solution.
- The superexponential decay **depletes** the variance very rapidly until all that is left is variance in the exponentially decaying mode  $(0 \rightarrow 1)$ .
- The superexponential phase thus ends when the variance at large wavenumbers equals that in mode  $(0 \rightarrow 1)$ .

# Transition from Superexponential to Exponential

- Now that the mechanism of exponential decay is understood, we can go back and describe the condition for **breakdown** of the superexponential solution.
- The superexponential decay **depletes** the variance very rapidly until all that is left is variance in the exponentially decaying mode  $(0 \ 1)$ .
- The superexponential phase thus ends when the variance at large wavenumbers equals that in mode  $(0 \ 1)$ .
- Assuming that the variance resides entirely in the  $\mathbf{k}_0 = (0 \ 1)$  mode initially, the condition for breakdown is

$$\mu^{i_2} = \exp \left( -\epsilon \left\| \mathbf{k}_0 \cdot \mathbb{M}^{-(i_2-1)} \right\|^2 \right),$$

where  $\mu^2$  is the decay factor of the variance in  $\mathbf{k}_0$ .

## Transition (continued)

---

After substituting  $\| \mathbf{k}_0 \cdot \mathbb{M}^{-(i_2-1)} \| \simeq \Lambda^{i_2-1}$ , solve numerically for  $i_2$ .

For  $K = 10^{-3}$  and  $\epsilon = 10^{-5}$ , we have  $i_2 \simeq 9.2$ , numerical results.

## Transition (continued)

---

After substituting  $\|\mathbf{k}_0 \cdot \mathbb{M}^{-(i_2-1)}\| \simeq \Lambda^{i_2-1}$ , solve numerically for  $i_2$ .

For  $K = 10^{-3}$  and  $\epsilon = 10^{-5}$ , we have  $i_2 \simeq 9.2$ , numerical results.

For  $\epsilon \ll 1$ , approximate solution given by

$$i_2 \simeq 1 + \log(\epsilon^{-1} \log \mu^{-1}) / \log \Lambda^2,$$

which gives  $i_2 \simeq 8$  for  $K = 10^{-3}$ ,  $\epsilon = 10^{-5}$ .

## Transition (continued)

---

After substituting  $\| \mathbf{k}_0 \cdot \mathbb{M}^{-(i_2-1)} \| \simeq \Lambda^{i_2-1}$ , solve numerically for  $i_2$ .

For  $K = 10^{-3}$  and  $\epsilon = 10^{-5}$ , we have  $i_2 \simeq 9.2$ , numerical results.

For  $\epsilon \ll 1$ , approximate solution given by

$$i_2 \simeq 1 + \log(\epsilon^{-1} \log \mu^{-1}) / \log \Lambda^2,$$

which gives  $i_2 \simeq 8$  for  $K = 10^{-3}$ ,  $\epsilon = 10^{-5}$ .

Subtracting  $i_1 = 1 + \log \epsilon^{-1} / \log \Lambda^2$ , the onset of the superexponential phase, we find the **duration of the phase** is

$$i_2 - i_1 \simeq \frac{\log \log \mu^{-1}}{\log \Lambda^2}.$$

# Duration of Superexponential Phase

---

$$i_2 - i_1 \simeq \frac{\log \log \mu^{-1}}{\log \Lambda^2}$$

- Independent of  $\epsilon$  ;

# Duration of Superexponential Phase

---

$$i_2 - i_1 \simeq \frac{\log \log \mu^{-1}}{\log \Lambda^2}$$

- Independent of  $\epsilon$  ;
- **Very** weak dependence on the decay rate  $\log \mu$  ;



# Duration of Superexponential Phase

---

$$i_2 - i_1 \simeq \frac{\log \log \mu^{-1}}{\log \Lambda^2}$$

- Independent of  $\epsilon$  ;
- **Very** weak dependence on the decay rate  $\log \mu$  ;
- Unless  $\mu$  is small (recall that  $0 < \mu < 1$ ), the superexponential phase is short and may not be noticeable at all, resembling instead a smooth transition;

# Duration of Superexponential Phase

---

$$i_2 - i_1 \simeq \frac{\log \log \mu^{-1}}{\log \Lambda^2}$$

- Independent of  $\epsilon$  ;
- **Very** weak dependence on the decay rate  $\log \mu$  ;
- Unless  $\mu$  is small (recall that  $0 < \mu < 1$ ), the superexponential phase is short and may not be noticeable at all, resembling instead a smooth transition;
- For  $\log \mu^{-1} > 1$  there is **no superexponential phase** at all;

# Duration of Superexponential Phase

---

$$i_2 - i_1 \simeq \frac{\log \log \mu^{-1}}{\log \Lambda^2}$$

- Independent of  $\epsilon$  ;
- **Very** weak dependence on the decay rate  $\log \mu$  ;
- Unless  $\mu$  is small (recall that  $0 < \mu < 1$ ), the superexponential phase is short and may not be noticeable at all, resembling instead a smooth transition;
- For  $\log \mu^{-1} > 1$  there is **no superexponential phase** at all;
- Observed in experiments? There  $\mu$  tends to be closer to unity, so unlikely. But...

# Variance Spectrum of the Eigenfunction

---

- The long-wavelength mode  $(0, 1)$  is the bottleneck that determines the decay rate, for small  $K$ .

# Variance Spectrum of the Eigenfunction

---

- The long-wavelength mode  $(0, 1)$  is the bottleneck that determines the decay rate, for small  $K$ .
- But this dominant mode does not determine the structure of the eigenfunction.

# Variance Spectrum of the Eigenfunction

---

- The long-wavelength mode  $(0 \ 1)$  is the bottleneck that determines the decay rate, for small  $K$ .
- But this dominant mode does not determine the structure of the eigenfunction.
- In fact, a very small amount of the total variance actually resides in that bottleneck mode: the variance is concentrated at small scales.

# Cascade

---

The variance is taken out of the  $(0 \ 1)$  mode by the map: there is a **cascade** to larger wavenumber through the action of  $\mathbb{M}^{-1}$ :

$$(0 \ 1) \rightarrow (-1 \ 2) \rightarrow (-3 \ 5) \rightarrow (-8 \ 13) \rightarrow \dots$$

These become more and more aligned with the stable (contracting) direction of the map.

# Cascade

---

The variance is taken out of the  $(0 \ 1)$  mode by the map: there is a **cascade** to larger wavenumber through the action of  $\mathbb{M}^{-1}$ :

$$(0 \ 1) \rightarrow (-1 \ 2) \rightarrow (-3 \ 5) \rightarrow (-8 \ 13) \rightarrow \dots$$

These become more and more aligned with the stable (contracting) direction of the map.

The **amplitude of the wavenumbers** is multiplied at each step by a factor  $\Lambda = (3 + \sqrt{5})/2 \simeq 2.618$ , the largest eigenvalue of  $\mathbb{M}^{-1}$ .



# Cascade

The variance is taken out of the  $(0 \ 1)$  mode by the map: there is a **cascade** to larger wavenumber through the action of  $\mathbb{M}^{-1}$ :

$$(0 \ 1) \rightarrow (-1 \ 2) \rightarrow (-3 \ 5) \rightarrow (-8 \ 13) \rightarrow \dots$$

These become more and more aligned with the stable (contracting) direction of the map.


The **amplitude of the wavenumbers** is multiplied at each step by a factor  $\Lambda = (3 + \sqrt{5})/2 \simeq 2.618$ , the largest eigenvalue of  $\mathbb{M}^{-1}$ .

But we have seen that the exponential decay rate suggests that the scalar concentration is in an **eigenfunction of the advection–diffusion operator**.

What is going on?

# Eigenfunction: One Iteration

The wavenumbers are mapped back to themselves, with their variance decreased by a uniform factor  $\mu^2 < 1$  (**vertical arrows**). But at the same time the modes are mapped to next one down the cascade following the **diagonal arrows**.



# Eigenfunction and Cascade

---

The decrease in variance for each of the diagonal arrows is diffusive and is given by the factor  $\nu_n = \exp(-2\epsilon \mathbf{k}_n^2)$ .

# Eigenfunction and Cascade

---

The decrease in variance for each of the diagonal arrows is diffusive and is given by the factor  $\nu_n = \exp(-2\epsilon \mathbf{k}_n^2)$ .

If we denote by  $\sigma_n^{(i)} := |\hat{\theta}_{\mathbf{k}_n}|^2$  the variance in mode  $\mathbf{k}_n$  at the  $i$ th iteration, we have

$$\sigma_n^{(i)} = \mu^2 \sigma_n^{(i-1)}, \quad n = 0, 1, \dots,$$

$$\sigma_n^{(i)} = \nu_{n-1} \sigma_{n-1}^{(i-1)}, \quad n = 1, 2, \dots$$

# Eigenfunction and Cascade

The decrease in variance for each of the diagonal arrows is diffusive and is given by the factor  $\nu_n = \exp(-2\epsilon \mathbf{k}_n^2)$ .

If we denote by  $\sigma_n^{(i)} := |\hat{\theta}_{\mathbf{k}_n}|^2$  the variance in mode  $\mathbf{k}_n$  at the  $i$ th iteration, we have

$$\sigma_n^{(i)} = \mu^2 \sigma_n^{(i-1)}, \quad n = 0, 1, \dots,$$

$$\sigma_n^{(i)} = \nu_{n-1} \sigma_{n-1}^{(i-1)}, \quad n = 1, 2, \dots$$

These two recurrences can be combined to give

$$\Sigma_n^{(i)} := \frac{\sigma_n^{(i)}}{\sigma_0^{(i)}} = \frac{\nu_{n-1} \nu_{n-2} \cdots \nu_0}{\mu^{2n}} = \mu^{-2n} \exp\left(-2\epsilon \sum_{m=0}^{n-1} \mathbf{k}_m^2\right),$$

where  $\Sigma_n^{(i)}$  is the **relative variance** in the  $n$ th mode.

## Eigenfunction and Cascade (cont'd)

---

The wavenumber is given by the exponential recursion,

$$\|\mathbf{k}_n\| \simeq \Lambda \|\mathbf{k}_{n-1}\| \quad \Longrightarrow \quad \|\mathbf{k}_n\| \simeq \Lambda^n \|\mathbf{k}_0\| = \Lambda^n .$$

## Eigenfunction and Cascade (cont'd)

---

The wavenumber is given by the exponential recursion,

$$\|\mathbf{k}_n\| \simeq \Lambda \|\mathbf{k}_{n-1}\| \quad \Longrightarrow \quad \|\mathbf{k}_n\| \simeq \Lambda^n \|\mathbf{k}_0\| = \Lambda^n .$$

Solve for  $n = \log \|\mathbf{k}_n\| / \log \Lambda$  and rewrite the relative variance as

$$\Sigma_n^{(i)} \simeq \|\mathbf{k}_n\|^{-2 \log \mu / \log \Lambda} \exp \left( -2\epsilon \mathbf{k}_n^2 / \Lambda^2 \right) ,$$

where we retained only the  $\mathbf{k}_{n-1}^2$  (last) term of the sum.

## Eigenfunction and Cascade (cont'd)

The wavenumber is given by the exponential recursion,

$$\|\mathbf{k}_n\| \simeq \Lambda \|\mathbf{k}_{n-1}\| \implies \|\mathbf{k}_n\| \simeq \Lambda^n \|\mathbf{k}_0\| = \Lambda^n.$$

Solve for  $n = \log \|\mathbf{k}_n\| / \log \Lambda$  and rewrite the relative variance as

$$\Sigma_n^{(i)} \simeq \|\mathbf{k}_n\|^{-2 \log \mu / \log \Lambda} \exp \left( -2\epsilon \mathbf{k}_n^2 / \Lambda^2 \right),$$

where we retained only the  $\mathbf{k}_{n-1}^2$  (last) term of the sum.


Does not (and should not) depend on the iteration number,  $i$ , and depends only on  $n$  through  $\mathbf{k}_n$ . Find

$$\Sigma(k) = k^{2\zeta} \exp \left( -2\epsilon k^2 / \Lambda^2 \right), \quad \zeta := -\log \mu / \log \Lambda,$$

the **spectrum of relative variance**.



# Spectrum of Variance



## Spectrum of Variance (cont'd)

---

$$\Sigma(k) = k^{2\zeta} \exp(-2\epsilon k^2 / \Lambda^2), \quad \zeta := -\log \mu / \log \Lambda$$

- Since  $\mu < 1$  and  $\Lambda > 1$ , we have  $\zeta > 0$ .

## Spectrum of Variance (cont'd)

---

$$\Sigma(k) = k^{2\zeta} \exp(-2\epsilon k^2 / \Lambda^2), \quad \zeta := -\log \mu / \log \Lambda$$

- Since  $\mu < 1$  and  $\Lambda > 1$ , we have  $\zeta > 0$ .
- This implies that there is more variance at the large wavenumbers than at the slowest-decaying mode  $k_0$ .

## Spectrum of Variance (cont'd)

$$\Sigma(k) = k^{2\zeta} \exp(-2\epsilon k^2 / \Lambda^2), \quad \zeta := -\log \mu / \log \Lambda$$

- Since  $\mu < 1$  and  $\Lambda > 1$ , we have  $\zeta > 0$ .
- This implies that there is more variance at the large wavenumbers than at the slowest-decaying mode  $k_0$ .

Find the maximum of  $\Sigma(k)$ ,

$$k_m = \Lambda (\zeta / 2\epsilon)^{1/2}, \quad \Sigma(k_m) = k_m^{2\zeta} e^{-\zeta} = k_m^{2\zeta} \mu^{\log \Lambda}.$$

The peak wavenumber thus scales as  $\epsilon^{-1/2}$ , the same scaling as the dissipation scale.

The relative variance in that peak wavenumber scales as  $\epsilon^{-\zeta}$ .

$k_m$  largest wavenumber that must be included in a numerical calculation to capture the decay of variance correctly (fewer?).

# Conclusions

---

- Three phases of chaotic mixing: constant variance, superexponential decay, exponential decay.

# Conclusions

---

- Three phases of chaotic mixing: constant variance, superexponential decay, exponential decay.
- Large-scale eigenmode dominates exponential phase, as for baker's map. [Fereday et al., Wonhas and Vassilicos, PRE (2002)]

# Conclusions

---

- Three phases of chaotic mixing: constant variance, superexponential decay, exponential decay.
- Large-scale eigenmode dominates exponential phase, as for baker's map. [Fereday et al., Wonhas and Vassilicos, PRE (2002)]
- The spectrum of this eigenmode is determined by a **balance** between the **eigenfunction** property and a **cascade** to large wavenumbers.

# Conclusions

---

- Three phases of chaotic mixing: constant variance, superexponential decay, exponential decay.
- Large-scale eigenmode dominates exponential phase, as for baker's map. [Fereday et al., Wonhas and Vassilicos, PRE (2002)]
- The spectrum of this eigenmode is determined by a **balance** between the **eigenfunction** property and a **cascade** to large wavenumbers.
- For our case of a map with nearly uniform stretching, most of the variance is concentrated at large wavenumbers.



# Conclusions

---

- Three phases of chaotic mixing: constant variance, superexponential decay, exponential decay.
- Large-scale eigenmode dominates exponential phase, as for baker's map. [Fereday et al., Wonhas and Vassilicos, PRE (2002)]
- The spectrum of this eigenmode is determined by a **balance** between the **eigenfunction** property and a **cascade** to large wavenumbers.
- For our case of a map with nearly uniform stretching, most of the variance is concentrated at large wavenumbers.
- The decay rate is unrelated to the Lyapunov exponent or its distribution.

# Conclusions

---

- Three phases of chaotic mixing: constant variance, superexponential decay, exponential decay.
- Large-scale eigenmode dominates exponential phase, as for baker's map. [Fereday et al., Wonhas and Vassilicos, PRE (2002)]
- The spectrum of this eigenmode is determined by a **balance** between the **eigenfunction** property and a **cascade** to large wavenumbers.
- For our case of a map with nearly uniform stretching, most of the variance is concentrated at large wavenumbers.
- The decay rate is unrelated to the Lyapunov exponent or its distribution.
- **Global structure matters!**

# Conclusions

---

- Three phases of chaotic mixing: constant variance, superexponential decay, exponential decay.
- Large-scale eigenmode dominates exponential phase, as for baker's map. [Fereday et al., Wonhas and Vassilicos, PRE (2002)]
- The spectrum of this eigenmode is determined by a **balance** between the **eigenfunction** property and a **cascade** to large wavenumbers.
- For our case of a map with nearly uniform stretching, most of the variance is concentrated at large wavenumbers.
- The decay rate is unrelated to the Lyapunov exponent or its distribution.
- **Global structure matters!**
- Large  $K$ ? Periodic orbits?