
Contents

Introduction	4
Plant anatomy	4
Tissue	4
Types of plant tissues	5
Meristematic Tissues	5
Introduction	5
Characteristics of meristematic tissues	5
Functions of meristematic tissues:	7
Classification of meristematic tissues	7
Origin	7
Position	7
Function	8
Plane of division	8
Promeristem	8
Primary Meristem	9
Secondary Meristem	9
Apical Meristem	10
Intercalary Meristem	11
Lateral Meristem	11
Types of lateral meristem	12
Primary Lateral Meristem	12
Marginal Meristem	12
Meristem on the basis of function	13
Protoderm	13
Procambium	13
Ground meristem	13
Plane of division	14
Rib Meristem	14
Function of rib meristem	14

Mass meristem	14
Function of mass meristem	14
Plate meristem	15
Function of plate meristem	15
Root Apical Meristem <i>R.A.M.</i>	15
Histogen theory for <i>R.A.M.</i>	15
Histogens	15
Components of Histogen	16
Role of histogens	16
Dermatogen of monocot	16
Dermatogen in dicot	16
Features of monocot root	17
Quiescent Center	17
Shoot Apical Meristem	18
Function of shoot apical meristem	18
Tissue Zones	18
Histogen Theory for shoot apical meristem	18
Statement of histogen theory	18
Histogen	18
Rejection of histogen theory	19
Tunica Corpus Theory	19
Statement of tunica corpus theory	19
Tunica	20
Corpus	20
Introduction	20
Origin	20
Structure	21
Structure in plant parts	21
Arrangement in dicot and monocot	21
Anatomical Location	21
Function	22
Types of Ground Tissue System	22
Extra Stelar Region	22

Description of Extra Stelar	22
Hypodermis	22
Cortex	23
Endodermis	23
Pericycle	24
Description of intra stelar	24
Pith	24
Medullary Rays	24
Conjunctive Tissue	25
Vascular tissue	25
Vascular bundles	25
Origin of vascular bundle	25
Location of procambium	25
Parts of vascular bundles	25
Function of vascular bundle	26
Open and closed vascular bundle	26
Open vascular bundles	26
Closed vascular bundle	26
Examples of closed vascular bundles	26
Arrangement of vascular bundles	26
Radial vascular bundles	27
Quantity of radial vascular bundle	27
Conjoint vascular bundles	27
Collateral	27
Example of collateral:	27
Bicollateral	27
Examples of bicollateral xylem	28
Concentric	28
Types of concentric vascular bundles	28
Amphicribal	28
Examples of amphicribal vascular bundles	29

Amphivasal	29
Examples of amphivasaal	29
Types of xylem	29
Protoxylem	29
Character of Wall thickening in protoxylem	30
Metaxylem	30
Character of wall thickneing of metaxylem	30
Division on the basis of position	30
Exarch	30
Location of exacrch xylem	31
Endacrh	31
Loaction of endarch	31
Mesarch	31
Location of mesarch	31
Centacrh	31
Location of centarch	31

Introduction

Plant anatomy

- **Definition:**
 - Plant anatomy is the study of:
 - * internal organization
 - * structure of plants
- **Father of Plant Anatomy:** Nehemiah Grew (N. Grew)

Tissue

- **Definition:** Tissues are :
 - group of similar cells
 - group of dissimilar cells
 - alike in

-
- * origin
 - * function

- **Histology:** **Histology** is the study of tissues.
- **Coined by:** N.Grew
- **Coined in:** 1682

Types of plant tissues

The types of plant tissues are:

- Meristematic Tissues
- Permanent Tissues
- Secretory Tissues

Meristematic Tissues

Introduction

- **Nature of Division:** Meristematic tissues divide continuously.
- **Location:** Growth Regions
 - Tip of plant
 - Apices of
 - * root
 - * shoot
- **Function:** Meristematic tissue helps in
 - growth of the plant.

Characteristics of meristematic tissues

- **Coined by:** C. Nagelli
- **Age:** Immature

-
- **Spacing:** Compact
 - Meristematic tissues have no intercellular spaces.
 - **Shape:** Isodiametric
 - Meristematic tissues have same
 - * length
 - * breadth
 - * height
 - **Geometrical shape:**
 - rounded
 - oval
 - polygonal
 - fusiform
 - * Cells of cambium have fusiform cells.
 - **Type of Nucleus:** multi nucleated
 - **Structure of cell wall:**
 - Cell walls are thin.
 - Cell wall have only primary cell wall.
 - The primary cell wall is cellulosic.
 - Cell wall is elastic.
 - * Cell wall is elastic because of immature cells.
 - **Type of plastids:** Pro plastids are present.
 - Proplastids are the precursor of plastids.
 - **Structure of mitochondria:**
 - Mitochondria have fewer cristae.
 - Mitochondria have shorter cristae.
 - Mitochondrial respiration rate is high.
-

-
- **Storage:** Meristematic cells do not store reserve food.
 - **Examples:**
 - root and shoot apices
 - cambium

Functions of meristematic tissues:

The functions of meristematic tissues are:

- Growth of plant.
 - Meristematic tissue does primary growth.
 - Meristematic tissue does the secondary growth.
 - Formation of new organs
 - * wood
 - * cork

Classification of meristematic tissues

Origin

The meristematic tissues on the basis of origin are:

- Promeristem
- Primary meristem
- Secondary meristem

Position

The meristematic tissues on the basis of position are:

- Apical meristem
- Intercalary meristem
- Lateral meristem

Function

The meristematic tissue on the basis of functions are:

- Protoderm
- Procambium
- Ground meristem

Plane of division

The meristematic tissue on the basis of plane of division are:

- Rib meristem
- Mass meristem
- Plate meristem

Promeristem

- **Origin:** Promeristem has embryonic origin.
- Promeristem is also called
 - Primordial meristem
 - Eumeristem
- **Age:** Promeristem are earliest and youngest.
- **Location:** Promeristem are located at:
 - Extreme tip of young growing root.
 - Extreme tip of young growing shoot.
- **Quantity:** Promeristem contain only few cells.
- **Function:** Promeristem give rise to
 - primary meristem.

Primary Meristem

- **Origin:** Primary meristem is originated from promeristem.
- **Age of formation:** Early in plant
- **Type of Tissues:**
 - active
 - dividing
- **Location:**
 - below promeristem in the root tip
 - below promeristem in the shoot tip
 - intercalary position
- **Function:**
 - Primary meristem gives rise to primary permanent tissue
 - * Primary permanent tissues are formed by differentiation.
 - Primary meristem gives rise to secondary permanent tissue.
 - * Secondary permanent tissue are formed by dedifferentiation.
 - Primary permanent tissues increases the
 - * length and size of root.
 - * length and size of stem.
- **Examples**
 - apical meristem
 - intercalary meristem
 - lateral meristem
 - intra fascicular cambium

Secondary Meristem

- **Time of origin:** Secondary meristem originates

-
- later in the life cycle of plant.
 - **Functions:**
 - Secondary meristem does the secondary growth of plant.
 - Secondary meristem gives rise to secondary permanent tissue.
 - **Examples of secondary permanent tissues:**
 - * secondary cortex
 - * secondary xylem
 - Secondary permanent tissue are originated by dedifferentiation.
 - **Examples:**
 - dicot stem
 - * inter fascicular cambium
 - * cork cambium
 - dicot root
 - * cork cambium
 - * vascular cambium

Apical Meristem

- **Location:**
 - Apices of plant meristem
 - Tip of root
 - Tip of stem
 - Tip of branches
 - Root apex
 - Shoot apex
- **Structural position:**
 - It is present in sub terminal position of root.

-
- * The terminal position of root contains root cap.
 - * Root cap is also termed as calyptragen.
 - It is present at the apical position in shoot.

- **Function:**

- Apical meristem increases the length of the plant body.

Intercalary Meristem

- **Location:**

- between permanent tissues
 - base of leaves
 - nodal regions

- **Function:**

- Intercalary meristem does the primary growth of the plant.

- **Examples:**

- Intercalary meristem are found at the base of pinus leaves.
 - * The intercalary meristem at the base of pinus leaves is basal meristem.
 - Above the sporophytes of anthoceros
 - Above the nodes of bamboo, grasses
 - Below nodes of mint
 - Stem of all monocots
 - Petiole of leaves
 - Leaf sheath of monocot

Lateral Meristem

- **Location:**

-
- Lateral side of stem
 - Lateral side of roots
 - Components of lateral meristem divide periclinally.
 - Periclinal division increases the diameter of an organ.
 - **Function:**
 - Lateral meristem is responsible of growth in thickness of plant body.
 - * Growth in thickness of plant body is secondary growth.

Types of lateral meristem

The types of lateral meristem are:

- Primary Lateral meristem
- Secondary lateral meristem
 - Secondary lateral meristem is also called marginal meristem.

Primary Lateral Meristem

The examples of primary lateral meristem are:

- Intra fascicular cambium
 - Intra fascicular cambium is present within the vascular bundle.
 - Intra fascicular cambium is present in dicot stem.

Marginal Meristem

- Location:
 - Margins of leaf blade
- **Function:**
 - Marginal meristem expand the leaf surface.
 - Marginal meristem do not expand the thickness.

Meristem on the basis of function

- The meristem were classified on the basis of function by Hanberlandt.
- Hanberlandt calssified meristem on the basis of function in 1914 .

Protoderm

- **Location:** Protoderm is the outermost meristematic tissue.

The function of protoderm are:

- Protection from mechanical injury.
- Protoderm gives rise to epidermis layer.

Procambium

- **Location:** Procambium is the innermost meristematic tissue.

The function of procambium are:

- Procambium transports water and nutrition
- Procambium gives rise to vascular tissues.
 - Vascular tissues are xylem and phloem.

Ground meristem

The function of ground meristem are:

- Ground meristem gives rise to the following components in monocot plants:
 - Hypodermis
 - Ground Parenchyma
- Ground meristem gives rise to the following components in dicot plants:
 - Cortex
 - Endodermis
 - Pericycle
 - Pith

Plane of division

The types of meristematic tissues on the basis of plane of division are:

- Rib meristem
- Mass meristem
- Plate meristem

Rib Meristem

- These cells divide only in one plane.
 - The division in only one plane is called anticlinal division.

Function of rib meristem

The function of rib meristem are:

- Rib meristem plays a role in young stem to develop:
 - Pith
 - Cortex

Mass meristem

- These cells divide in three planes.

Examples of mass meristem are:

- Early embryo

Function of mass meristem

The function of mass meristem is:

- Mass meristem plays a role in:
 - early development of endosperm
 - early development of cortex
 - early development of pith

Plate meristem

- Plate meristems divide in two planes.

Function of plate meristem

The function of plate meristem are:

- Plate meristem develops leaves.
 - The development of leaves by plate meristem doesnot increase the thickness.

Root Apical Meristem *R.A.M.*

- Root apical meristem is also known as root apex.
- **Structure:** Root apical meristem has inverted cup like structure.
- **Content:** Root apical meristem is a group of initial cells.
- **Origin:** Root apical meristem is embryonic in origin.
- **Location:**
 - Root apical meristem is present at the sub terminal region of growing root tip.
 - Root apical meristem lies below the root cap.

Function:

- Root apical meristem produces the tissues of primary root.

Histogen theory for *R.A.M.*

- Histogen theory was given by Hanstein.
- Histogen theory was given in 1868 .

Histogens

- The cell initiating region of root apical meristem are called histogens.

Components of Histogen

- The histogens are:
 - dermatogen
 - periblem
 - plerome

Role of histogens

- Histogen forms components present in a mature root.
- The role of histogen is described as:
 - Dermatogen forms the epidermis.
 - Periblem forms the ground tissue.
 - * cortex
 - * endodermis
 - * pith
 - * pericycle
 - Plerome forms the vascular cylinder.
 - * xylem
 - * phloem

Dermatogen of monocot

The following condition is seen in monocot plants:

- Dermatogen generates only the root cap in monocots.
 - The dermatogen of monocot is called calyptragen.

Dermatogen in dicot

The following components are generated by dermatogen in dicot plants:

-
- Dermatogen generates protoderm in dicots.
 - Dermatogen generates root cap in dicots.
 - The dermatogen of dicot is called dermatocalyptrogen.

Features of monocot root

- **Quantity of Xylem:** The quantity of xylem groups vary from 8 to 20.
- **Role of pericycle:** Pericycle gives rise to lateral roots only.
- **Contents:** Cambium is absent.
- **Nature of growth:** There is no secondary growth.
- **Nature of pith:**
 - The pith is larger.
 - The pith is well developed.

Quiescent Center

- **Location:** Quiescent center is found in root promeristem.
- **Discovery:**
 - Quiescent center was discovered by Clowes.
 - Quiescent center was discovered in 1956 – 1958 .
 - Quiescent center was discovered in the root tip of *Zea Mays* .
- **Shape:**
 - Closed meristem
 - * Quiescent center is hemispherical in closed meristem.
 - Open meristem
 - * Quiescent center is disc shaped in open meristem.
- **Nature of cells:** Cells in quiescent center are inactive.
- **Division of cells:** Cells in quiescent center do not divide.

Shoot Apical Meristem

Function of shoot apical meristem

- Shoot apical meristem produces lateral organs.
- The lateral organs produced by shoot apical meristem are:
 - leaves
 - branch
 - flowers

Tissue Zones

The tissue zones of shoot apical meristem are the meristematic tissues divided on the basis of function.

Histogen Theory for shoot apical meristem

- The histogen theory was given by J. Hanstein.
- The histogen theory was given in 1868 .

Statement of histogen theory

- The plant body doesnot originate from a single superficial cells.
- The plant body originates from mass of meristematic cells.
- The meristematic region contain three distinct zones.
- The distinct zones of meristematic region are called histogens.

Histogen

The histogens proposed by Hanstein are:

- **Dermatogen:**
 - *Location:* Dermatogen is the outermost layer.

-
- *Function:* Dermatogen gives rise to the epidermis.
 - **Periblem:**
 - **Location:** Periblem lies inner to dermatogen.
 - **Function:** Periblem gives rise to cortex.
 - **Plerome:**
 - *Location:* Plerome lies at the center.
 - *Function:* Plerome gives rise to
 - * vascular tissue
 - * endodermis

Rejection of histogen theory

The reasons for rejection of histogen theory are:

- There is no strict zonal differentiation between histogens.
- Dermatogen , Periblem and Plerome cannot be distinguished from each other.

Tunica Corpus Theory

- Tunica Corpus Theory was proposed by Schmidst.
- Tunica Corpus Theory was proposed in 1924 .

Statement of tunica corpus theory

- There are two distinct tissue zones in the apical region of the shoot.
- The distinct tissue zones are:
 - Tunica
 - Corpus

Tunica

- **Size of cells:** The cells of tunica are smaller than corpus.
- **Nature of division:** The cells of tunica show anticlinal division.
- **Contents:** Tunica contains one or more peripheral layer of cells.
- **Functions:**
 - Tunica produces :
 - * cortex
 - * endodermis
 - * pericycle
 - * vascular tissue
 - * pith
 - Tunica increases surface area.

Corpus

- **Size of cells:** The cells of corpus are larger than tunica.
- **Contents:** Corpus contains inner mass of cells.
- **Functions:**
 - Corpus produces tissues that are not produced by tunica.
 - Corpus increases the volume.

Introduction

Origin

- **Ground Tissue System** originates from **ground meristem**

Structure

- **Ground Tissue System** *histologically* is made up of:
 - *Parenchyma*
 - *Collenchyma*
 - *Sclerenchyma*

Structure in plant parts

- **Ground** tissue system is present in leaf as:
 - **Pallisade**
 - **Spongy Mesophyll Cells**
- These **tissues** are *parenchymatous* .

Arrangement in dicot and monocot

- **Monocot** system consists of:
 - **Hypodermis**
 - **Ground Parenchyma**
- **Dicot** system consists of:
 - **Hypodermis**
 - **Cortex**
 - **Endodermis**
 - **Pith**
 - **Medullary rays**

Anatomical Location

- **Ground tissue system** is present *inner* to the **epidermis** and **stele** .

Function

- **Ground** tissue system:
 - fills the **gap** in between.
 - provides **mechanical** strength.

Types of Ground Tissue System

Extra Stelar Region

- In **extra** stelar region ground tissue are categorized into:
 - Hypodermis
 - Cortex
 - Endodermis
 - Pericycle
- In **intra** stelar region ground tissue are categorized into:
 - Pith
 - Medullary rays
 - Conjunctive Tissue

Description of Extra Stelar

Hypodermis

- **Hypodermis** lies inner to *epidermis* .
- It is made up of cells of type:
 - **collenchymatous**
 - **sclerenchymatous**
- Hypodermis provides **mechanical strength**.

Cortex

- **Location:** It lies inner to *hypodermis* .
- **Structure:** It is made up of **parenchymatous** cells.
- **Content:**
 - Starch grains
 - Resin ducts
 - Oils
 - Tanins
 - Lactiferous ducts
- **Function:**
 - Storage
 - Photosynthesis
 - Protection
 - Water pumping to inner plant parts.

Endodermis

- **Structure:** It consists of cells of type :
 - * Single Layered
 - * Barrel shaped
 - * **Living**
- **Anatomical Structure:**
 - * Only in **roots** there is presence of:
 - Band of **suberin** .
 - This band is present at the *lateral* and *tangential* wall.
 - This band was discovered by **Caspary** .
 - It was discovered on **1866** .
 - This band is also called **casparian strips** .

– **Function:**

- * Checks the entry of **water**.
- * Checks the entry of **air** in parts of xylem.
- * Acts as *internal* **protective tissue**.

Pericycle

- **Location:** Between **endodermis** and **vascular strands** .
- **Type of cells:**
 - *Stem:* **parenchymatous** or **sclerenchymatous** .
 - *Root:* **parenchymatous**
- **Function:** Acts as **outer vascular** strands.

Description of intra stelar

Pith

- **Types of cells:** Parenchymatous cells without **chlorophyll** .
- **Contents:** *Tanin* and *Crystals* .
- **Function:** *Storage* of food.

Medullary Rays

- **Location:** Between vascular bundles.
- **Anatomical Structure:** *Narrow parenchymatous strips*.
- **Function:**
 - **Meristematic medullary rays:** Formation of interfascicular cambium.
 - **Parenchymatous medullary rays:** Transportation of **food** and **water** from *pith* to *cortex* .

Conjunctive Tissue

- **Background:** *Xylem* and *Phloem* are present in separate *patches* .
- **Location:** Between **patches** of **vascular tissue**.
- **Structure of cells:** Parenchymatous

Vascular tissue

- The other name for vascular tissue is fascicular tissue.
- Vascular tissue is defined as:
 - central cylinder of root and shoot
 - This system is surrounded by ground tissue system
 - The ground tissue system may be pericycle pith
 - Stele is the central cylinder of this structure.

Vascular bundles

- Vascular bundles are tissue system

The location for vascular bundles is: - vascular bundles are distributed in the stele.

Origin of vascular bundle

Vascular bundles are originated from procambium.

Location of procambium

Procambium lies under primary meristem.

Parts of vascular bundles

The parts of vascular bundles are - Xylem - Phloem - Cambium

Function of vascular bundle

The function of vascular bundle are:

- Vascular bundles conduct water from root to leaves
- Vascular bundles conduct nutrients from root to leaves
- These materials such as root and nutrients are carried by xylem
- The vascular tissues translocate food prepared in the leaves
- This translocation is done by pholem.

Open and closed vascular bundle

Open vascular bundles

- The cambium is present between the xylem and pholem.
- This types of vascular bundle is called open vascular bundle.
- The cambium of open vascular bundle forms the sencondary xylem and pholem tissue.

Closed vascular bundle

- There is the absence of cambium in closed vascular bundle
- The plants containing closed vascular bundles don not form the seondary growth

Examples of closed vascular bundles

- Closed vascular bundles is present in monocots.

Arrangement of vascular bundles

The types of vascular bundles on the basis of arrangement are - radial vascular bundles - conjoint vascular bundles

Radial vascular bundles

- There is the presence of xylem and phloem in radial vascular bundle
- Xylem are arranged alternatively in different bundles.
- Phloem are arranged alternatively in different bundles with xylem.
- The arrangement of xylem and phloem is present on different radii.

Quantity of radial vascular bundle

Dicots : - Dicot plants have 2-6 radial vascular bundle - They are also called as di arch to hex arch

Monocots: - Monocot plants have 8-20 radial vascular bundles - They are also called polyarch

Conjoint vascular bundles

Types of conjoint vascular bundles Conjoint vascular bundles are of three type: - collateral - bicollateral - concentric

Collateral

- There is the presence of xylem and phloem.
- Phloem is present in the outer side
- Xylem is present in the inner side

Example of collateral:

- sunflower

Bicollateral

- There is the presence of phloem and xylem
- Phloem is present on both sides of xylem

-
- There is a presence of vascular cambium
 - The anatomy of the vascular cambium is a strip
 - The vascular cambium strip is present at the outer sides of the xylem
 - The vascular cambium strip is present at the inner sides of the xylem

Examples of bicollateral xylem

- Stem of cucurbita
- Solanaceace family

Concentric

Concentric vascular bundle have xylem and phloem The phloem surrounds the xylem or the xylem surrounds the phloem One of them forms the central core The other of them surrounds the central core

Types of concentric vascular bundles

Concentric vascular bundles are of two types

- Amphivasal
- Amphicribal

Amphicribal

- The amphicribal vascular bundles are also called hadrocentric
- The xylem forms the central core
- The phloem surrounds the central core of the xylem

Examples of amphi-ribal vascular bundles

- ferns
- aquatic angiosperms
- dicots

Amphivasal

- The other term for amphivasal is leptocentric
- the phloem forms the central core.
- The xylem surrounds the phloem in amphivasal

Examples of amphivasal

- Dracena
- Yucca

Types of xylem

- The xylem can be divided in two types.
- This division can be done on the basis of development
- The types of xylem are
- Protoxylem
- Metaxylem

Protoxylem

- Protoxylem is the first formed xylem
- The protoxylem contains tracheids and vessels.
- There is the presence of lumens in tracheids and vessels of protoxylem.
- The lumen of tracheids and vessels of proto xylem is narrow.
- This narrowness results in the implication of smaller size of proto xylem

Character of Wall thickening in protoxylem

- The wall thickening character in protoxylem may be
- annular
- spiral

Metaxylem

- Metaxylem are formed later.
- The metaxylem have presence of tracheids and vessels.
- There is the presence of lumen in the tracheids and vessels.
- The lumen in the tracheids and vessels of the metaxylem is larger in size.
- This implies that the metaxylem have a large size

Character of wall thickening of metaxylem

- The wall thickening character of metaxylem can be described as:
- reticulate
- scalariform pitted
- simple
- bordered

Division on the basis of position

Xylem are also divided on the basis of position

The types of xylem on the basis of position are: - exarch - endarch - mesarch - centarch

Exarch

- There are protoxylems and metaxylems in exarch
- The presence of protoxylem is towards the periphery in exarch.
- The presence of metaxylem is towards the center in exarch

Location of exarch xylem

The exarch xylem are present in the - roots of monocot plants - roots of dicot plants

Endarch

- There is the presence of protoxylem and metaxylem at endarch
- The protoxylem are located at the center in endarch.
- The metaxylem are located at the periphery in endarch

Location of endarch

Endarch xylem s are present in - stem of monocot plants - stem of dicot plants

Mesarch

- There is the presence of protoxylem and metaxylem in mesarch
- The protoxylem is situated in between the metaxylem

Location of mesarch

- Mesarch xylem is present in
- Pteridophytes The pteridophytes which have mesarch xylem are ferns

Centarch

- There is the presence of protoxylem and metaxylem in centarch
- The protoxylem are located at the center
- The metaxylem surrounds the protoxylem

Location of centarch

- Centarch is present in pteridophytes
- The pteridophytes having the presence of centarch are sellagenaella