M&S Monthly Colloquium Meeting

Fixed Raters, Random Selection:
A Correlation-Based Framework for Agreement ICCs in Two-Way Mixed-Effects Designs

Tzu-Yao Lin, Francis Tuerlinckx, Sophie Vanbelle

Maastricht University KU LEUVEN

05/10/2026

How to measure reliability?

- Reliability asks whether a score can support the same conclusion across replications of a specified measurement procedure.
- Classical Test Theory (Lord et al., 1968)

$$Y_{ij} = T_{ij} + \varepsilon_{ij}, \quad \sigma_Y^2 = \sigma_T^2 + \sigma_\varepsilon^2$$

- The reliability coefficient and **two interpretations**:

$$\rho = \frac{\sigma_T^2}{\sigma_T^2 + \sigma_\epsilon^2}$$

Correlation Approach

(under the parallel assumptions)

$$\rho = \mathrm{Cor}\left(Y_{ij}, Y_{ij'}\right)$$

The degree of similarity between two replicates.

Variance Partitioning Approach

$$\rho = \frac{\sigma^2_{\text{person}}}{\sigma^2_{\text{person}} + \sigma^2_{\text{error}}}$$

How well can we distinguish targets from error?

Agreement and consistency



	Rater A	Rater B
S_1	20	30
S_2	35	45
S_3	45	55
S_4	67	77
⋮		
S_{10}	80	90

- Would another rater support the same decision?

1. Do the raters give interchangeable scores?

   ⇒ Interrater **Agreement**

2. Do the raters rank patients similarly?

   ⇒ Interrater **Consistency**

McGraw and Wong's (1996) ICCs

Model	Unit	Decision	
		Agreement	Consistency
Two-way random-effects	Single	$\text{ICC(A, 1)} = \frac{\sigma_s^2}{\sigma_s^2+\sigma_R^2+\sigma_\varepsilon^2}$	$\text{ICC(C, 1)} = \frac{\sigma_s^2}{\sigma_s^2+\sigma_\varepsilon^2}$
	Average	$\text{ICC(A, k)} = \frac{\sigma_s^2}{\sigma_s^2+\frac{\sigma_R^2}{k}+\frac{\sigma_\varepsilon^2}{k}}$	$\text{ICC(C, k)} = \frac{\sigma_s^2}{\sigma_s^2+\frac{\sigma_\varepsilon^2}{k}}$
Two-way mixed-effects	Single	$\text{ICC(A, 1)} = \frac{\sigma_s^2}{\sigma_s^2+\theta_R^2+\sigma_\varepsilon^2}$	$\text{ICC(C, 1)} = \frac{\sigma_s^2}{\sigma_s^2+\sigma_\varepsilon^2}$
	Average	$\text{ICC(A, k)} = \frac{\sigma_s^2}{\sigma_s^2+\frac{\theta_R^2}{k}+\frac{\sigma_\varepsilon^2}{k}}$	$\text{ICC(C, k)} = \frac{\sigma_s^2}{\sigma_s^2+\frac{\sigma_\varepsilon^2}{k}}$

$\theta_R^2 = \frac{1}{k-1}\sum R_j^2$: Rater dispersion for fixed rater effects

Two-way random-effects designs

	R_1	R_2	R_3	…	R_k
S_1	x	x	x		x
S_2	x	x	x		x
S_3	x	x	x		x
S_4	x	x	x		x
⋮				⋱	
S_n	x	x	x		x

- Person: $i = 1, \ldots, n$, Rater $j = 1, \ldots, k$
- Two-way random-effects model:


$$Y_{ij} = \mu + S_i + R_j + \varepsilon_{ij}$$

  - $S_i \sim (0, \sigma_S^2)$; $R_j \sim (0, \sigma_R^2)$; $\varepsilon_{ij} \sim (0, \sigma_\varepsilon^2)$
  - Cannot separate the person×rater interaction from residual error.
- Total variance:

$$\mathrm{Var}(Y_{ij}) = \sigma_S^2 + \sigma_R^2 + \sigma_\varepsilon^2$$

Agreement vs. consistency


	Rater	Operational Score
S_1	X	$Y_i = \mu + S_i + R + \varepsilon_i$
S_2	X	
S_3	X	
⋮		
⋮		
S_n	X	



Agreement vs. consistency

	Rater
S_1	X
S_2	X
S_3	X
⋮	
⋮	
S_n	X

Operational Score
Y_i
$=$
μ
$+$
S_i
$+$
R
$+$
ε_i



$$\rho = \frac{\sigma^2_{\text{person}}}{\sigma^2_{\text{person}} + \sigma^2_{\text{error}}}$$

$$\sigma^2_{\text{abs.error}} = \sigma_R^2 + \sigma_\varepsilon^2$$

$$\text{ICC}(A,1) = \frac{\sigma_s^2}{\sigma_s^2 + \sigma_R^2 + \sigma_\varepsilon^2}$$


$$\sigma^2_{\text{rel.error}} = \sigma_\varepsilon^2$$

$$\text{ICC}(C,1) = \frac{\sigma_s^2}{\sigma_s^2 + \sigma_\varepsilon^2}$$

Single vs. average scores

	R_1	$\cdots$	R_k
S_1	X		X
S_2	X		X
S_3	X		X
$\vdots$			
$\vdots$			
S_n	X		X

Operational Score
$\bar{Y}_{i.}$
$=$
μ
$+$
S_i
$+$
$\frac{R_1 + \cdots + R_k}{k}$
$+$
$\frac{\varepsilon_1 + \cdots + \varepsilon_k}{k}$



$$\rho = \frac{\sigma^2_{\text{person}}}{\sigma^2_{\text{person}} + \sigma^2_{\text{error}}}$$

$$\sigma^2_{\text{abs.error}} = (\sigma_R^2 + \sigma_\varepsilon^2)/k$$

$$\text{ICC}(A,k) = \frac{\sigma_s^2}{\sigma_s^2 + (\sigma_R^2 + \sigma_\varepsilon^2)/k}$$

$$\sigma^2_{\text{rel.error}} = \sigma_\varepsilon^2/k$$


$$\text{ICC}(C,k) = \frac{\sigma_s^2}{\sigma_s^2 + \sigma_\varepsilon^2/k}$$

McGraw and Wong's (1996) ICCs

Model	Unit	Decision	
		Agreement	Consistency
Two-way random-effects	Single	ICC(A, 1) $= \mathrm{Cor}(Y_{ij}, Y_{ij'})$	ICC(C, 1) $=???$
	Average	ICC(A, k) $= \mathrm{Cor}(\bar{Y}_{i.}, \bar{Y}_{i.'})$	ICC(C, k) $=???$
Two-way mixed-effects	Single	ICC(A, 1) $= \frac{\sigma_S^2}{\sigma_S^2+\theta_R^2+\sigma_\varepsilon^2}$	ICC(C, 1) $= \frac{\sigma_S^2}{\sigma_S^2+\sigma_\varepsilon^2}$
	Average	ICC(A, k) $= \frac{\sigma_S^2}{\sigma_S^2+\frac{\theta_R^2}{k}+\frac{\sigma_\varepsilon^2}{k}}$	ICC(C, k) $= \frac{\sigma_S^2}{\sigma_S^2+\frac{\sigma_\varepsilon^2}{k}}$

$\theta_R^2 = \frac{1}{k-1}\sum R_j^2$: Rater dispersion for fixed rater effects

Agreement ICC (Shrout & Fleiss, 1979)




Hypothetical replicate

	$R^{(1)}$	$R^{(2)}$	$R^{(3)}$	...
$S^{(1)}$	$Y^{(1)}$			
$S^{(2)}$		$Y^{(2)}$		
$S^{(3)}$			$Y^{(3)}$	
⋮				⋱

	$R'^{(1)}$	$R'^{(2)}$	$R'^{(3)}$	...
$S^{(1)}$	$Y'^{(1)}$			
$S^{(2)}$		$Y'^{(2)}$		
$S^{(3)}$			$Y'^{(3)}$	
⋮				⋱

$$\mathrm{Cor}(Y_{ij}, Y_{ij'}) = \frac{\sigma_s^2}{\sigma_s^2 + \sigma_R^2 + \sigma_\varepsilon^2} = \mathrm{ICC(A,1)}$$

Consistency ICC (Vangeneugden et al., 2005)



Specify two raters all the time

Hypothetical replicate


	R	R	R	…
$S^{(1)}$	$Y^{(1)}$			
$S^{(2)}$		$Y^{(2)}$		
$S^{(3)}$			$Y^{(3)}$	
⋮				⋱

	R'	R'	R'	…
$S^{(1)}$	$Y'^{(1)}$			
$S^{(2)}$		$Y'^{(2)}$		
$S^{(3)}$			$Y'^{(3)}$	
⋮				⋱

$$\mathrm{Cor}(Y_{ij}, Y_{ij'}|j, j') = \frac{\sigma_s^2}{\sigma_s^2 + \sigma_\varepsilon^2} = \mathrm{ICC}(C, 1)$$

McGraw and Wong's (1996) ICCs

Model	Unit	Decision	
		Agreement	Consistency
Two-way random-effects	Single	ICC(A, 1) $= \mathrm{Cor}\left(Y_{ij}, Y_{ij'}\right)$	ICC(C, 1) $= \mathrm{Cor}\left(Y_{ij}, Y_{ij'} \mid j, j'\right)$
	Average	ICC(A, k) $= \mathrm{Cor}\left(\bar{Y}_{i.}, \bar{Y}_{i.'}\right)$	ICC(C, k) $= \mathrm{Cor}\left(\bar{Y}_{i.}, \bar{Y}_{i.'} \mid \forall j, \forall j'\right)$
Two-way mixed-effects	Single	ICC(A, 1) $= \mathrm{Cor}\left(Y_{ij}, Y_{ij'}\right)$	ICC(C, 1) $= \mathrm{Cor}\left(Y_{ij}, Y_{ij'} \mid j, j'\right)$
	Average	ICC(A, k) $= \mathrm{Cor}\left(\bar{Y}_{i.}, \bar{Y}_{i.'}\right)$	ICC(C, k) $= \mathrm{Cor}\left(\bar{Y}_{i.}, \bar{Y}_{i.'} \mid \forall j, \forall j'\right)$



$\theta_R^2 = \frac{1}{k-1}\sum R_j^2$: Rater dispersion for fixed rater effects

Two-way mixed-effects designs

Fixed raters of interest

	R_1	R_2	R_3	…	R_K
S_1	x	x	x		x
S_2	x	x	x		x
S_3	x	x	x		x
S_4	x	x	x		x
⋮				⋱	
S_n	x	x	x		x

- Person: $i = 1, \ldots, n$, Rater $j = 1, \ldots, K$
- Model:

$$Y_{ij} = \mu + S_i + R_j + \varepsilon_{ij}$$

  - $S_i \sim (0, \sigma_S^2)$; $\varepsilon_{ij} \sim (0, \sigma_\varepsilon^2)$
  - $\sum_{j=1}^{K} R_j = 0$, R_j's are unknown constants.
  - Cannot separate the person×rater interaction from residual error.

- Total variance:

$$\mathrm{Var}(Y_{ij}) = \sigma_S^2 + \sigma_\varepsilon^2$$

Fixed-rater agreement paradox (1)

- Rater effects $R_1, \ldots, R_K$ are the fixed, unknown constants, $\sum_{j=1}^{K} R_j = 0$.
  - There is no probability variance,

$$\mathrm{Cor}\left(Y_{ij}, Y_{ij'}\right) = \frac{\sigma_s^2}{\sigma_s^2 + \sigma_\varepsilon^2} = \mathrm{ICC}(C, 1) \qquad \text{when } j \neq j'$$

$$\nparallel$$

$$\mathrm{ICC}(\mathrm{A}, 1) = \frac{\sigma_s^2}{\sigma_s^2 + \theta_R^2 + \sigma_\varepsilon^2}$$

(Recall that $\mathrm{ICC}(A, 1) = \mathrm{Cor}\left(Y_{ij}, Y_{ij'}\right)$ in two-way random-effects models.)


Fixed-rater agreement paradox (2)

- Rater effects $R_1, \ldots, R_K$ are the fixed, unknown constants, $\sum_{j=1}^{K} R_j = 0$.

	R_1	…	R_K
S_1	x		x
S_2	x		x
S_3	x		x
S_4	x		x
⋮		⋱	
S_n	x		x

Operational Score
$\bar{Y}_{i\cdot} = \mu + S_i + \frac{R_1 + \cdots + R_K}{K} + \frac{\varepsilon_1 + \cdots + \varepsilon_K}{K}$

$\frac{R_1 + \cdots + R_K}{K} = 0$



$$\mathrm{Cor}(\bar{Y}_{i\cdot}, \bar{Y}'_{i\cdot}) = \frac{\sigma_S^2}{\sigma_S^2 + \frac{\sigma_\varepsilon^2}{K}} = \mathrm{ICC}(C, K)$$

$\neq$

$$\mathrm{ICC}(\mathrm{A}, K) = \frac{\sigma_S^2}{\sigma_S^2 + \frac{\theta_R^2}{K} + \frac{\sigma_\varepsilon^2}{K}}$$

$\sigma^2_{\text{abs.error}}$?

Fixed-rater dispersion still matters

- Dispersion among fixed raters θ_R^2
  - Fixed raters can differ systematically in severity.
  - Under a variance-partitioning interpretation, the dispersion θ_R^2 is included in the denominator as **a source of disagreement.**
  - However, with rater identities given, fixed effects contribute no random variance.

How can these rater differences contribute to a correlation between replicate scores?

Fixed rater, random selection (1)

- K: the total number of fixed raters of interest
- k: the number of raters used for the average scores, $1 \leq k \leq K$
  - $k = 1$: single score, $k = K$: all-rater averages
- J: a random index selected uniformly from $\{1, \ldots, K\}$
  - $\Pr(J = j) = \frac{1}{K}$
- R_J: a random variable
  - R_j: a specific rater effect and an unknown constant
  - $\mathrm{E}[R_J] = 0$: finite-population mean
  - $\delta_R^2 = \mathrm{Var}(R_J) = \frac{1}{K}\sum_{j=1}^{K} R_j^2$: finite-population variance


(vs. $\theta_R^2 = \frac{1}{K-1}\sum_{j=1}^{K} R_j^2$: ANOVA-scaled dispersion)

Fixed rater, random selection (2)

- $\mathcal{J}_1, \mathcal{J}_2 \subseteq \{1, \ldots, K\}$: the rater index sets with size k for two replicates
- $M = |\mathcal{J}_1 \cap \mathcal{J}_2|$: the size of overlap raters between two replicates
  - m: the realized value, $\max\{0, 2k - K\} \leq m \leq k$
  - M could be a specific value m
  - or be a random variable, following a Hypergeometric(K, k, k)
- Average scores for two replicates
  - First: $\bar{Y}_{i\cdot} = \mu + S_i + \bar{R}_{\mathcal{J}_1} + \bar{\varepsilon}_{i\cdot}$; Second: $\bar{Y}'_{i\cdot} = \mu + S_i + \bar{R}_{\mathcal{J}_2} + \bar{\varepsilon}'_{i\cdot}$
  - $\mathrm{Var}\left(\bar{R}_{\mathcal{J}_1}\right) = \mathrm{Var}\left(\bar{R}_{\mathcal{J}_2}\right) = \frac{\delta_R^2}{k}\left(\frac{K-k}{K-1}\right)$
  - $\mathrm{Cov}\left(\bar{R}_{\mathcal{J}_1}, \bar{R}_{\mathcal{J}_2} \mid M = m\right) = \frac{\delta_R^2}{k^2}\left(\frac{mK-k^2}{K-1}\right)$
  - $\mathrm{Cov}\left(\bar{R}_{\mathcal{J}_1}, \bar{R}_{\mathcal{J}_2}\right) = 0$

Agreement ICC for single scores (1)

- **Single score $k = 1, K = 3$**



	R_1	R_2	R_3
$S^{(1)}$	x		
$S^{(2)}$		x	
$S^{(3)}$			x
$S^{(4)}$	x		
⋮			

Hypothetical replicate


	R_1	R_2	R_3
$S^{(1)}$			x
$S^{(2)}$	x		
$S^{(3)}$		x	
$S^{(4)}$		x	
⋮			

Between: Sampling without replacement
i.e., $J_1 \neq J_2$ or $M = 0$

$$\mathrm{ICC}^*(\mathrm{A}, k = 1 \mid M = 0) = \mathrm{Cor}(Y_{iJ_1}, Y_{iJ_2} \mid J_1 \neq J_2) = \frac{\sigma_s^2 - \delta_R^2/2}{\sigma_s^2 + \delta_R^2 + \sigma_\varepsilon^2}$$

Agreement ICC for single scores (2)

- **Single score $k = 1, K = 3$**



	R_1	R_2	R_3
$S^{(1)}$	x		
$S^{(2)}$		x	
$S^{(3)}$			x
$S^{(4)}$	x		
⋮			

Hypothetical replicate

	R_1	R_2	R_3
$S^{(1)}$		x	
$S^{(2)}$		x	
$S^{(3)}$	x		
$S^{(4)}$	x		
⋮			


Between: Sampling with replacement
i.e., M could be 0 or 1

vs. $\mathrm{ICC}(\mathrm{A},1) = \dfrac{\sigma_s^2}{\sigma_s^2 + \theta_R^2 + \sigma_\varepsilon^2}$

$$\mathrm{ICC}^*(A, k=1) = \mathrm{Cor}(Y_{iJ_1}, Y_{iJ_2}) = \frac{\sigma_s^2}{\sigma_s^2 + \delta_R^2 + \sigma_\varepsilon^2}$$

Agreement ICC for all-rater average scores

- **Average score $k = K = 3$**



	R_1	R_2	R_3
$S^{(1)}$	x	x	x
$S^{(2)}$	x	x	x
$S^{(3)}$	x	x	x
$S^{(4)}$	x	x	x
⋮			

Hypothetical replicate


	R_1	R_2	R_3
$S^{(1)}$	x	x	x
$S^{(2)}$	x	x	x
$S^{(3)}$	x	x	x
$S^{(4)}$	x	x	x
⋮			

Within: Sampling without replacement
Between: Sampling MUST with replacement. i.e., $M = 3$

$$\mathrm{ICC}^*(A, k = 3) = \mathrm{ICC}^*(A, k = 3 \mid M = 3) = \mathrm{Cor}(\bar{Y}_{i\cdot}, \bar{Y}'_{i\cdot}) = \frac{\sigma_s^2}{\sigma_s^2 + \sigma_\varepsilon^2/3}$$

Agreement ICC for partial-rater average scores

- **Average score $k = 2, K = 3$**



	R_1	R_2	R_3
$S^{(1)}$	x	x	
$S^{(2)}$	x	x	
$S^{(3)}$	x		x
$S^{(4)}$	x		x
⋮			

Hypothetical replicate

	R_1	R_2	R_3
$S^{(1)}$	x		x
$S^{(2)}$	x		x
$S^{(3)}$	x	x	
$S^{(4)}$	x	x	
⋮			

Within: Sampling without replacement
Between: Sampling with partial replacement. i.e., $M = 1$

$$\mathrm{ICC}^*(A, k = 2 \mid M = 1) = \frac{\sigma_s^2 - \delta_R^2/8}{\sigma_s^2 + \delta_R^2/4 + \sigma_\varepsilon^2/2}$$

The family of agreement ICCs

Distinct rater reliability

Under specific planning-missing designs

Common settings

replicates


$\bar{Y}_{i,\mathcal{J}_1} \leftrightarrow \bar{Y}'_{i,\mathcal{J}_2}$

Within replicates	Without replacement (Non-overlap, $m = 0$)	With partial replacement (Partial overlap, $1 \leq m \leq k$)	With replacement (Random overlap, M is RV)
Without replacement (k distinct rater)	$\mathrm{ICC}^*(A,k \mid M=0) = \frac{\sigma_S^2 - \frac{\delta_R^2}{K-1}}{\sigma_S^2 + \frac{\delta_R^2}{k}\left(\frac{K-k}{K-1}\right) + \frac{\sigma_\varepsilon^2}{k}}$	$\mathrm{ICC}^*(A,k \mid M=m) = \frac{\sigma_S^2 + \frac{\delta_R^2}{k^2}\left(\frac{mK-k^2}{K-1}\right)}{\sigma_S^2 + \frac{\delta_R^2}{k}\left(\frac{K-k}{K-1}\right) + \frac{\sigma_\varepsilon^2}{k}}$	$\mathrm{ICC}^*(A,k) = \frac{\sigma_S^2}{\sigma_S^2 + \frac{\delta_R^2}{k}\left(\frac{K-k}{K-1}\right) + \frac{\sigma_\varepsilon^2}{k}}$
With replacement (k iid sampling)	—	—	$\mathrm{ICC}^*_{\mathrm{iid}}(A,k) = \frac{\sigma_S^2}{\sigma_S^2 + \frac{\delta_R^2}{k} + \frac{\sigma_\varepsilon^2}{k}}$

Structurally identical to conventional ICC(A, k)

Link to conventional agreement ICC

- **Average score $k = K = 3$**



	R_1	R_2	R_3
$S^{(1)}$	xx	x	
$S^{(2)}$		xxx	
$S^{(3)}$		x	xx
$S^{(4)}$	xx		x
⋮			

Hypothetical replicate

	R_1	R_2	R_3
$S^{(1)}$	x	x	x
$S^{(2)}$	x		xx
$S^{(3)}$	xx	x	
$S^{(4)}$		x	xx
⋮			


Within: Sampling with replacement
Between: Sampling with replacement. i.e., M is a RV


$$\mathrm{ICC}^{*}_{\mathrm{iid}}(A,\mathrm{K}) = \frac{\sigma_S^2}{\sigma_S^2 + \frac{\delta_R^2}{K} + \frac{\sigma_\varepsilon^2}{K}}$$

vs. $$\mathrm{ICC}(\mathrm{A},\mathrm{K}) = \frac{\sigma_s^2}{\sigma_s^2 + \frac{\theta_R^2}{K} + \frac{\sigma_\varepsilon^2}{K}}$$

Flowchart for selecting an ICC

McGraw and Wong (1996)




Scope
1. Within replicates
2. Target score unit (k)
3. Between replicates
Selected coefficient
Independent draws with replacement
Between-replicate draws inde-
pendent by convention; specify k
IID rater benchmark
$\mathrm{ICC}^{*}_{\mathrm{iid}}(A, k)$
$k = K$
Full-panel average
Complete overlap
$m = k = K$ is mandatory
Full-panel composite
$\mathrm{ICC}^{*}(A, K) = \mathrm{ICC}(3, K)$
Two-way
mixed-effect
Fixed K rater
Absolute agre
ment target
er test–retest
1 | $M = 1$)
ter agreement
$_{\mathrm{iff}}(A, 1)$
t reassignment
$_{\mathrm{nd}}(A, 1)$
nt panels
k | $M = 0$)
1. Two-way mixed-effects design & Agreement
2. Number of the fixed rater: K
3. Unit of the operational score: k
4. Replacement mechanism
• Within: without / with
• Between: without / partial / with
$1 < k < K$
Partial-panel
average
Partial panel overlap
$0 < m < k$;
feasible fixed m
Partially retained panels
$\mathrm{ICC}^{*}(A, k \mid M = m)$
Retain assigned panel
$m = k$
Fully retained panel
$\mathrm{ICC}^{*}(A, k \mid M = k)$
Independent
panel assignment
Random overlap M
Independent panels
$\mathrm{ICC}^{*}(A, k)$

Ex 1: Single ratings from 3 fixed raters

- **Math achievement measures.**

  $n = 161$ (Delacruz, 2011. pp. 33, 127)

  The three open-ended items on the math assessment were scored using three raters who were trained on how to apply the rubrics…….
  Each rater used the final version of the rubric to score responses from **20 students**.
  Interrater reliability for each item was determined by calculating ICC for each item, which were applied to single measurement scores (i.e., individual ratings of items by individual judges)……
  A two-way, mixed-effect model was used to examine the absolute agreement of measurements between the three raters for each of the three open-ended items.

1. Care about interrater agreement
2. Fixed raters, K=3
3. Single rating, k = 1
4. Replication mechanism?

Different raters for two replicates

$$\mathrm{ICC}^*(A, 1 \mid M = 0)$$

Independent reassignment

$$\mathrm{ICC}^*(A, 1)$$

Ex 2: All-rater average ratings

- **SBS knowledge. $n = 118$**

  (Verhees et al., 2019. pp. 201, 203).

  Children are instructed to tell a story using these prompt words as if the story is about themselves. … Transcribed are scored in terms of SBS knowledge on a scale from 1 to 7. ……
  For the current study, stories were double coded by two trained coders ……
  Interrater agreement was established by separately rating the same ASA stories for **30 participants** from the current sample. … using the two-way mixed model and absolute agreement for average measures. After establishing sufficient interrater reliability both coders …, all stories were thus double coded. … the mean score of the two raters was used for further analyses.

1. Care about interrater agreement
2. Fixed raters, K=2
3. Average rating, k = 2
4. Replication mechanism
   - Within: without
   - Between: with (completed overlap)

Full-panel average

$$\mathrm{ICC}^*(A, k = 2 \mid M = 2) = \mathrm{ICC}^*(A, k = 2)$$

Ex 3: Partial-rater average ratings

- **Attachment script assessment.**
  $n = 149$. (Verhees et al., 2019. PP. 206).

  Stories were double coded by four trained coders who were blind to participants' scores on the other measures. The **same 30 ASA stories** were independently rated by all four coders to establish interrater agreement.

  ……

  After establishing interrater agreement, two coders separately rated one half of the remaining stories, and the other two coders separately rated the other half, so that all stories were double coded.

1. Care about interrater agreement
2. Fixed raters, K=4
3. Average rating, k = 2
4. Replication mechanism
   - Within: without
   - Between: without? Non-overlap

	R_1	R_2	R_3	R_4
S_1	x	x		
⋮	x	x		
S_{71}			x	x
⋮			x	x

	R_1	R_2	R_3	R_4
S_1			x	x
⋮			x	x
S_{71}	x	x		
⋮	x	x		

Disjoint pair on replication

$$\mathrm{ICC}^*(A, k = 2 \mid M = 0)$$

Discussion

- Main contributions
  - Correlation-based definition for ICC
  - Fixed effects and random assignment
  - A family of agreement ICCs for two-way mixed-effects designs
  - To select a proper agreement ICC, researchers should specify the unit of the score, within- and between-replicates sampling mechanism.

- Limitations and Future Work
  - Depend on the model assumptions
  - Estimation properties
  - Practical selection and reporting guidelines
  - Extensions beyond uniform assignment