

1. TOPOLOGICAL TIME SERIES ANALYSIS (TTSA)

1.1. Persistent Cup Products for QuasiPeriodicity Detection. A function $f : \mathbb{R} \rightarrow \mathbb{C}$ is said to be quasiperiodic if it can be written as $f(t) = F(t\omega_1, \dots, t\omega_N)$ for a continuous function $F : \mathbb{T}^N \rightarrow \mathbb{C}$ on the N -torus $\mathbb{T}^N = S^1 \times \dots \times S^1$, and \mathbb{Q} -linearly independent real numbers $\omega_1, \dots, \omega_N$. For these types of functions and appropriate parameters $d \in \mathbb{N}$ and $\tau > 0$, the sliding window point cloud

$$\mathbb{SW}_{d,\tau}f = \{SW_{d,\tau}f(t) : t \in \mathbb{R}\} \quad , \quad SW_{d,\tau}f(t) = \begin{bmatrix} f(t) \\ f(t+\tau) \\ \vdots \\ f(t+d\tau) \end{bmatrix}$$

can be shown to be dense in an N -torus embedded in \mathbb{C}^{d+1} .

Let $\mathbf{dgm}_j^{\mathcal{R}}(\mathbb{SW}_{d,\tau}f)$ denote the persistence diagram for the j -th persistent cohomology over a field \mathbb{F} , $j \geq 0$, of the Rips filtration on the Sliding window point cloud $\mathbb{SW}_{d,\tau}f$. In TTSA the diagrams $\mathbf{dgm}_j^{\mathcal{R}}(\mathbb{SW}_{d,\tau}f)$ are used to certify that f is quasiperiodic (i.e., that the closure of $\mathbb{SW}_{d,\tau}f$ is an N -torus) – by checking that there are $\binom{N}{j}$ highly persistent features. The main issue with this approach is that even if the dimension of the cohomology vector spaces looks right, one will not have an N -torus unless the cup product structure is also correct. The goal of this project is to bridge this gap; the challenge is to define and implement a quasiperiodicity score involving persistent cup products.

A place to start:

- (1) Confirm the advertised cup product structure on \mathbb{T}^2 . Use a small triangulation.

References.

- (1) Ripser: Efficient Computation of Vietoris-Rips Persistence Barcodes, U. Bauer,
<https://arxiv.org/pdf/1908.02518.pdf>
- (2) Dualities in Persistent (Co)Homology, V. de Silva, D. Morozov, M. Vejdemo Johanson,
<https://arxiv.org/pdf/1107.5665.pdf>
- (3) Sliding Window Persistence of QuasiPeriodic Functions, H. Gakhar and J. A. Perea,
<https://arxiv.org/pdf/2103.04540.pdf>
- (4) (Quasi)Periodicity Quantification in Video Data, Using Topology, by Christopher J. Tralie and Jose A. Perea,
<https://arxiv.org/pdf/1704.08382.pdf>
- (5) Topological Time Series Analysis, J. A. Perea,
<https://www.ams.org/journals/notices/201905/rnoti-p686.pdf>
- (6) Persistent Cup-Length,
<https://arxiv.org/pdf/2107.01553.pdf>