

Constant additive and multiplicative seasonal figures

PD Dr. Ralf Stecking and **Abigail Opokua Asare**

Department of Business Administration,
Economics and Law


Institute of Economics
Carl von Ossietzky University Oldenburg

11 November, 2025


Overview

1. Phase average method
2. Seasonal dummies
3. Regression with dummy variables
4. Empirical example: Retail sales
5. Holt's Winter seasonal method
6. Forecasting error measures

Retail sales per Month (1950-2019)



Retail sales per quarter (1994-2010)



Constant additive seasonal figures

- ▶ **Assumption 1:** When there are stable periodic fluctuations for the whole observation period.
- ▶ **Assumption 2:** The time series can be represented by an additive model:

$$X = T + C + S + U$$

- ▶ **Assumption 3:** Seasonal influence increases or lowers the observation value by a constant factor.

$$X = T + C + S + U$$

- ▶ **Example:** In case of quarterly data we find four constant values

$$S_1, S_2, S_3, S_4!$$

Method I

In order to “free” the time series $X - G$ from its residual component U we introduce the **phase average method**:

- 1 Identification of the smooth component $G = T + C$ as
 - ▶ linear function of time,
 - ▶ polynomial of higher degree,
 - ▶ exponential function or
 - ▶ by use of moving average methods.
- 2 The difference $X - G = S + U$ just includes the seasonal component S and the residual component U .

Method II

- 3 For each phase (e.g. month, quarter) the mean deviation D_{ph} from the smooth component is computed: \rightarrow the residual component U is “averaged to zero”.
- 4 In case of **quarterly data** one gets four values D_1, D_2, D_3, D_4 .
- If these four values do not add up to zero, they will be adjusted by its mean value $\bar{d} = \frac{1}{4} \sum D_{ph}$ which lead to seasonal index numbers


$$S_{ph} := D_{ph} - \bar{d},$$

that add up to zero: $S_1 + S_2 + S_3 + S_4 = 0$

- A **seasonally adjusted time series** is obtained by subtracting the seasonal index number from the non adjusted time series!

Example

Retail sales per quarter (2006-2010)



Calculation of additive seasonal index numbers – Retail Sales

Year	Quarter	x	G	x-G
2006	1	251.9	NA	NA
2006	2	268.0	NA	NA
2006	3	262.7	269.4250	-6.7250
2006	4	295.9	268.5125	27.3875
2007	1	250.3	267.2375	-16.9375
2007	2	262.3	266.0625	-3.7625
2007	3	258.2	266.0375	-7.8375
2007	4	291.0	267.4500	23.5500
2008	1	255.0	269.4750	-14.4750
2008	2	268.9	271.0750	-2.1750
2008	3	267.8	270.0375	-2.2375
2008	4	294.2	267.4625	26.7375
2009	1	243.5	264.8125	-21.3125
2009	2	259.8	262.3000	-2.5000
2009	3	255.7	261.8875	-6.1875
2009	4	286.2	262.9750	23.2250
2010	1	248.2	264.7125	-16.5125
2010	2	263.8	266.9375	-3.1375
2010	3	265.6	NA	NA
2010	4	294.1	NA	NA

Quarter	Mean	adjusted	N
1	-17.309375	-17.128125	4
2	-2.893750	-2.712500	4
3	-5.746875	-5.565625	4
4	25.225000	25.406250	4
Total	-0.725	0	16

Regression analysis

- ▶ Calculation of seasonal index numbers by means of regression analysis.

Regression with **dummy variables**.

- ▶ Dummy variables are auxiliary variables with two outcomes (e.g., 0 and 1) for measuring certain qualitative impacts → **binary** or **dichotomous** variables.
- ▶ **Four** quarters of a year can be represented by four seasonal dummies.

Calculation of additive seasonal index numbers

t	1	2	3	4	...	68
x_t	x_1	x_2	x_3	x_4	...	x_{68}
constant	1	1	1	1	...	1
$Dummy_1$	1	0	0	0	...	0
$Dummy_2$	0	1	0	0	...	0
$Dummy_3$	0	0	1	0	...	0
$Dummy_4$	0	0	0	1	...	1

Linear regression approach II

- ▶ By using the general approach with constant b_0 just **three** seasonal dummies are adequate:

$$x_t - g_t = b_0 + b_1 \cdot \text{Dummy}_1 + b_2 \cdot \text{Dummy}_2 + b_3 \cdot \text{Dummy}_3 + u_t$$

with

$$D_1 = b_0$$


$$D_2 = b_0 + b_1$$

$$D_3 = b_0 + b_2$$

$$D_4 = b_0 + b_3$$

- ▶ The seasonal index numbers are obtained by adjusting the phase averages D_{ph} for their means.
- ▶ The residuals of the regression equation are representing the residual component U . The time series $x_t^{SA} = g_t + u_t$ is referred to as **seasonally adjusted**.

Retail sales per quarter (1994-2010)



Linear regression

	WITHOUT constant term	WITH constant term
Constant		244.682*** (1.271)
quarter1	244.682*** (1.271)	
quarter2	256.241*** (2.006)	11.559*** (2.375)
quarter3	253.700*** (1.868)	9.018*** (2.259)
quarter4	288.765*** (1.035)	44.082*** (1.639)
Num.Obs.	68	68
R2	0.999	0.872
R2 Adj.	0.999	0.866
AIC	455.2	455.2
BIC	466.3	466.3
RMSE	6.39	6.39

Empirical example: seasonal index numbers

- ▶ The regression equation reads:

$$x_t - g_t = b_0 + b_1 \cdot \text{Dummy}_1 + b_2 \cdot \text{Dummy}_2 + b_3 \cdot \text{Dummy}_3 + u_t$$

with

$$D_1 = b_0 = 244.682$$

$$D_2 = b_0 + b_1 = 244.682 + 11.559 = 256.241$$

$$D_3 = b_0 + b_2 = 244.682 + 9.018 = 253.7$$




$$D_4 = b_0 + b_3 = 244.682 + 44.082 = 288.764$$

- ▶ The seasonal index numbers are obtained by adjusting the phase averages D_{ph} for their means, i.e.:

$$\bar{d} = \frac{1}{4} \sum D_{ph} = \frac{1}{4}(244.682 + \dots + 253.7) = 260.8468$$

and therefore $S_1 = D_1 - \bar{d} = -16.1648$; $S_2 = -4.6068$; $S_3 = -7.1468$; $S_4 = 27.9172$

Retail sales / seasonal component



Regression with trend component

- ▶ The real advantage of the regression method is, that it can be used to determine seasonal **and** trend component simultaneously.
- ▶ Therefore seasonal dummies and trend function are combined in the regression equation.
- ▶ The approach for the **linear** trend now reads:

$$x_t = b_0 + b_1 \cdot Dummy_1 + b_2 \cdot Dummy_2 + b_3 \cdot Dummy_3 + b_4 \cdot t + u_t$$




- ▶ for the **linear** and **quadratic** trend:

$$x_t = b_0 + b_1 \cdot Dummy_1 + b_2 \cdot Dummy_2 + b_3 \cdot Dummy_3 + b_4 \cdot t + b_5 \cdot t^2 + u_t$$


Regression with linear and quadratic trend component

	linear trend	quadratic trend
Constant	236.332*** (1.156)	235.045*** (1.653)
trend	0.253*** (0.027)	0.364*** (0.104)
$trend^2$		-0.002 (0.002)
quarter2	11.306*** (1.510)	11.303*** (1.500)
quarter3	8.512*** (1.374)	8.508*** (1.392)
quarter4	43.323*** (1.352)	43.323*** (1.351)
Num.Obs.	68	68
R2	0.949	0.950
R2 Adj.	0.946	0.946
AIC	394.4	395.1
BIC	407.7	410.7
RMSE	4.03	3.99


seasonal + linear trend component




seasonal + linear and quadratic trend component




Seasonal + linear/Quadratic trend component



Residual Component




Retail sales per Month (1950-1990)




How to visually inspect

Retail Sales — Additive Seasonality (Constant Amplitude)



Retail Sales — Multiplicative Seasonality (Expanding Amplitude)



Constant multiplicative seasonal figures

- ▶ **Assumption 1:** When there is a proportional or percentage increase in periodic fluctuations for the whole observation period.
- ▶ **Assumption 2:** The time series can be represented by a multiplicative model: model:

$$X = T \cdot C \cdot S \cdot U$$

- ▶ **Assumption 3:** Seasonal influence increases or lowers the observation value by a proportional or percentage factor.

$$X = T + C + S + U$$

- ▶ **Example** In case of quarterly data we find four constant values

$$S_1, S_2, S_3, S_4!$$

Phase average method I

In order to “free” the time series $\frac{X}{G}$ from its residual component U we introduce the **phase average method**:

- 1 After identification of the smooth component G as

$$\frac{X}{G} = S \cdot U$$

is used, including just the seasonal and the residual component.

- 2 The quotients

$$\frac{x_1}{g_1}, \frac{x_2}{g_2}, \dots, \frac{x_T}{g_T}$$

show the observation values in relation to their smooth component. A quotient of 1.08 means, that the current value exceeds the smooth component by 8%.

Phase average method II

- 3 For each phase the average quotient Q_{ph} is computed as geometrical mean of the quotients $\frac{X}{G}$ from the same phases. In practice: often arithmetic mean or median is used.
 - 4 In case of **quarterly data** one gets four values Q_1, Q_2, Q_3, Q_4 .
- The mean of the seasonal factors must be equal to one (the neutral element of multiplication). Adjusting by the mean value $\bar{q} = \sqrt[4]{\prod Q_{ph}}$ leads to seasonal factors

$$S_{ph} := \frac{Q_{ph}}{\bar{q}}$$

with a geometrical mean of one.

- A **seasonally adjusted time series** is obtained by dividing the non adjusted time series by the seasonal index number!

Calculation of multiplicative seasonal index numbers

Year	Quarter	x	G	x/G
2006	1	251.9	NA	NA
2006	2	268.0	NA	NA
2006	3	262.7	269.4250	0.9750
2006	4	295.9	268.5125	1.1020
2007	1	250.3	267.2375	0.9366
2007	2	262.3	266.0625	0.9859
2007	3	258.2	266.0375	0.9705
2007	4	291.0	267.4500	1.0881
2008	1	255.0	269.4750	0.9463
2008	2	268.9	271.0750	0.9920
2008	3	267.8	270.0375	0.9917
2008	4	294.2	267.4625	1.1000
2009	1	243.5	264.8125	0.9195
2009	2	259.8	262.3000	0.9905
2009	3	255.7	261.8875	0.9764
2009	4	286.2	262.9750	1.0883
2010	1	248.2	264.7125	0.9376
2010	2	263.8	266.9375	0.9882
2010	3	265.6	NA	NA
2010	4	294.1	NA	NA

Mean quotient X/G per quarter

Quarter	Geometric Mean	Adjusted	N
1	0.935	0.936	4
2	0.989	0.990	4
3	0.978	0.979	4
4	1.09	1.10	4
Total	0.9992872	1	16

Holt-Winters with Additive Seasonal Fluctuations

- ▶ It is suitable for time series with trend but with additive seasonality.

$$F_{t+h} = \ell_t + b_t h + s_{t+h-m}$$

- ▶ three updating equations:

$$\ell_t = \alpha(x_t - s_{t-m}) + (1 - \alpha)(\ell_{t-1} + b_{t-1})$$

$$b_t = \beta(\ell_t - \ell_{t-1}) + (1 - \beta)b_{t-1}$$

$$s_t = \gamma(x_t - \ell_{t-1} - b_{t-1}) + (1 - \gamma)s_{t-m}$$

- ▶ m which indicates seasonal indices used for forecasting is from the final year of the sample.

Holt-Winters with Additive Seasonal Fluctuations

- ▶ m is the length of the seasonality.
- ▶ We assume seasonality with m to be the number of the seasons in the first half of the sample.
- ▶ α is the smoothing constant, β is the slope parameter and γ is the parameter for adjusting seasonal fluctuations.
- ▶ We need m start values for the seasonal factors.
- ▶ We also need start values for g_t and b_t .

Holt-Winters with Multiplicative Seasonal Fluctuations

- ▶ It is suitable for time series with trend but with multiplicative seasonality.

$$F_{t+h} = (g_t + b_th)s_{t+h-m}$$

- ▶ Updating equations:

$$\ell_t = \alpha \frac{x_t}{s_{t-m}} + (1 - \alpha)(\ell_{t-1} + b_{t-1})$$

$$b_t = \beta(\ell_t - \ell_{t-1}) + (1 - \beta)b_{t-1}$$

$$s_t = \gamma \frac{x_t}{(\ell_{t-1} + b_{t-1})} + (1 - \gamma)s_{t-m}$$

- ▶ In multiplicative seasonality, values in different seasons differ by percentage amount

Evaluating Forecasting Model Performance

Forecasters are usually concerned about the accuracy of their forecast.

- ▶ Measuring the forecasting error serves several purposes:
 - as a **target criterion** for model and parameter estimation,
 - in order to assess **goodness of fit** and **adequacy** of the model,
 - in order to **compare** different forecasting approaches.

Forecasting Error

1. Forecasting error:

$$e_t = X_t - \hat{X}_{t|t-1}$$

2. Absolute error:

$$AE_t = |e_t|$$

3. Percentage error:

$$PE_t = \frac{e_t}{X_t} \cdot 100$$

4. Absolute percentage error:

$$APE_t = \left| \frac{e_t}{X_t} \right| \cdot 100$$

Accuracy Measure I

1. Mean error:

$$ME = \frac{1}{T} \sum_{t=1}^T e_t$$

2. Mean absolute error:

$$MAE/MAD = \frac{1}{T} \sum_{t=1}^T |e_t|$$

3. Mean square error:

$$MSE = \frac{1}{T} \sum_{t=1}^T e_t^2$$

Accuracy Measure II

1. Mean error in percent:

$$MPE = \frac{1}{T} \sum_{t=1}^T \frac{e_t}{X_t} \cdot 100$$

2. Mean absolute error in percent:

$$MAPE = \frac{1}{T} \sum_{t=1}^T \left| \frac{e_t}{X_t} \right| \cdot 100$$

3. Root mean square error::

$$RMSE = \sqrt{MSE} = \sqrt{\frac{1}{T} \sum_{t=1}^T e_t^2}$$