



Presentation:

Algin[®] Tablet: Each film-coated tablet contains 50 mg Tiemonium Methylsulphate

Algin® Injection: Each 2 ml injection contains 5 mg Tiemonium Methylsulphate

Indications:

Algin® (Tiemonium Methylsulphate) is an antispasmodic agent and is used in the relief of visceral pain. It reduces muscle spasm of the intestine, biliary system, bladder and uterus. Algin® (Tiemonium Methylsulphate) is indicated for pain in gastrointestinal and biliary diseases e.g. hepato-colic, colitis, gastric and duodenal ulcer. Algin® can also be used in gynecology and renal colic.

Algin®(Tiemonium Methylsulphate) is also indicated for acute pain.

Dosage and Administration:

Adults:

Oral: The usual dosage is 100 to 300 mg (2 to 6 tablets) per day in divided doses.

Parenteral: The usual dosage is 5 mg (1 ampoule) three times daily by intramuscular

or slow intravenous injection.

Contra-indication:

Tiemonium Methylsulphate is contraindicated in Hyperthyroidism, Glaucoma, Prostatic hypertrophy, Paralytic ileus, Pyloric stenosis, Tachycardia, Acute edema of the lung and Myocardial infarction.

Precautions:

Tiemonium Methylsulphate should be used with caution in patients suffering from chronic bronchitis, coronary insufficiency, ambient hyperthermia, severe hepatic insufficiency, severe renal insufficiency, pregnancy and in breast-feeding. To avoid the risk of hypotension and tachycardia, the injection should be administered slowly.

Storage:

Store in a cool, dry place and away from light & children.

How Supplied:

Algin® Tablet: Box containing 5×10's film coated tablets in blister pack.

Algin® Injection: Box containing 1×5's ml ampoules in blister pack.



C-Code No.: 5989/V 02

Size: 60 x 120mm